

ON THE REFLEXIVITY OF PAIRS OF CONTRACTIONS

MAREK KOSIEK, ALFREDO OCTAVIO, AND MAREK PTAK

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. We consider pairs of commuting contractions such that the joint left essential spectrum is dominating for the algebra $H^\infty(\mathbf{D}^2)$. It is also assumed, in the first case, that one of them is C_0 , and the second one is absolutely continuous. In the second case, we assume that the pair is diagonally extendable. It will be shown that such pairs are reflexive.

1. INTRODUCTION

The dual algebras technique, after a great achievement in the problem of invariant subspaces for single operators, is being applied to pairs or N -tuples of operators. Theorems 5.3 and 5.4 of [KP] show the reflexivity for pairs of commuting C_0 - or doubly commuting contractions with dominating joint left essential spectrum. The purpose of our paper is to weaken those assumptions. For the first result we need the C_0 - assumption for one operator only. For the second one doubly commutativity is replaced by a weaker one, diagonal extendibility (for a definition see section 4).

Throughout the paper $L(\mathcal{H})$ denotes the algebra of all linear, bounded operators in a complex separable, infinite-dimensional Hilbert space \mathcal{H} . $I_{\mathcal{H}}$ or I stands for the identity in \mathcal{H} . By a subspace of \mathcal{H} , we mean a closed subspace, and by an algebra of operators - a subalgebra of $L(\mathcal{H})$ with unit $I_{\mathcal{H}}$. If \mathcal{S} is a subset of $L(\mathcal{H})$, then $\mathcal{W}(\mathcal{S})$, $\mathcal{A}(\mathcal{S})$, and $Lat \mathcal{S}$ stand for the WOT (=weak operator topology)-closed algebra, the weak-star closed algebra, generated by \mathcal{S} and I , and the lattice of all invariant subspaces for \mathcal{S} , respectively. $Alg Lat \mathcal{S}$ is the algebra of all operators on \mathcal{H} , which leave invariant all subspaces from $Lat \mathcal{S}$. An algebra \mathcal{W} is called *reflexive* if $\mathcal{W} = Alg Lat \mathcal{W}$. A family $\mathcal{S} \subset L(\mathcal{H})$ is called *reflexive* if so is $\mathcal{W}(\mathcal{S})$.

As usual, $A(\mathbf{D}^N)$ will denote the polydisc algebra, i.e., the algebra of all functions analytic on \mathbf{D}^N and continuous on its closure, and $H^\infty(\mathbf{D}^N)$ the algebra of all bounded analytic functions on \mathbf{D}^N . A subset E of \mathbf{D}^N is called *dominating for $H^\infty(\mathbf{D}^N)$* if $\sup\{|h(z)| : z \in \mathbf{D}^N\} = \sup\{|h(z)| : z \in E\}$ for all $h \in H^\infty(\mathbf{D}^N)$. For a complex function f on $F \subset \mathbf{D}^N$ denote by f^\sim the function $f^\sim(z) = f(\bar{z})$, $z \in F$.

Received by the editors April 1, 1993 and, in revised form, July 30, 1993.

1991 *Mathematics Subject Classification.* Primary 47A15; Secondary 47D27.

Key words and phrases. Essential spectrum, reflexive algebras, reflexive operators.

The first and the third author were partially supported by KBN grant.

Let T_1, \dots, T_N be commuting operators on \mathcal{H} . Denote by $C(\mathcal{H})$ the Calkin algebra and by π the quotient map $\pi : L(\mathcal{H}) \rightarrow C(\mathcal{H})$. Recall that the *joint left essential spectrum* $\sigma_{le}(T_1, \dots, T_N)$ of T_1, \dots, T_N is defined as the joint left spectrum of $\pi(T_1), \dots, \pi(T_N)$. Let us recall (see [KP, Lemma 2.1]) that $\lambda = (\lambda_1, \dots, \lambda_N) \in \sigma_{le}(T_1, \dots, T_N)$, if and only if there exists an orthonormal sequence $\{x_n\}$ such that $\lim_{n \rightarrow \infty} \|(T_i - \lambda_i)x_n\| = 0$, for $i = 1, \dots, N$. Moreover, using standard techniques we can prove

Lemma 1.1. *Let T_1, \dots, T_N be commuting operators on \mathcal{H} ; then $\lambda = (\lambda_1, \dots, \lambda_N) \in \sigma_{le}(T_1, \dots, T_N)$, if and only if there exists a sequence $\{x_n\}$ such that $x_n \rightarrow 0$ weakly, $\|x_n\| = 1$, and $\lim_{n \rightarrow \infty} \|(T_i - \lambda_i)x_n\| = 0$, for $i = 1, \dots, N$.*

A contraction $T \in L(\mathcal{H})$ is *absolutely continuous* (a.c.) if it has no singular unitary part.

2. REPRESENTATIONS

Let us recall that the algebra homomorphism $\Phi : A(\mathbf{D}^2) \rightarrow L(\mathcal{H})$ is a *representation* if $\|\Phi(f)\| \leq \|f\|$ for $f \in A(\mathbf{D}^2)$. It is a consequence of standard techniques that for every $x, y \in \mathcal{H}$ there exists a complex, Borel, regular measure $\mu_{x,y}$ on $\overline{\mathbf{D}^2}$ such that $(\Phi(u)x, y) = \int u d\mu_{x,y}$ for $x, y \in \mathcal{H}, u \in A(\mathbf{D}^2)$. We say that Φ is *absolutely continuous* (a.c.), if it has a system of *elementary measures* $\{\mu_{x,y}\}_{x,y \in \mathcal{H}}$ such that each element of the system is absolutely continuous with respect to some (positive) representing measure $\nu_z, z \in \mathbf{D}^2$. By [G, VI.1.2, II.7.5], the above definition is equivalent to that given in [KP, Chapters 3 and 4], which uses the terminology of bands of measures.

Having the pair $\{T_1, T_2\}$ of commuting contractions in $L(\mathcal{H})$, let us construct, using Ando Theorem [SNF, I.6.4] like in [KP, Chapter 5], the representation Φ of the algebra $A(\mathbf{D}^2)$ generated by $\{T_1, T_2\}$, i.e., $\Phi(z_i) = T_i, i = 1, 2$. We say that the pair $\{T_1, T_2\}$ is a.c. if Φ is a.c. In the same way we can construct the representation Φ^+ of $A(\mathbf{D}^2)$ generated by $\{T_1^*, T_2^*\}$. Moreover we have

Lemma 2.1. (1) *The representation Φ is a.c. if and only if Φ^+ is a.c.*

(2) *If $\{T_1, T_2\}$ is an a.c. pair of commuting contractions and $f \in H^\infty(\mathbf{D}^2)$ then for any vectors x, y we have*

$$(f(T_1, T_2)y, x) = (y, f^\sim(T_1^*, T_2^*)x).$$

Proof. Let $u \in A(\mathbf{D}^2)$, and $\mu_{y,x}$ be an elementary measure of Φ for $y, x \in \mathcal{H}$ absolutely continuous with respect to a representing measure ν_z for some $z \in \mathbf{D}^2$. It is obvious that $u^\sim \in A(\mathbf{D}^2)$.

For a complex measure μ on $\overline{\mathbf{D}^2}$ denote by μ^\sim the Borel measure $\mu^\sim(\cdot) = \overline{\mu(\overline{\Pi(\cdot)})}$, where $\Pi : z \rightarrow \bar{z}$ is a homeomorphism of $\overline{\mathbf{D}^2}$ onto itself. Then for $u \in A(\mathbf{D}^2)$ we have

$$\begin{aligned} (u(T_1^*, T_2^*)x, y) &= (x, u(T_1^*, T_2^*)^*y) = (x, u^\sim(T_1, T_2)y) \\ &= \overline{(u^\sim(T_1, T_2)y, x)} = \int u^\sim d\mu_{y,x} \\ &= \int \overline{u^\sim} d\overline{\mu_{y,x}} = \int u d\mu_{y,x}^\sim. \end{aligned}$$

So $\eta_{x,y} \stackrel{\text{df}}{=} \mu_{y,x}^\sim$ is an elementary measure of Φ^+ for vectors $x, y \in \mathcal{H}$. Since $\mu_{y,x}$ is absolutely continuous with respect to ν_z , then $\eta_{x,y}$ is absolutely continuous with respect to ν_z^\sim , and an easy calculation shows that ν_z^\sim is a representing measure for $\bar{z} \in \mathbf{D}^2$, which finishes the proof of (1).

Next, for (2), one can easily see that $f^\sim \in H^\infty(\mathbf{D}^2)$. Let, like in (1), $\mu_{y,x}$ and $\mu_{y,x}^\sim$ be elementary measures of Φ and Φ^+ , respectively. Notice that $\sigma_z \stackrel{\text{df}}{=} \nu_z + \nu_z^\sim$ is positive and symmetric with respect to adjoint. Let $d\mu_{y,x} = h_{y,x} d\sigma_z$. Then $d\mu_{y,x}^\sim = h_{y,x}^\sim d\sigma_z$. Hence, the following finishes the proof

$$\begin{aligned} (f(T_1, T_2)y, x) &= \int f d\mu_{y,x} = \int f(\lambda)h_{y,x}(\lambda) d\sigma_z(\lambda) \\ &= \int f(\bar{\lambda})h_{y,x}(\bar{\lambda}) d\sigma_z(\bar{\lambda}) = \overline{\int f^\sim(\lambda)h_{y,x}^\sim(\lambda) d\sigma_z(\lambda)} \\ &= \overline{\int f^\sim d\mu_{y,x}^\sim} = \overline{(f^\sim(T_1^*, T_2^*)x, y)} = (y, f^\sim(T_1^*, T_2^*)x). \end{aligned}$$

Having the representation constructed above, using [K1, Proposition] (see also [K1, Sec.2], [K2, Sec.3]), we can decompose Φ and \mathcal{H} into orthogonal sums as follows

$$(2.1) \quad \Phi = \Phi_0 \oplus \Phi_1 \oplus \Phi_2 \oplus \Phi_3, \quad \mathcal{H} = \mathcal{H}_0 \oplus \mathcal{H}_1 \oplus \mathcal{H}_2 \oplus \mathcal{H}_3,$$

where Φ_i ($i = 0, 1, 2, 3$) is the restriction of Φ to the subspace \mathcal{H}_i reducing all the values of Φ . Moreover we get

- Φ_0 is an absolutely continuous representation,
- $T_i|_{\mathcal{H}_i}$ ($i = 1, 2$) is a unitary operator with singular spectral measure,
- $T_i|_{\mathcal{H}_3}$ ($i = 1, 2$) are unitary operators.

So we get

Lemma 2.2. *If T_1 is c.n.u. and T_2 is a.c. then the pair $\{T_1, T_2\}$ is a.c.*

3. THE FIRST REFLEXIVITY RESULT

The purpose of the section is the following

Theorem 3.1. *Let $\{T_1, T_2\} \subset L(\mathcal{H})$ be a pair of commuting contractions. Assume also that $T_1 \in \mathbf{C}_0$ and T_2 is absolutely continuous. If $\sigma_{1e}(T_1, T_2) \cap \mathbf{D}^2$ is dominating for $H^\infty(\mathbf{D}^2)$, then the algebra $\mathcal{W}(T_1, T_2)$ is reflexive.*

The proof of the theorem uses the technique of dual algebras. Thus we start with some definitions and notation. The Banach space of trace class operators in $L(\mathcal{H})$ with the trace norm will be denoted by (τc) . Recall that the bilinear form $\langle A, T \rangle = \text{tr}(AT)$, $A \in L(\mathcal{H})$, $T \in (\tau c)$, on $L(\mathcal{H}) \times (\tau c)$ allows us to identify $L(\mathcal{H})$ with $(\tau c)^*$ and the weak-star topology on $L(\mathcal{H})$ under this identification coincides with the ultraweak operator topology. If \mathcal{A} is any ultraweakly closed subspace of $L(\mathcal{H})$, then its preannihilator ${}^\perp\mathcal{A}$ is a closed subspace of (τc) . \mathcal{A} can be identified with the dual of $Q_{\mathcal{A}} = (\tau c) / {}^\perp\mathcal{A}$ via the bilinear form $\langle A, T \rangle = \text{tr}(AT)$ on $\mathcal{A} \times Q_{\mathcal{A}}$. The quotient norm in $Q_{\mathcal{A}}$ (so-called $Q_{\mathcal{A}}$ -norm) will be denoted by $\|\cdot\|_{Q_{\mathcal{A}}}$. For simplicity we denote Q_T and Q_{T_1, T_2}

instead of $Q_{\mathcal{A}(T)}$ and $Q_{\mathcal{A}(T_1, T_2)}$ respectively for $T, T_1, T_2 \in L(\mathcal{H})$. In what follows, for $x, y \in \mathcal{H}$, $x \otimes y$ denotes rank one operator $(x \otimes y)z = (z, y)x$.

Proof of Theorem 3.1. The representation generated by $\{T_1, T_2\}$ is, by Lemma 2.2, absolutely continuous. We apply the technique used in [KP, Chapter 4 and 5]. Note that [KP, Theorem 4.5] is stated for absolutely continuous representations. The careful reader can notice that all the lemmas in [KP, Chapter 4] are true not only for c.n.u. contractions but also for any pair of contractions which generate an absolutely continuous representation.

Let, as usual, $[C_\lambda]$, $\lambda \in \mathbb{D}^2$, denote the element of $Q \stackrel{df}{=} Q_{\mathcal{A}(T_1, T_2)}$ corresponding to a weak-star continuous functional of evaluation at a point λ . Notice also that like in [KP] we can obtain that $\overline{\text{aco}}\{[C_\lambda] : \lambda \in \sigma_{le}(T_1, T_2) \cap \mathbb{D}^2\}$ contains the closed unit ball in Q . Let us recall that a dual algebra $\mathcal{A} \subset L(\mathcal{H})$ has property $X_{0,1}$ if for all $[L] \in Q$, $\|[L]\| \leq 1$, we have the following:

(*) there exist sequences $\{x_n\}_{n=1}^\infty, \{y_n\}_{n=1}^\infty \subset \mathcal{H}$ with $\|x_n\| \leq 1, \|y_n\| \leq 1$ for $n = 1, 2, \dots$ such that

$$(3.1) \quad \lim_{n \rightarrow \infty} \|[x_n \otimes y_n] - [L]\|_Q = 0,$$

$$(3.2) \quad \lim_{n \rightarrow \infty} \|[x_n \otimes z]\|_Q = 0, \quad \text{for all } z \in \mathcal{H},$$

$$(3.3) \quad \lim_{n \rightarrow \infty} \|[z \otimes y_n]\|_Q = 0, \quad \text{for all } z \in \mathcal{H}.$$

It is well known [BFP] that any dual algebra with property $X_{0,1}$ is reflexive. Also the set of functionals $[L] \in Q$ satisfying (*) is absolutely convex and closed. Therefore, for the reflexivity of $\mathcal{A}(T_1, T_2)$, hence also $\mathcal{W}(T_1, T_2)$, it is enough to prove (*) for $[C_\lambda]$ where $\lambda \in \sigma_{le}(T_1, T_2)$

Let $\lambda \in \sigma_{le}(T_1, T_2)$. In [KP, Lemma 4.10, 4.11], (3.1) and (3.2) were proved with our assumptions and even without C_0 , thus we need to show the statement (3.3). It will be shown below in Lemma 3.3, strengthening [KP, Lemma 5.1, 4.12].

It is an easy observation that in [R, Lemma 3.4] the assumption c.n.u. is not necessary, only the absolute continuity of T is needed. Moreover, weaker assumptions for the sequence $\{x_n\}$ are sufficient. So we have

Lemma 3.2. *Assume that $T \in L(\mathcal{H})$ is an a.c. contraction generating an isometric functional calculus. If x_n is a sequence such that $x_n \rightarrow 0$ weakly, $\|x_n\| = 1$, and $\|(T - \lambda)x_n\| \rightarrow 0$, then $\|[y \otimes x_n]\|_{Q_T} \rightarrow 0$ for all $y \in \mathcal{H}$.*

Now we can formulate

Lemma 3.3. *Assume T_1, T_2 generate an a.c. isometric representation and $T_1^n \rightarrow 0$ strongly. If x_n is a sequence such that $x_n \rightarrow 0$ weakly, $\|x_n\| = 1$, and $\|(T_2 - \lambda_2)x_n\| \rightarrow 0$, then $\|[y \otimes x_n]\|_Q \rightarrow 0$ for all $y \in \mathcal{H}$.*

Proof. By [SNF, Theorem II.2.1], choosing $V_1 \in L(\mathcal{H})$ a minimal isometric dilation of T_1^* , V_1 is a unilateral shift of uncertain multiplicity and also we have $T_1 = V_1^*|_{\mathcal{H}}$. On the other hand, by the commutant lifting theorem of Sz.-Nagy and Foias (see [SNF], [P, p. 484]) there is an operator W_2 (not necessarily isometry) preserving the norm of T_2^* , such that the pair $\{V_1, W_2\}$ dilates $\{T_1^*, T_2^*\}$. Let $\varepsilon > 0$. Choose $M > 0$ such that $\|(I - P_{\ker V_1^{*M}})y\| \leq \frac{\varepsilon}{3}$,

and denote $y_1 = (P_{\ker V_1^{*M}})y$, $y_2 = (I - P_{\ker V_1^{*M}})y$. By Hahn-Banach theorem, for each n , there is $f_n \in H^\infty(\mathbf{D}^2)$ such that

$$(3.4) \quad \|[y \otimes x_n]\|_Q = \langle f_n(T_1, T_2), [y \otimes x_n] \rangle = \langle f_n(T_1, T_2)y, x_n \rangle, \quad \|f_n\| = 1.$$

For each n we can decompose f_n as follows

$$(3.5) \quad f_n(z_1, z_2) = \sum_{k=0}^{M-1} a_{nk}(z_2)z_1^k + z_1^M h_n(z_1, z_2).$$

The functions a_{nk} are measurable. Moreover, since $a_{nk}(z_2)$ is the k -th Fourier coefficient of $f_n(\cdot, z_2)$, then $|a_{nk}(z_2)| \leq 1$, for $z_2 \in \mathbf{D}$, by (3.4). Thus $\|a_{nk}\| \leq 1$ and consequently $\|h_n\| \leq M + 1$. It is easy to check that the negative Fourier coefficients of every $a_{n,k}$ vanish. Hence for every n, k we get $a_{n,k} \in H^\infty(\mathbf{D})$, and $h_n \in H^\infty(\mathbf{D}^2)$.

Applying the Lebesgue type decomposition (2.1) to space \mathcal{H} and an easy calculation on elementary measures we get $\mathcal{H} \subset \mathcal{H}_0$. So, by the minimality of V_1 , we have $\mathcal{H} = \mathcal{H}_0$, hence the pair $\{V_1, W_2\}$ is a.c. By Lemma 2.1, also $\{V_1^*, W_2^*\}$ is a.c. By [K1, Proposition], the representation generated by T_2 is a.c. and so is W_2^* .

The above facts give us the existence of the functional calculus for all above-mentioned pairs and single operators. So, by Lemma 2.1(2) applied to the pairs $\{T_1, T_2\}$ and $\{V_1^*, W_2^*\}$, we have

$$(3.6) \quad \begin{aligned} \|[y \otimes x_n]\|_Q &= |\langle f_n(T_1, T_2)y, x_n \rangle| = |(y, \tilde{f}_n(T_1^*, T_2^*)x_n)| \\ &\leq |(y, \tilde{f}_n(V_1, W_2)x_n)| = |\langle f_n(V_1^*, W_2^*)y, x_n \rangle| \\ &\leq |\langle f_n(V_1^*, W_2^*)y_1, x_n \rangle| + |\langle f_n(V_1^*, W_2^*)y_2, x_n \rangle| \\ &\leq \sum_{k=0}^{M-1} |(a_{nk}(W_2^*)V_1^{*k}y_1, x_n)| \\ &\quad + |(V_1^{*M}h_n(V_1^*, W_2^*)y_1, x_n)| + \|f_n\| \|y_2\| \|x_n\|. \end{aligned}$$

The vector y_1 was defined such that $V_1^{*M}y_1 = 0$, thus the second component is equal 0. Observe that for all k

$$|(a_{nk}(W_2^*)V_1^{*k}y_1, x_n)| \leq \|[V_1^{*k}y_1 \otimes x_n]\|_{Q_{W_2^*}}.$$

On the other hand, by [SNF, Theorem II.2.3], W_2^* is an extension of T_2 , thus not only $\|(W_2^* - \lambda_2)x_n\| \rightarrow 0$, but also W_2^* generates an isometric representation, since T_2 does. Hence, Lemma 3.2 shows that $\|[V_1^{*k}y_1 \otimes x_n]\|_{Q_{W_2^*}} \rightarrow 0$ ($n \rightarrow \infty$). Hence we can choose n_0 such that for all $n > n_0$ we have $\|[V_1^{*k}y_1 \otimes x_n]\|_{Q_{W_2^*}} \leq \frac{\varepsilon}{3M}$ for $k = 0, 1, \dots, M - 1$. Coming back to the estimation of $\|[y \otimes x_n]\|_Q$, using (3.6) and the estimation of $\|y_2\|$, we obtain for $n > n_0$

$$\|[y \otimes x_n]\|_Q \leq \sum_{k=0}^{M-1} \|[V_1^{*k}y_1 \otimes x_n]\|_{Q_{W_2^*}} + \|f_n\| \|y_2\| \|x_n\| \leq \varepsilon.$$

The proof of the lemma is finished.

Let us notice in the end of the section that by taking adjoints using Theorem 3.1 we can obtain reflexivity assuming the dominating property for the right essential joint spectrum and C_0 assumption for one operator.

4. THE SECOND REFLEXIVITY RESULT

This section deals with pairs of operators which extend the case of doubly commuting operators. For any pair $\{T_1, T_2\}$ of commuting contractions we can construct a minimal isometric dilation $\{V_1, V_2\}$ of the pair $\{T_1^*, T_2^*\}$ (see [SNF]). One can see that V_i^* is a coisometric extension of T_i , $i = 1, 2$. We call $\{V_1^*, V_2^*\}$ a *joint coisometric extension* of $\{T_1, T_2\}$. It can be seen that it is minimal in standard meaning, for the details see [O]. We say that a pair $\{T_1, T_2\} \subset L(\mathcal{H})$ is *diagonally extendable* if there exist a Hilbert space $\mathcal{K} \supset \mathcal{H}$ and a minimal joint coisometric extension $\{B_1, B_2\} \subset L(\mathcal{K})$ of $\{T_1, T_2\}$ such that, for either $j = 1$ or $j = 2$, if \mathcal{K} is decomposed as $\mathcal{K} = \mathcal{S}_j \oplus \mathcal{R}_j$, relative to which the matrix for B_j has the form

$$B_j = \begin{pmatrix} S_j^* & 0 \\ 0 & R_j \end{pmatrix},$$

where $S_j^* \in L(\mathcal{S}_j)$ is a (unilateral) backward shift and $R_j \in L(\mathcal{R}_j)$ is a unitary operator, then the matrix for B_k , for $k \neq j$, relative to the decomposition $\mathcal{K} = \mathcal{S}_j \oplus \mathcal{R}_j$ has the form

$$B_k = \begin{pmatrix} A_s & 0 \\ 0 & A_r \end{pmatrix},$$

for some $A_s \in L(\mathcal{S}_j)$ and $A_r \in L(\mathcal{R}_j)$.

Let us recall the result of [O, Theorem 2.5] and [S, Lemma 1] as

Proposition 4.1. *With the above notation, a pair $\{T_1, T_2\} \subset L(\mathcal{H})$ is diagonally extendable if any of the following conditions holds:*

- (a) R_1 has no part of uniform multiplicity \aleph_0 ,
- (b) R_2 has no part of uniform multiplicity \aleph_0 ,
- (c) T_1 and T_2 doubly commute.

The main result of this section is the following

Theorem 4.2. *Let $\{T_1, T_2\} \subset L(\mathcal{H})$ be a pair of a.c. commuting contractions. Assume also that $\{T_1, T_2\} \subset L(\mathcal{H})$ is diagonally extendable. If $\sigma_{le}(T_1, T_2) \cap \mathbf{D}^2$ is dominating for $H^\infty(\mathbf{D}^2)$, then the algebra $\mathcal{W}(T_1, T_2)$ is reflexive.*

The proof is based on the same considerations as a proof of Theorem 3.1, using Lemma 4.3, given below, instead of Lemma 3.3.

Lemma 4.3. *Let $\{T_1, T_2\}$ be a diagonally extendable pair of contractions generating an a.c. isometric representation. If x_n is a sequence such that $x_n \rightarrow 0$ weakly, $\|x_n\| = 1$, and $\|(T_i - \lambda_i)x_n\| \rightarrow 0$ for $i = 1, 2$, then $\|[y \otimes x_n]\|_Q \rightarrow 0$ for all $y \in \mathcal{H}$.*

Proof. Without loss of generality, we assume that the space $\mathcal{K} \supset \mathcal{H}$ and operator $B_i \in L(\mathcal{K})$ extend T_i for $i = 1, 2$. Moreover, B_1 is a coisometry, $\mathcal{K} = \mathcal{S}_1 \oplus \mathcal{R}_1$,

$$B_1 = \begin{pmatrix} S_1^* & 0 \\ 0 & R_1 \end{pmatrix},$$

where $S_1^* \in L(\mathcal{S}_1)$ is a (unilateral) backward shift and $R_1 \in L(\mathcal{R}_1)$ is a unitary operator, and B_2 has the form

$$B_2 = \begin{pmatrix} A_s & 0 \\ 0 & A_r \end{pmatrix},$$

for some $A_s \in L(\mathcal{S}_1)$ and $A_r \in L(\mathcal{R}_1)$.

By decomposition (2.1) and the minimality of $\{B_1, B_2\}$ we conclude similarly as in the proof of Lemma 3.3 that $\{B_1, B_2\}$ is an a.c. pair. It also generates an isometric functional as an extension of the pair $\{T_1, T_2\}$. Thus we have

$$\begin{aligned} \|[y \otimes x_n]\|_Q &= \sup\{|\langle h(T_1, T_2)y, x_n \rangle| : h \in H^\infty(\mathbf{D}^2), \|h\|_\infty = 1\} \\ &= \sup\{|\langle h(B_1, B_2)y, x_n \rangle| : h \in H^\infty(\mathbf{D}^2), \|h\|_\infty = 1\} \\ &= \|[y \otimes x_n]\|_{Q_{B_1, B_2}}. \end{aligned}$$

So we need to show that the last tends to zero.

For any contraction T and $\lambda \in \mathbf{D}$, let T^λ denote the operator $(T - \lambda) \cdot (I - \bar{\lambda}T)^{-1}$. An easy calculation, based on [SNF], shows that the decomposition $\mathcal{X} = \mathcal{S}_1 \oplus \mathcal{R}_1$ is also a decomposition of $(B_1^*)^{\lambda_1}$ for a pure isometry and a unitary part. Let us note that $\mathcal{S}_1, \mathcal{R}_1$ reduce $B_2^{\lambda_2}$. Moreover, $B_i^{\lambda_i}$ is a coisometric extension of $T_i^{\lambda_i}$, for $i = 1, 2$. Let $x_n = x_n^s \oplus x_n^r, y = y^s \oplus y^r$ be the orthogonal sums with respect to the decomposition $\mathcal{X} = \mathcal{S}_1 \oplus \mathcal{R}_1$. Since $\|(T_1 - \lambda_1)x_n\| \rightarrow 0$, thus $\|T_1^{\lambda_1}x_n\| \rightarrow 0$. One can easily see that

$$\begin{aligned} (4.1) \quad \|x_n^r\| &\leq \|x_n - P_{\text{Ker}B_1^{\lambda_1}}x_n\| = \|(I - P_{\text{Ker}B_1^{\lambda_1}})x_n\| \\ &= \|(B_1^*)^{\lambda_1}B_1^{\lambda_1}x_n\| = \|T_1^{\lambda_1}x_n\| \rightarrow 0, \end{aligned}$$

Thus,

$$\begin{aligned} \|[y \otimes x_n]\|_{Q_{B_1, B_2}} &= \sup\{|\langle h(B_1, B_2)y, x_n \rangle| : h \in H^\infty(\mathbf{D}^2), \|h\|_\infty = 1\} \\ &\leq \sup\{|\langle h(S_1^*, A_s)y^s, x_n^s \rangle| : h \in H^\infty(\mathbf{D}^2), \|h\|_\infty = 1\} \\ &\quad + \sup\{|\langle h(R_1, A_r)y^r, x_n^r \rangle| : h \in H^\infty(\mathbf{D}^2), \|h\|_\infty = 1\} \\ &\leq \|[y^s \otimes x_n^s]\|_{Q_{S_1^*, A_s}} + \|y^r\| \|x_n^r\|. \end{aligned}$$

The second component converges to 0. To proof that the first one converges to 0, it is enough to show that the pair $\{S_1^*, A_s\}$ and sequence $z_n = x_n^s / \|x_n^s\|$ fulfill the assumption of Lemma 3.3. The operator S_1^* converges strongly to 0 as a backward shift. Since B_2 is an extension of T_2 , thus $\|(B_2 - \lambda_2)x_n\| \rightarrow 0$. Since $\|(B_2 - \lambda_2)x_n\|^2 = \|(A_s - \lambda_2)x_n^s\|^2 + \|(A_r - \lambda_2)x_n^r\|^2$, thus $\|(A_s - \lambda_2)x_n^s\| \rightarrow 0$ and $\|(A_s - \lambda_2)z_n\| \rightarrow 0$, since $\|x_n^s\| \rightarrow 1$. Hence, we can finish the proof of Lemma and Theorem 4.1 with the following

Lemma 4.4. *The pair $\{S_1^*, A_s\}$ generates a.c. isometric representation.*

Proof. To see that the pair $\{S_1^*, A_s\}$ generates an a.c. representation it is enough to notice that it is a restriction of the pair $\{B_1, B_2\}$, which is a.c. pair, and then make an easy calculation on elementary measures. Now we prove that $\sigma_{le}(T_1, T_2) \cap \mathbf{D}^2 \subset \sigma_{le}(S_1^*, A_s) \cap \mathbf{D}^2$. Let $(\lambda_1, \lambda_2) \in \sigma_{le}(T_1, T_2) \cap \mathbf{D}^2$, and $\{x_n\}$ is an orthonormal sequence such that $\lim_{n \rightarrow \infty} \|(T_i - \lambda_i)x_n\| = 0$, for $i = 1, 2$.

Hence, $\lim_{n \rightarrow \infty} \|(B_i - \lambda_i)x_n\| = 0$, for $i = 1, 2$, too. Let $x_n = x_n^s \oplus x_n^r$ is the orthogonal sum with respect to the decomposition $\mathcal{H} = \mathcal{S}_1 \oplus \mathcal{P}_1$. We can prove as above that $\|(A_s - \lambda_2)x_n^s\| \rightarrow 0$ and in the same way that $\|(S_1^* - \lambda_1)x_n^s\| \rightarrow 0$. Like in (4.1) it can be shown that $\|x_n^r\| \rightarrow 0$, hence $\|x_n^s\| \rightarrow 1$. Let $z_n = x_n^s / \|x_n^s\|$. It is easy to check that $\|(A_s - \lambda_2)z_n\| \rightarrow 0$ and $\|(S_1^* - \lambda_1)z_n\| \rightarrow 0$. Thus $(\lambda_1, \lambda_2) \in \sigma_{le}(S_1^*, A_s) \cap \mathbf{D}^2$ by Lemma 1.1. Hence, if $\sigma_{le}(T_1, T_2) \cap \mathbf{D}^2$ is dominating for $H^\infty(\mathbf{D}^2)$, then $\sigma_{le}(S_1^*, A_s) \cap \mathbf{D}^2$ also does. Hence, by [KP, Lemmas 4.1 and 4.4], the proof of the lemma is finished.

Remark 4.5. If a pair of operators fulfills assumptions of Theorem 3.1 or 4.2, then it has a common nontrivial invariant subspace.

REFERENCES

- [BFP] H. Bercovici, C. Foiaş, C. Pearcy, *Dual algebras with applications to invariant subspaces and dilation theory*, CBMS-NSF Regional Conf. Ser. in Appl. Math., vol. 56, Amer. Math. Soc., Providence, RI, 1985.
- [G] T. W. Gamelin, *Uniform algebras*, Prentice-Hall, Englewood Cliffs, NJ, 1969.
- [K1] M. Kosiek, *Lebesgue-type decomposition of a pair of commuting Hilbert space operators*, Bull. Acad. Polon. Sci. Ser. Sci. Math **27** (1979), 583–589.
- [K2] ———, *Representation generated by a finite number of Hilbert space operators*, Ann. Polon. Math. **44** (1984), 309–315.
- [KP] M. Kosiek and M. Ptak, *Reflexivity of N -tuples of contractions with rich joint left essential spectrum*, Integral Equations Operator Theory **13** (1990), 395–420.
- [O] A. Octavio, *Coisometric extension and functional calculus for pairs of commuting contractions* (to appear).
- [P] S. Parrot, *Unitary dilation for commuting contractions*, Pacific J. Math. **34** (1970), 481–490.
- [R] G. Robel, *On the structure of (BCP)-operators and related algebras*, J. Operator Theory **12** (1984), 23–45.
- [S] M. Słociński, *Isometric dilation of doubly commuting contractions and related models*, Bull. Acad. Sci. Pol. **25** (1977), 1233–1242.
- [SNF] B. Sz.-Nagy and C. Foiaş, *Analyse harmonique des operateurs de l'espace de Hilbert*, Masson et Cie, Academai Kiado, 1967.

INSTITUTE OF MATHEMATICS, JAGIELLONIAN UNIVERSITY, REYMONTA 4, 30-059 KRAKÓW, POLAND

E-mail address: mko@im.uj.edu.pl

DEPARTAMENTO DE MATEMATICAS, I. V. I. C., CARACAS, VENEZUELA

E-mail address: aoctavio@ivic.ivic.ve

INSTITUTE OF MATHEMATICS, UNIVERSITY OF AGRICULTURE, KRÓLEWSKA 6, 30-045 KRAKÓW, POLAND

E-mail address: rmptak@cyf-kr.edu.pl