

UNIVERSAL EQUIVARIANT BUNDLES

MITUTAKA MURAYAMA AND KAZUHISA SHIMAKAWA

(Communicated by Thomas Goodwillie)

ABSTRACT. We present a method of functorially constructing universal equivariant bundles.

INTRODUCTION

Let G and A be topological groups, and suppose G acts on A via a continuous homomorphism $\alpha: G \rightarrow \text{Aut } A$. In [16], tom Dieck introduced a notion of (G, α, A) -bundles which includes the notions of usual G -equivariant bundles [13], [18], [2], [5] (the case α is trivial) and Atiyah's vector bundles with Reality [1] (α is the complex conjugation $\mathbf{Z}/2\mathbf{Z} \rightarrow \text{Aut } U(n)$). He proved the existence of a universal (G, α, A) -bundle in the case G is a compact Lie group and A is an inductive limit of Lie groups with finitely many connected components. However, his construction as well as other constructions involving choice of representatives for equivalence classes of local objects (e.g., [17], [5]) is far from being functorial in G and A .

In this article, we present a new construction of universal (G, α, A) -bundles under the assumption that G is a compact Lie group or a discrete group and A is any topological group. It has three advantages: it is quite explicit, does not use the (generally unknown) classification of local objects, and, especially, is functorial in (G, α, A) , the meaning of which will be made clear in due course. Our method is a generalization of the functor category approach used in [15, §3] and is closely connected with the mapping space approach used by May [8] (cf. the final remark in §3).

1. NUMERABLE (G, α, A) -BUNDLES

In this section we recall the definition and basic properties of (G, α, A) -bundles. We write $ga = \alpha(g, a)$ for $g \in G$ and $a \in A$.

Definition 1.1. A principal (G, α, A) -bundle is a locally trivial principal A -bundle $p: E \rightarrow B$ enjoying the following properties:

- (1) Both E and B are left G -spaces, and p is G -equivariant.
- (2) For every $g \in G$, $a \in A$, and $e \in E$, we have $g(e \cdot a) = ge \cdot ga$.

Received by the editors February 16, 1993 and, in revised form, July 20, 1993.
1991 *Mathematics Subject Classification*. Primary 55R91.

©1995 American Mathematical Society
0002-9939/95 \$1.00 + \$.25 per page

The latter property is equivalent to saying that the *semi-direct product* $G \times_{\alpha} A$, the product $G \times A$ viewed as a topological group under the multiplication

$$(g, a)(g', a') = (gg', ga' \cdot a),$$

acts on the left of E by $(g, a)e = ge \cdot a$, $(g, a) \in G \times_{\alpha} A$.

There are evident notions of (G, α, A) -bundle maps and induced bundles. If F admits left G and A actions satisfying $g(a \cdot x) = ga \cdot gx$ then we can form the associated (G, α, A) -bundle $E \times_A F \rightarrow B$ in such a way that the (diagonal) G -action on the total space is compatible with the A -actions on its fibres.

Definition 1.2. For any closed subgroup H of G , a principal (G, α, A) -bundle over the homogeneous space G/H is called a local object.

A continuous map $\phi: H \rightarrow A$ is called a *crossed homomorphism* if $\phi(hk) = \phi(h) \cdot h\phi(k)$ holds for $h, k \in H$. Let $p: E \rightarrow G/H$ be a local object and let $e \in p^{-1}(H)$. Then the equation $he = e \cdot \phi(h)$ determines a crossed homomorphism $\phi: H \rightarrow A$, and the isotropy group of e under the $G \times_{\alpha} A$ -action on E is given as the graph of the map $h \mapsto \phi(h)^{-1}$. Since $G \times_{\alpha} A$ acts transitively on E , there is a $G \times_{\alpha} A$ -equivariant homeomorphism of E with $G \times_{\alpha} A/H$, the orbit space of $G \times_{\alpha} A$ with respect to the right H -action

$$(g, a) \mapsto (g, a)(h, \phi(h)^{-1}) = (gh, g\phi(h)^{-1} \cdot a), \quad h \in H.$$

We have shown:

Lemma 1.3. For every local object $p: E \rightarrow G/H$, there exist a crossed homomorphism $\phi: H \rightarrow A$ and a (G, α, A) -bundle equivalence $E \cong G \times_{\alpha} A/H$ over G/H .

More generally, a (G, α, A) -bundle with fibre F is called *trivial* if it is equivalent to a (G, α, A) -bundle of the form

$$(G \times_{\alpha} F) \times_H V \rightarrow G \times_H V,$$

where V is a left H -space and H acts on the right of $G \times_{\alpha} F = G \times F$ by $(g, f) \mapsto (gh, g\phi(h)^{-1} \cdot f)$, ϕ a crossed homomorphism $H \rightarrow A$. Clearly, a principal (G, α, A) -bundle is trivial if and only if there is a (G, α, A) -bundle map to a local object.

Definition 1.4. A (G, α, A) -bundle $p: E \rightarrow B$ is called locally trivial if there is an open cover $\{V_{\beta}\}$ of B such that for each β ,

- (1) V_{β} is G -invariant, and
- (2) $p^{-1}(V_{\beta}) \rightarrow V_{\beta}$ is a trivial (G, α, A) -bundle.

If, moreover, there exists a subordinate partition of unity $\{\lambda_{\beta}\}$ such that each λ_{β} is G -invariant then p is called a numerable (G, α, A) -bundle.

If G and A are compact Lie groups then every (G, α, A) -bundle is locally trivial if its base space B is completely regular and is numerable if B is paracompact. (See [5, Corollary 1.5] for an argument using the existence of a *slice* [3].) In general, local triviality restricts the type of G -action on the base space.

2. CLASSIFICATION OF EQUIVARIANT BUNDLES

A numerable principal (G, α, A) -bundle $p: E \rightarrow B$ is called *universal* if for any G -space X , the equivalence classes of numerable (G, α, A) -bundles over X are in bijective correspondence with the equivariant homotopy classes of G -maps of X into B under the correspondence given by induced bundles. The goal of this section is to prove the following. (Compare [16, Satz 5.1] and [5, Theorem 2.14].)

Theorem 2.1. *Let $p: E \rightarrow B$ be a numerable principal (G, α, A) -bundle. Then p is universal if and only if for every closed subgroup H and every crossed homomorphism $\phi: H \rightarrow A$, E is H -contractible under the H -action $e \mapsto he \cdot \phi(h)^{-1}$, $h \in H$.*

Write G -SEP and G -CHP for G -section extension property and G -covering homotopy property respectively. Arguing as in [5, Proposition 2.3, Corollaries 2.2 and 2.9], we have following three lemmas.

Lemma 2.2. *Let $p: E \rightarrow B$ be a G -map and $\{V_\beta\}$ be a numerable G -cover of B . If p has the G -SEP over each V_β then p has the G -SEP.*

Lemma 2.3. *Let H be a closed subgroup of G . Let V and F be H -spaces and give $V \times F$ the diagonal H -action. If F is H -contractible, the projection $p: G \times_H (V \times F) \rightarrow G \times_H V$ has the G -SEP.*

Lemma 2.4. *Let G be a compact Lie group or a discrete group, and let X be a G -space. If E is a trivial (G, α, A) -bundle over $X \times I$ then $E \cong (E|X) \times I$, hence has the G -CHP.*

As a consequence of Lemma 2.4 we have:

Corollary 2.5. *A numerable (G, α, A) -bundle has the G -CHP.*

We are now ready to prove Theorem 2.1. Assume that $p: E \rightarrow B$ satisfies the condition of the theorem. By Corollary 2.5, we get a natural map

$$\gamma: [X, B]^G \rightarrow \{\text{Equivalence classes of } (G, \alpha, A)\text{-bundles over } X\}$$

which assigns to each $f: X \rightarrow B$ the induced bundle $f^*p: f^*E \rightarrow X$.

We first show that γ is surjective. Given a numerable principal (G, α, A) -bundle $q: D \rightarrow X$, let $\langle q, p \rangle: \langle D, E \rangle \rightarrow X$ be the ‘functional bundle’ [4, 7.5] whose fibre on $x \in X$ consists of all admissible maps of D_x into E . Under the correspondence

$$(f: D_x \rightarrow E) \mapsto [d, f(d)] \in (D_x \times E)/A, \quad d \in D_x,$$

$\langle D, E \rangle$ is naturally isomorphic to the orbit space $(D \times E)/A$. In other words, $\langle q, p \rangle$ is the associated bundle of D with fibre E , where A acts on the left of E by $a \mapsto e \cdot a^{-1}$, $a \in A$. Since q is numerable, $\langle q, p \rangle$ is a numerable (G, α, A) -bundle.

By the definition, a section in $\langle q, p \rangle$ over $U \subset X$ is the same as a bundle map $q|U \rightarrow p$. Thus, to prove the surjectivity of γ , we need only show that $\langle q, p \rangle$ has the G -SEP. Moreover, by Lemma 2.2 we may assume that q is a trivial bundle $(G \times_\alpha A) \times_H V \rightarrow G \times_H V$.

Now the total space of $\langle q, p \rangle$ can be identified with $G \times_H (V \times E)$, where H acts on E by $(h, e) \mapsto he \cdot \phi(h)^{-1}$ and diagonally on $V \times E$, through a bundle isomorphism

$$\begin{aligned} ((G \times_\alpha A) \times_H V) \times_A E &\rightarrow G \times_H (V \times E), \\ [[g, a, v], e] &\mapsto [g, v, g^{-1}(e \cdot a^{-1})]. \end{aligned}$$

By Lemma 2.3, we see that $\langle q, p \rangle$ has the G -SEP.

The injectivity of γ is proved by the similar argument with X replaced by $X \times I$. Thus $p: E \rightarrow B$ is a universal (G, α, A) -bundle.

Conversely, assume that $p: E \rightarrow B$ is a universal (G, α, A) -bundle. As we shall see in the next section, there exists a (G, α, A) -bundle which satisfies the condition of the theorem, and so, is universal. Since any two universal bundles are equivalent, we conclude that p satisfies the condition of the theorem. \square

3. FUNCTOR CATEGORY MODEL FOR UNIVERSAL EQUIVARIANT BUNDLES

We present here a functorial construction of universal (G, α, A) -bundles by generalizing the method of [15, §3].

Let $p: UA \rightarrow BA$ be a universal A -bundle. By using a functorial construction of universal bundles (e.g., [11], [10], [12], [7]), we may assume that p is a (G, α, A) -bundle with respect to the G -action induced by α .

Let $\mathcal{G}A$ be a topological category with

$$\text{ob } \mathcal{G}A = BA, \quad \text{mor } \mathcal{G}A = \langle UA, UA \rangle = (UA \times UA)/A,$$

where $[b, a] \in \langle UA, UA \rangle$ is regarded as an arrow from $p(a)$ to $p(b)$, and with composition law $[c, b] \circ [b, a] = [c, a]$. (With suitable choice of representatives, every composable pair of arrows in $\mathcal{G}A$ is of the form $([c, b], [b, a])$.)

Similarly, let $\mathcal{S}A$ be a topological category with

$$\text{ob } \mathcal{S}A = UA, \quad \text{mor } \mathcal{S}A = \langle UA, UA \rangle \times_{BA} UA,$$

where $([b, a], a) \in \langle UA, UA \rangle \times_{BA} UA$ is regarded as an arrow from a to b , and with composition law induced by that of $\mathcal{G}A$.

$\mathcal{S}A$ is fibred over $\mathcal{G}A$ by a functor $\pi: \mathcal{S}A \rightarrow \mathcal{G}A$ induced by p . Moreover, under the natural isomorphism $UA \times UA \cong \langle UA, UA \rangle \times_{BA} UA$, $(b, a) \mapsto ([b, a], a)$, $\mathcal{S}A$ is identified with the category with a unique isomorphism between each pair of elements of UA . Hence $\mathcal{S}A$ is equivalent to the trivial category with one object and one morphism.

Denote by EG the translation category of G , that is, a contractible category with $\text{ob } EG = G$ and $\text{mor } EG = G \times G$, where $(y, x) \in G \times G$ is viewed as a unique isomorphism $x \rightarrow y$. Let $\text{Cat}(EG, \mathcal{S}A)$ be the category of continuous functors and continuous natural transformations of EG into $\mathcal{S}A$. Since $\mathcal{S}A$ has a unique isomorphism between each pair of its objects, a functor $f: EG \rightarrow \mathcal{S}A$ is completely determined by its restriction to objects, $\text{ob } f: G \rightarrow UA$. Thus the nerve of the contractible category $\text{Cat}(EG, \mathcal{S}A)$ is given by

$$N_k \text{Cat}(EG, \mathcal{S}A) = \text{Map}(G, UA)^{k+1}, \quad k \geq 0,$$

where $\text{Map}(G, UA)$ is the set of continuous maps of G into UA , together with face and degeneracy operators induced by projections and diagonal embeddings respectively.

Let $G \times_\alpha A$ act on $\text{Cat}(EG, \mathcal{S}A)$ via its action on $\text{Map}(G, UA)$,

$$(g, a)f(x) = gf(xg) \cdot a, \quad (g, a) \in G \times_\alpha A.$$

We topologize $\text{Cat}(EG, \mathcal{S}A)$ in two ways. The *strong topology* is obtained by imposing on each $N_k \text{Cat}(EG, \mathcal{S}A)$ the strongest topology such that its $G \times_\alpha A$ -action is continuous. On the other hand, the compact-open topology on $\text{Map}(G, UA)$ (retopologized, if necessary, so as to be compactly generated [9, §2]) gives rise to another topology on $\text{Cat}(EG, \mathcal{S}A)$, called *natural topology*. With respect to either topology, $G \times_\alpha A$ acts continuously on $\text{Cat}(EG, \mathcal{S}A)$ and the functor

$$\eta: \text{Cat}(EG, \mathcal{S}A) \rightarrow \text{Cat}(EG, \mathcal{G}A)$$

induced by $\pi: \mathcal{S}A \rightarrow \mathcal{G}A$ is a G -equivariant continuous functor. In fact, $N_k \eta$ is the natural map of $\text{Map}(G, UA)^{k+1}$ onto the orbit space

$$N_k \text{Cat}(EG, \mathcal{G}A) = \text{Map}(G, UA)^{k+1}/A, \quad k \geq 0,$$

with respect to the diagonal A -action.

Recall from [14, Appendix A] that (the nerve of) a topological category C has the naive realization $|C|$ and also the fat realization $\|C\|$. With the strong topology on $\text{Cat}(EG, \mathcal{S}A)$, we will prove:

Theorem 3.1.

$$\|\eta\|: \|\text{Cat}(EG, \mathcal{S}A)\| \rightarrow \|\text{Cat}(EG, \mathcal{G}A)\|$$

is a universal (G, α, A) -bundle.

Remark. It is desirable if we can prove Theorem 3.1 with the natural topology on $\text{Cat}(EG, \mathcal{S}A)$ and/or with the naive realization instead of the fat one. This is certainly the case if both G and A are compact Lie groups.

Before proving the theorem, let us examine the functorial property of $\|\eta\|$ with respect to (G, α, A) .

A morphism of (G, α, A) to another such system (K, γ, C) is a pair of homomorphisms $l: K \rightarrow G$ and $r: A \rightarrow C$ such that $r(l(k)a) = kr(a)$ holds for $k \in K, a \in A$. Such a morphism $\mu = (l, r)$ functorially gives rise to a commutative square

$$\begin{array}{ccc} \|\text{Cat}(EG, \mathcal{S}A)\| & \xrightarrow{U\mu} & \|\text{Cat}(EK, \mathcal{S}C)\| \\ \eta \downarrow & & \downarrow \eta \\ \|\text{Cat}(EG, \mathcal{G}A)\| & \xrightarrow{B\mu} & \|\text{Cat}(EK, \mathcal{G}C)\| \end{array}$$

where $U\mu$ and $B\mu$ are induced by the functors $EK \rightarrow EG$ and $UA \rightarrow UC$ associated with l and r respectively.

Let A act on C via r and let $K \times_{\gamma} C$ act on $\|\text{Cat}(EG, \mathcal{S}A)\| \times_A C$ by $(k, c)[e, d] = [l(k)e, kd \cdot c]$, $(k, c) \in K \times_{\gamma} C$. Then

$$\|\text{Cat}(EG, \mathcal{S}A)\| \times_A C \rightarrow \|\text{Cat}(EK, \mathcal{S}C)\|, \quad [e, c] \mapsto U\mu(e) \cdot c,$$

is a (K, γ, C) -bundle map over $B\mu$ or, equivalently, $B\mu$ is a classifying map for the associated bundle $\|\text{Cat}(EG, \mathcal{S}A)\| \times_A C \rightarrow \|\text{Cat}(EG, \mathcal{S}A)\|$ regarded as a principal (K, γ, C) -bundle. In particular, if l is the inclusion of a closed subgroup K of G and r the identity of A then $B\mu$ corresponds to the restriction of the group action from G to K and, hence, is a K -equivalence. On the other hand, if l is the identity and r the inclusion of a closed subgroup A of C then $B\mu$ corresponds to the reduction of the structure groups (in the equivariant setting).

To prove Theorem 3.1, it suffices to show the following two lemmas.

Lemma 3.2. $\|\eta\|$ is a numerable principal (G, α, A) -bundle.

Lemma 3.3. For any crossed homomorphism $\phi: H \rightarrow A$, $\|\text{Cat}(EG, \mathcal{S}A)\|$ is H -contractible under the H -action $e \mapsto he \cdot \phi(h)^{-1}$, $h \in H$.

Lemma 3.2 is proved by the argument of [4, §8] together with the fact that

$$N_k \eta: N_k \text{Cat}(EG, \mathcal{S}A) \rightarrow N_k \text{Cat}(EG, \mathcal{S}A)$$

is a disjoint union of local objects, as follows by the property of the strong topology.

We now prove Lemma 3.3. In fact, we will show that $\text{Cat}(EG, \mathcal{S}A)$ has an H -fixed terminal object, implying both the naive and the fat realizations of $\text{Cat}(EG, \mathcal{S}A)$ are H -contractible with respect to any topology such that the $G \times_{\alpha} A$ -action is continuous. (See [10, Proposition 1.2] for the construction of H -contraction.)

Let $q: G \times_{\alpha} A/H \rightarrow G/H$ be the local object corresponding to a crossed homomorphism ϕ . Since $p: UA \rightarrow BA$ is universal, there exists an A -bundle map $F: G \times_{\alpha} A/H \rightarrow UA$. Define an object $f \in \text{ob Cat}(EG, \mathcal{S}A) = \text{Map}(G, UA)$ by

$$f(x) = x^{-1}F([x, 1]) \in UA, \quad x \in G.$$

Then one easily checks that $(hf \cdot \phi(h)^{-1})(x) = f(x)$ holds for any $h \in H$ and $x \in G$. Thus f is fixed by the H -action $e \mapsto he \cdot \phi(h)^{-1}$ on $\text{Cat}(EG, \mathcal{S}A)$.

Since there exists a unique isomorphism $f' \rightarrow f$ for any object f' in $\text{Cat}(EG, \mathcal{S}A)$, f is an H -fixed terminal (and initial) object. This proves Lemma 3.3. \square

Notice that in the argument above, we only used the property that $p: UA \rightarrow BA$ classifies principal A -bundles over homogeneous spaces G/H . In particular, if G is finite, we can take as p the trivial bundle $A \rightarrow *$. In this case, $\mathcal{S}A$ is the translation category EA of A and $\mathcal{S}A$ is the group A regarded as a category with one object. Since $\text{Map}(G, A)$ is a product of copies of A , the nerve of $\text{Cat}(EG, \mathcal{S}A)$ as well as that of $\text{Cat}(EG, \mathcal{S}A)$ has particularly simple structure and can be regarded as an equivariant generalization of the geometric bar construction [10], [7].

Remark. May showed in [8] that if G and A are compact Lie groups then the natural map

$$\text{Map}(UG, UA) \rightarrow \text{Map}(UG, UA)/A$$

is a model for universal (G, α, A) -bundle. This mapping space model is connected with our functor category model through a $G \times_{\alpha} A$ -equivalence

$$\begin{aligned} |\text{Cat}(EG, \mathcal{S}A)| &\cong |\text{Map}_{\bullet}(N_{\bullet}EG, N_{\bullet}\mathcal{S}A)| \rightarrow \text{Map}(|EG|, |\mathcal{S}A|) \\ &\simeq \text{Map}(UG, UA), \end{aligned}$$

where $\text{Map}_{\bullet}(-, -)$ denotes the simplicial mapping space. We hope to discuss elsewhere a further generalization of our functor category model to the case of *generalized equivariant bundles* introduced by [6] and its connection with the corresponding generalization of the mapping space model [8].

REFERENCES

1. M. F. Atiyah, *K-theory and reality*, Quart. J. Math. Oxford Ser. (2) **17** (1966), 367–386.
2. E. Bierstone, *The equivariant covering homotopy property for differentiable G -fibre bundles*, J. Differential Geom. **8** (1973), 615–622.
3. G. Bredon, *Introduction to compact transformation groups*, Pure Appl. Math., vol. 46, Academic Press, New York, 1972.
4. A. Dold, *Partitions of unity in the theory of fibrations*, Ann. of Math. (2) **78** (1963), 223–255.
5. R. K. Lashof, *Equivariant bundles*, Illinois J. Math. **26** (1982), 257–271.
6. R. K. Lashof and J. P. May, *Generalized equivariant bundles*, Bull. Soc. Math. Belg. Sér. B **38** (1986), 265–271.
7. J. P. May, *Classifying spaces and fibrations*, Mem. Amer. Math. Soc., vol. 155, Amer. Math. Soc., Providence, RI, 1975.
8. ———, *Some remarks on equivariant bundles and classifying spaces*, Astérisque **191** (1990), 239–253.
9. M. C. McCord, *Classifying spaces and infinite symmetric products*, Trans. Amer. Math. Soc. **146** (1969), 273–298.
10. R. Milgram, *The bar construction and abelian H -spaces*, Illinois J. Math. **11** (1967), 242–250.
11. J. W. Milnor, *Construction of universal bundles. II*, Ann. of Math. (2) **63** (1956), 430–436.
12. G. Segal, *Classifying spaces and spectral sequences*, Inst. Hautes Études. Sci. Publ. Math. **34** (1968), 105–112.
13. ———, *Equivariant K -theory*, Inst. Hautes Études. Sci. Publ. Math. **34** (1968), 129–151.
14. ———, *Categories and cohomology theories*, Topology **13** (1974), 293–312.
15. K. Shimakawa, *Infinite loop G -spaces associated to monoidal G -graded categories*, Publ. Res. Inst. Math. Sci. **25** (1989), 239–262.
16. T. tom Dieck, *Faserbündel mit gruppenoperation*, Arch. Math. **20** (1969), 136–143.
17. S. Waner, *Equivariant classifying spaces and fibrations*, Trans. Amer. Math. Soc. **258** (1980), 385–405.
18. S. Wasserman, *Equivariant differential topology*, Topology **8** (1969), 127–150.

DEPARTMENT OF MATHEMATICS, TOKYO INSTITUTE OF TECHNOLOGY, O-OKAYAMA, MEGURO-KU, TOKYO, JAPAN

E-mail address: murayama@math.titech.ac.jp

DEPARTMENT OF MATHEMATICS, OKAYAMA UNIVERSITY, OKAYAMA 700, JAPAN

E-mail address: yfae0866@ccews2.cc.okayama-u.ac.jp