

SOBOLEV IMBEDDING THEOREMS IN BORDERLINE CASES

NICOLA FUSCO, PIERRE LOUIS LIONS, AND CARLO SBORDONE

(Communicated by Barbara Lee Keyfitz)

ABSTRACT. An imbedding theorem is given for functions whose gradient belongs to a class slightly larger than $L^n(\Omega)$, $\Omega \subset \mathbb{R}^n$.

1. INTRODUCTION

We wish here to come back to the well-known imbedding theorem for $W^{1,n}(\Omega)$ functions due to N. Trudinger [T] and J. Moser [Mo].

One way to shed some light on the phenomena involved consists in extending this result to other spaces closely related to $W^{1,n}$, as in [ALT].

We consider here spaces of functions that are larger than $W^{1,n}$, but are contained in $\bigcap_{1 < p < n} W^{1,p}$. More precisely, we consider functions u whose gradient Du satisfies for some $\sigma > 0$

$$(1) \quad \int_{\Omega} |Du|^n \log^{-\sigma}(e + |Du|) dx < \infty,$$

a class of functions motivated by recent work on the regularity properties of Jacobians (see [BFS], [CLMS], [Mu], [IS], [Gr]).

In this paper we prove, in particular, that if $u \in W_0^{1,1}$ and (1) holds, then

$$(2) \quad \int_{\Omega} \exp\left(\frac{|u(x)|^\alpha}{\lambda}\right) dx < \infty$$

for any $\lambda > 0$, where $\alpha = \frac{n}{n-1+\sigma}$.

This result relies on the simple observation that condition (2) is equivalent to saying that

$$\lim_{p \rightarrow \infty} \frac{1}{p} \left(\int_{\Omega} |u|^{\alpha p} dx \right)^{1/p} = 0$$

(where \int_{Ω} stands for $\frac{1}{|\Omega|} \int_{\Omega}$).

More generally, our results apply to functions such that

$$\sup_{\varepsilon > 0} \varepsilon^\sigma \int_{\Omega} |Du|^{n-\varepsilon} dx < \infty$$

(see Theorem 2).

Received by the editors July 7, 1993 and, in revised form, September 12, 1994.
1991 *Mathematics Subject Classification*. Primary 46E35.

2. SOME PRELIMINARY RESULTS

Let Ω be an open set in \mathbb{R}^n , with finite measure $|\Omega|$. We denote by $\text{EXP} = \text{EXP}(\Omega)$ the set of functions $g : \Omega \rightarrow \mathbb{R}$ such that there exists $\lambda > 0$ for which

$$\int_{\Omega} \exp\left(\frac{|g|}{\lambda}\right) dx < \infty.$$

One way to test if a function f belongs to EXP is given by the following proposition, whose proof is an easy exercise (see [G, Chapter VI, Example 17]).

Proposition 1. *Let $g : \Omega \rightarrow \mathbb{R}$ be a measurable function. Set*

$$E(g) = e \limsup_{p \rightarrow \infty} \frac{1}{p} \left(\int_{\Omega} |g|^p dx \right)^{1/p}.$$

Then

$$E(g) = \inf \left\{ \lambda > 0 : \int_{\Omega} \exp\left(\frac{|g|}{\lambda}\right) dx < \infty \right\}.$$

Remark. This proposition says that $g \in \text{EXP}$ if and only if $E(g) < \infty$. In particular, if $f \in L^\infty(\Omega)$ we have $E(f) = 0$ and $E(g - f) = E(g)$ for any $g \in \text{EXP}$.

We recall that EXP is a Banach space under the norm (see [RR])

$$\|g\|_{\text{EXP}} = \inf \left\{ \lambda > 0 : \int_{\Omega} \exp\left(\frac{|g|}{\lambda}\right) dx \leq 2 \right\}.$$

We also remark that L^∞ is not a dense subspace of EXP . Indeed one can easily prove the following

Proposition 2. *$E(g) = 0$ if and only if there exists a sequence $f_h \in L^\infty$ such that $\|f_h - g\|_{\text{EXP}} \rightarrow 0$.*

Proof. If f_h verifies $\|f_h - g\|_{\text{EXP}} \rightarrow 0$, then by the previous remark:

$$E(g) = E(f_h - g) \leq \|f_h - g\|_{\text{EXP}}.$$

Conversely, assume that $E(g) = 0$, and let $\varepsilon > 0$. By definition, we have

$$\int_{\Omega} \exp\left(\frac{|g|}{\varepsilon}\right) dx < \infty;$$

then there exists h_ε such that, for $h > h_\varepsilon$,

$$\frac{1}{|\Omega|} \int_{\{|g|>h\}} \exp\left(\frac{|g|}{\varepsilon}\right) dx \leq 1.$$

If we set

$$f_h(x) = \begin{cases} g(x) & \text{if } |g(x)| \leq h, \\ 0 & \text{otherwise,} \end{cases}$$

then

$$\int_{\Omega} \exp\left(\frac{|f_h - g|}{\varepsilon}\right) dx = \frac{1}{|\Omega|} \int_{\{|g|>h\}} \exp\left(\frac{|g|}{\varepsilon}\right) dx + \frac{|\{|g| \leq h\}|}{|\Omega|} \leq 2.$$

Therefore we have

$$\|f_h - g\|_{\text{EXP}} \leq \varepsilon \quad \text{for } h > h_\varepsilon.$$

3. THE MAIN RESULTS

The first Sobolev type result we want to deduce is the following

Theorem 1. *If $u \in W_0^{1,1}(\Omega)$ satisfies for some $\sigma > 0$ the condition*

$$(3) \quad \lim_{\varepsilon \rightarrow 0^+} \varepsilon^\sigma \int_{\Omega} |Du|^{n-\varepsilon} dx = 0,$$

then, for any $c > 0$, we have

$$(4) \quad \int_{\Omega} \exp\left(\frac{|u|^\alpha}{c}\right) dx < \infty$$

with $\alpha = \frac{n}{n-1+\sigma}$, i.e. $E(|u|^\alpha) = 0$.

Let us recall that the Riesz potential If of a function $f \in L^1(\Omega)$ is defined as

$$If(x) = \int_{\Omega} f(y)|x - y|^{1-n} dy.$$

We will use the following theorem ([GT, Theorem 7.34]).

Theorem. *Let $1 \leq p, q \leq \infty$ and assume that $0 \leq \delta = \frac{1}{p} - \frac{1}{q} < \frac{1}{n}$. Then*

$$(5) \quad \|If\|_{L^q(\Omega)} \leq \left(\frac{1-\delta}{\frac{1}{n}-\delta}\right)^{1-\delta} \omega_n^{1-1/n} |\Omega|^{1/n-\delta} \|f\|_{L^p(\Omega)}$$

where ω_n is the measure of the unit ball in \mathbb{R}^n .

We can now pass to the

Proof of Theorem 1. Since

$$|u(x)| \leq \frac{1}{n\omega_n} I(|Du|)(x),$$

using Proposition 1, it will be enough to prove that

$$(6) \quad \lim_{\varepsilon \rightarrow 0^+} \varepsilon^{1/\alpha} \left(\int_{\Omega} [I(|Du|)]^{1/\varepsilon} dx \right)^\varepsilon = 0.$$

In (5) let us take $q = \frac{1}{\varepsilon}, p = n - \varepsilon$, and note that, if $0 < \varepsilon \leq \frac{1}{n}$, then certainly $\delta < \frac{1}{n}$; hence we have

$$\varepsilon^{1/\alpha} \left(\int_{\Omega} [I(|Du|)]^{1/\varepsilon} dx \right)^\varepsilon \leq \varepsilon^{1/\alpha} \left(\frac{1-\delta}{\frac{1}{n}-\delta}\right)^{1-\delta} \omega_n^{1-1/n} |\Omega|^{1/n-\delta-\varepsilon} \|Du\|_{L^{n-\varepsilon}(\Omega)}.$$

It is easy to check that, if $0 < \varepsilon \leq \frac{1}{n}$, then

$$\left(\frac{1-\delta}{\frac{1}{n}-\delta}\right)^{1-\delta} \leq c(n)\varepsilon^{-(n-1)/n},$$

so we have

$$\begin{aligned} & \varepsilon^{1/\alpha} \left(\int_{\Omega} [I(|Du|)]^{1/\varepsilon} dx \right)^\varepsilon \\ & \leq c(n, |\Omega|) \varepsilon^{1/\alpha - (n-1)/n} \varepsilon^{-\sigma/(n-\varepsilon)} \left(\varepsilon^\sigma \int_{\Omega} |Du|^{n-\varepsilon} dx \right)^{1/(n-\varepsilon)} \\ & \leq c(n, |\Omega|) \varepsilon^{1/\alpha - (n-1)/n - \sigma/n} \left(\varepsilon^\sigma \int_{\Omega} |Du|^{n-\varepsilon} dx \right)^{1/(n-\varepsilon)} \end{aligned}$$

which proves (6), since $\frac{1}{\alpha} - \frac{n-1}{n} - \frac{\sigma}{n} = 0$.

Remark 1. If $g \in L^1_{loc}(\Omega)$ satisfies

$$\int_{\Omega} |g|^n \log^{-\sigma}(e + g) \, dx < +\infty,$$

then (see [BFS, Lemma 3])

$$\lim_{\varepsilon \rightarrow 0^+} \varepsilon^\sigma \int_{\Omega} |g|^{n-\varepsilon} \, dx = 0.$$

From this it follows that if $u \in W_0^{1,1}(\Omega)$ and $|Du|^n \log^{-\sigma}(e + |Du|) \in L^1(\Omega)$, then

$$\int_{\Omega} \exp\left(\frac{|u|^\alpha}{c}\right) < \infty$$

for any $c > 0$, where α is given by Theorem 1. Note that, in any dimension, if $\sigma = 1$, then $\alpha = 1$.

Remark 2. Theorem 1 is optimal in the sense that the exponent α cannot be improved. In fact, if

$$u(x) = \frac{\log|x|}{|\log|\log|x||^\vartheta}$$

for $|x|$ small, with $\vartheta > 1/n$, then $|Du|^n \log^{-1}(e + |Du|)$ is in L^1 , while, for any $c > 0, \delta > 0$

$$\int \exp\left(\frac{|u|^{1+\delta}}{c}\right) = \infty.$$

We conclude by proving an imbedding theorem in terms of the quantity on the left-hand side of (3), which can be regarded as an extension of Trudinger’s imbedding theorem.

Theorem 2. *Let $u \in W_0^{1,1}(\Omega)$ satisfy for some $\sigma \geq 0$*

$$(7) \quad M = \sup_{0 < \varepsilon \leq 1} \left(\varepsilon^\sigma \int_{\Omega} |Du|^{n-\varepsilon} \, dx \right)^{1/(n-\varepsilon)} < \infty.$$

Then, if $\alpha = \frac{n}{n-1+\sigma}$, there exist $c_1 = c_1(n, \sigma)$, $c_2 = c_2(n, \sigma)$ such that

$$(8) \quad \int_{\Omega} \exp\left(\frac{|u|}{c_1 M |\Omega|^{1/n}}\right)^\alpha \, dx \leq c_2.$$

Proof. Arguing as in the proof of Theorem 1, we have for any $0 < \varepsilon \leq \frac{1}{n}$

$$\begin{aligned} \varepsilon^{1/\alpha} \left(\int_{\Omega} |u(x)|^{1/\varepsilon} \right)^\varepsilon &\leq \frac{\varepsilon^{1/\alpha}}{n\omega_n} \left(\int_{\Omega} [I(|Du|)]^{1/\varepsilon} \, dx \right)^\varepsilon \\ &\leq c(n) \varepsilon^{1/\alpha - (n-1)/n} |\Omega|^{1/n} \left(\varepsilon^\sigma \int_{\Omega} |Du|^{n-\varepsilon} \, dx \right)^{1/(n-\varepsilon)} \\ &\leq c(n) \varepsilon^{1/\alpha + 1/n - 1 - \sigma/n} |\Omega|^{1/n} M \\ &= c(n) |\Omega|^{1/n} M. \end{aligned}$$

From this inequality, taking $\alpha\varepsilon = \frac{1}{p}$, we get

$$\sup_{p \geq \frac{n}{\alpha}} \frac{1}{p} \left(\int_{\Omega} (|u|^{\alpha})^p dx \right)^{1/p} \leq [c(n)|\Omega|^{1/n}M]^{\alpha}.$$

The result now follows, noting that if

$$\lambda = \sup_{p \geq p_0} \frac{1}{p} \left(\int_{\Omega} |u|^p dx \right)^{1/p},$$

where $p_0 \geq 1$, then

$$\int_{\Omega} \exp\left(\frac{|u|}{2e\lambda}\right) dx \leq c(p_0).$$

Remark 3. If $|Du| \in L^{n,\infty} = \text{weak-}L^n$, then (7) holds with $\sigma = 1$ (see [IS]) and then (8) holds with $\alpha = 1$. For this and similar results, see [ALT]. Note that Theorem 2 holds true also in the case $\sigma = 0$. In this case the above result reduces to Trudinger's theorem (see [T]).

REFERENCES

- [ALT] A. Alvino, P. L. Lions, and G. Trombetti, *On optimization problems with prescribed rearrangements*, *Nonlinear Anal.* TMA **13** (1989), 185–220. MR **90c**:90236
- [BFS] H. Brèzis, N. Fusco, and C. Sbordone, *Integrability for the Jacobian of orientation preserving mappings*, *J. Funct. Anal.* **115** (1993), 425–431. MR **94j**:26024
- [CLMS] R. Coifman, P. L. Lions, Y. Meyer, and S. Semmes, *Compensated compactness and Hardy spaces*, *J. Math. Pures Appl.* **72** (1993), 247–286. MR **95d**:46033
- [G] J. B. Garnett, *Bounded analytic functions*, Academic Press, New York, 1981. MR **83g**:30037
- [GT] D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*, Springer, Berlin and New York, 1977. MR **57**:13109
- [Gr] L. Greco, *A remark on the equality $\det Df = \text{Det } Df$* , *Differential and Integral Equations* **6** (1993), 1089–1100. MR **95a**:49104
- [IS] T. Iwaniec and C. Sbordone, *On the integrability of the Jacobian under minimal hypotheses*, *Arch. Rational Mech. Anal.* **119** (1992), 129–143. MR **93i**:49057
- [Mo] J. Moser, *A sharp form of an inequality by N. Trudinger*, *Indiana Univ. Math. J.* **20** (1971), 1077–1092. MR **46**:662
- [Mu] S. Müller, *Higher integrability of determinants and weak convergence in L^1* , *J. Reine Angew. Math.* **412** (1990), 20–34. MR **92b**:49026
- [RR] M. M. Rao and Z. D. Ren, *Theory of Orlicz spaces*, Marcel Dekker, New York, 1991. MR **92e**:46059
- [T] N. S. Trudinger, *On imbeddings into Orlicz spaces and some applications*, *J. Math. Mech.* **17** (1967), 473–483. MR **35**:7121

(N. Fusco and C. Sbordone) DIPARTIMENTO DI MATEMATICA E APPLICAZIONI, UNIVERSITÀ DI NAPOLI, VIA CINTIA, 80126 NAPOLI, ITALY

(P. L. Lions) CEREMADE, PLACE DU MARÉCHAL DE LATTRE DE TASSIGNY, 75775 PARIS CEDEX 16, FRANCE