

FIXED POINTS IN BOUNDARIES OF NEGATIVELY CURVED GROUPS

PHILIP L. BOWERS AND KIM RUANE

(Communicated by James E. West)

An important feature of the natural action of a negatively curved (= word hyperbolic) group on its boundary is that the fixed points of hyperbolic (= infinite order) elements form a dense subset of the boundary. Gromov states this in his influential treatise *Hyperbolic Groups* [4, 8.2 D] and his explanation involves Stallings's theorem on groups with infinitely many ends. This short note provides an elementary proof of this fact. The proof relies on a simple enhancement of the Pumping Lemma from the theory of finite state automata and on the fact that the geodesic words with respect to a fixed generating set for a negatively curved group form a regular language. For background material on negatively curved groups see [1] and on automata and regular languages on groups see [2]. For a different proof see [5].

Definition. Let L be a language over the finite alphabet A . An infinite sequence $\{a_i\}$ of letters from A is an L -ray provided, for every integer $n > 0$, there exists an integer $N \geq n$ such that $a_1 \cdots a_N$ is a word in the language L .

The following lemma, which we term an *Ultra-Pumping Lemma* (UPL), is our aforementioned simple enhancement of the Pumping Lemma.

Ultra-Pumping Lemma. Let $\{a_i\}$ be an L -ray where L is a regular language over the finite alphabet A . There exists an integer $M \geq 0$ such that, for each integer $n > 0$, there exists an integer $N \geq n$ such that $(a_1 \cdots a_M)(a_{M+1} \cdots a_{M+N})^r \in L$ for every integer $r \geq 0$.

Proof. Let W be a finite state automaton accepting L . The L -ray $\{a_i\}$ describes an infinite path in W and, as there are only finitely many (accept) states in W and infinitely many initial segments of $\{a_i\}$ in L , this path visits some accept state s_∞ infinitely often. Let M be the step at which this path first visits s_∞ and choose $N \geq n$ such that $a_{M+1} \cdots a_{M+N}$ describes a loop in W based at s_∞ . \square

We use the term δ -negatively curved in the sense of Gromov's δ -hyperbolic condition [4, p. 89]. A group G is δ -negatively curved with respect to a finite generating set $A = A^{-1}$ if the corresponding Cayley graph $\Gamma_A(G)$ with the usual word metric has a δ -hyperbolic inner product based at the identity; see [1, Definition 1.1, p. 8]. A group G is *negatively curved* (or *word hyperbolic*) if it is δ -negatively curved with

Received by the editors May 31, 1994.

1991 *Mathematics Subject Classification.* Primary 20F32.

Key words and phrases. Negatively curved group, Pumping Lemma, boundary of negatively curved group.

respect to some finite generating set for some $\delta \geq 0$. For an infinite order element g of the negatively curved group G , let $g^{\pm\infty}$ denote the two fixed points of the action of g on $G \cup \partial G$, where ∂G denotes the boundary of G . Recall that $g^{\pm\infty}$ are elements of ∂G and $\lim g^i = g^{+\infty}$ and $\lim g^{-i} = g^{-\infty}$.

Theorem [4, 8.2D]. *Let G be a negatively curved group with boundary ∂G . The set*

$$\{g^{\pm\infty} : g \in G, \text{ order}(g) = \infty\}$$

is a dense subset of ∂G .

Proof. Recall that ∂G is metrizable [1] and as such sequences suffice to verify denseness. Let G be δ -negatively curved with respect to the finite generating set $A = A^{-1}$ and let $x \in \partial G$. Choose an infinite geodesic path $a_1 a_2 \cdots$ based at 1 in the Cayley graph $\Gamma_A(G)$ such that $\lim \overline{a_1 \cdots a_i} = x$. Here, $a_1 \cdots a_i$ denotes a word over A and $\overline{a_1 \cdots a_i}$ denotes the element of G obtained by multiplying. The sequence $\{a_i\}$ is an L -ray, where L is the language of geodesic words of G with respect to A . Recall that L is a regular language [2] and apply the UPL to obtain an integer M and a strictly increasing sequence $0 < N(1) < N(2) < \dots$ of integers such that for all integers $i > 0$ and $r \geq 0$,

$$(a_1 \cdots a_M)(a_{M+1} \cdots a_{M+N(i)})^r \in L.$$

Let $a = a_1 \cdots a_M$, $b_i = a_{M+1} \cdots a_{M+N(i)}$, and $g_i = \overline{a b_i a}^{-1}$. Our claim is that the sequence $\{g_i^{+\infty}\}$ converges to x . The point is that since $ab_i^r \in L$ for all integers $r > 0$, the path $ab_i b_i \cdots$, where b_i is repeated infinitely, is a geodesic path from 1 to $g_i^{+\infty}$ that shares the initial segment ab_i of length $M + N(i)$ with the geodesic path $a_1 a_2 \cdots$ from 1 to x . Since $N(i) \rightarrow \infty$ as $i \rightarrow \infty$, this forces the convergence of $\{g_i^{+\infty}\}$ to x . To be precise, the Gromov inner product (extended to the boundary [1]) applied to x and $g_i^{+\infty}$ yields

$$(x, g_i^{+\infty}) \geq M + N(i) - 2\delta;$$

see [1, Lemma 4.6(4), p. 51]. Hence $(x, g_i^{+\infty}) \rightarrow \infty$ as $i \rightarrow \infty$ and therefore $\{g_i^{+\infty}\}$ converges to x as desired. \square

Observe that the proof of the theorem yields the following interesting fact: an infinite geodesic path $a_1 a_2 \cdots$ based at 1 may be approximated in $\Gamma_A(G) \cup \partial G$ by traveling along an initial segment $(a_1 \cdots a_M)$ and then infinitely often repeating large segments of the path. The UPL and the proof of the theorem may be applied to verify that an infinite quasi-geodesic path may be approximated by traveling on an initial segment of the path and then repeating a subpath, chosen arbitrarily large, infinitely often.

As an application of the theorem, we give a quick proof of a known result.

Corollary. *If H is an infinite normal subgroup of the negatively curved group G , then the limit set $\Lambda(H) = \partial G \cap \text{Cl}_{G \cup \partial G}(H)$ is equal to ∂G .*

Proof. Let $x \in \partial G$ and let U be an open neighborhood of x in ∂G . Let h be an infinite order element of H [3] and choose an infinite order element g of G such that $g^{+\infty} \in U$. Since $g^{+\infty}$ is an attractive fixed point of the action of g on $G \cup \partial G$ (induced by left multiplication), there is an integer $N > 0$ such that $g^N \cdot h^{+\infty} \in U$. Since $\lim(g^N h^i g^{-N}) = g^N \cdot h^{+\infty}$ and H is normal, $g^N \cdot h^{+\infty} \in \Lambda(H)$. Since U is an arbitrary neighborhood of x , we conclude that $x \in \Lambda(H)$. \square

REFERENCES

1. J.M. Alonso, T. Brady, D. Cooper, V. Ferlini, M. Lustig, M. Mihalik, M. Shapiro, and H. Short (Editor), *Notes on word hyperbolic groups*, Group Theory from a Geometrical Viewpoint (E. Ghys, A. Haefliger, and A. Verjovsky, eds.), World Scientific, Singapore, 1991, pp. 3–63. MR **93g**:57001
2. D.B.A. Epstein, J.W. Cannon, D.F. Holt, S.V.F. Levy, M.S. Paterson, and W.P. Thurston, *Word Processing in Groups*, Jones and Bartlett, Boston, 1992. MR **93i**:20036
3. E. Ghys and P. de la Harpe (eds.), *Sur les Groupes Hyperboliques d'après Mikhael Gromov*, Progress in Math., vol. 83, Birkhäuser, 1990. MR **92f**:53050
4. M. Gromov, *Hyperbolic groups*, Essays in Group Theory (S.M. Gersten, Editor, ed.), vol. 8, Springer-Verlag, MSRI Publ, 1987, pp. 75–263. MR **89e**:20070
5. E. Swenson, *Negatively curved groups and related topics*, PhD Thesis, Brigham Young University, Provo, Utah (1993).

DEPARTMENT OF MATHEMATICS, FLORIDA STATE UNIVERSITY, TALLAHASSEE, FLORIDA 32306-3027

E-mail address: bowers@math.fsu.edu

E-mail address: ruane@math.fsu.edu