

SIMPLE PROOF OF CALABI-BERNSTEIN'S THEOREM ON MAXIMAL SURFACES

ALFONSO ROMERO

(Communicated by Christopher Croke)

Dedicated to the memory of Professor P. Bobillo

The following uniqueness result, called Calabi-Bernstein's Theorem on maximal surfaces, is well known ([Ca], [C-Y], [K], [E-R]).

Theorem. *The only entire solutions to the maximal surface equation*

$$\operatorname{div} \left(\frac{Du}{\sqrt{1 - |Du|^2}} \right) = 0, \quad |Du| < 1,$$

are affine functions.

In the references cited above, this result appears as either a particular case of some much more general theorems or stated in terms of local complex representation of the surface. However, a direct simple proof would be desirable to be easily understood for beginning researchers. The proof we present here uses only Liouville's Theorem on harmonic functions on \mathbb{R}^2 . Thus, it is simple and complex function theory is not needed. This proof is inspired by [Ch]. Roughly, the key steps of our proof are: (1) On any maximal surface there exists a positive harmonic function, which is constant if and only if the surface is totally geodesic. (2) The metric of any spacelike graph is globally conformally related to a metric g^* , which is complete when the graph is entire. (3) On any maximal graph the metric g^* is flat.

1. PRELIMINARIES

Consider the Lorentz-Minkowski space \mathbb{L}^3 with its Lorentzian metric $\langle \cdot, \cdot \rangle = dx_1^2 + dx_2^2 - dx_3^2$, given in the usual coordinate system. Let $x : M \rightarrow \mathbb{L}^3$ be a spacelike immersion of a two-dimensional manifold M in \mathbb{L}^3 . Note that M must be orientable. Let N be a globally defined unit timelike normal vector field on M . Suppose that the mean curvature of x vanishes; then M is called a maximal surface in \mathbb{L}^3 .

For each vector $a \in \mathbb{L}^3$ we can consider on M the smooth function $\langle N, a \rangle$. Let a^T be the vector field on M induced from the tangential component of a at any

Received by the editors August 11, 1994.

1991 *Mathematics Subject Classification.* Primary 53C42, 53C50.

Key words and phrases. Maximal surface, entire maximal graph, Lorentz-Minkowski space.

Research partially supported by DGICYT Grant No. PB91-0731.

point of M , that is, $a^T = a + \langle N, a \rangle N$. Standard computations give the identities

$$(1) \quad \Delta \langle N, a \rangle = \operatorname{tr}(A^2) \langle N, a \rangle,$$

$$(2) \quad \nabla \langle N, a \rangle = -A(a^T),$$

where A is the Weingarten endomorphism associated to N , and Δ and ∇ are respectively the Laplacian and the gradient relative to the induced Riemannian metric g on M .

From (2) we obtain

$$(3) \quad |\nabla \langle N, a \rangle|^2 = (1/2) \operatorname{tr}(A^2) \{ \langle N, a \rangle^2 + \langle a, a \rangle \}.$$

Now, choose $a \neq 0$ satisfying $\langle a, a \rangle = 0$. Note that in this case $\langle N, a \rangle$ never vanishes and thus we may assume $\langle N, a \rangle > 0$. From (1) and (3) we get

$$(4) \quad \Delta(1/\langle N, a \rangle) = 0.$$

On the other hand, if we choose now $b \in \mathbb{L}^3$ such that $\langle b, b \rangle = -1$ and $\langle N, b \rangle > 0$, then, using again (1) and (3) we obtain

$$(5) \quad \Delta \log(\langle N, b \rangle + 1) = (1/2) \operatorname{tr}(A^2)$$

which means, taking into account the Gauss equation $2K = \operatorname{tr}(A^2)$ for x , that the metric $g^* = (\langle N, b \rangle + 1)^2 g$ on M is flat.

2. PROOF OF THE THEOREM

We begin by observing that on a spacelike graph in \mathbb{L}^3 of any entire function $u = u(x, y)$, $|Du| < 1$, the metric $g' = (1/(1 - |Du|^2))g$ is complete. This easily follows from the inequality $L' \geq (1/\sqrt{2})L_0$, where L' (resp. L_0) denotes the length of a curve on the graph with respect to g' (resp. the metric induced from the usual Riemannian one of \mathbb{R}^3). Now take $b = (0, 0, -1)$ so that $\langle N, b \rangle = 1/\sqrt{1 - |Du|^2}$. Consider now the Riemannian metric g^* introduced above. This metric is complete on a spacelike graph of any entire function. Assume the graph is maximal. On the other hand, from (5) we see that g^* is flat. Using Cartan's Theorem we have a global isometry from the Euclidean plane \mathbb{R}^2 onto the maximal entire graph endowed with the metric g^* . The invariance of harmonic functions under conformal changes of metric and this isometry permit us to induce $1/\langle N, a \rangle$ on a positive harmonic function on \mathbb{R}^2 which must be constant by Liouville's Theorem. Hence $\langle N, a \rangle$ is constant on the graph, which implies, using (1), that the maximal graph is totally geodesic.

3. REMARK

Consider as in §1, a maximal surface M in \mathbb{L}^3 . The metric g^* on M is also complete if the induced metric g is complete. Thus, the same reasoning as in the previous section gives a proof of the well-known parametric version of this Theorem: "The only complete maximal surfaces in \mathbb{L}^3 are the spacelike planes."

REFERENCES

- [Ca] E. Calabi, *Examples of Bernstein problems for some non-linear equations*, Proc. Sympos. Pure Math. **15** (1970), 223–230. MR **41**:8806
- [Ch] S. S. Chern, *Simple proofs of two theorems on minimal surfaces*, Enseign. Math. **15** (1969), 53–61. MR **39**:7516
- [C-Y] S. T. Cheng, S. T. Yau, *Maximal spacelike hypersurfaces in the Lorentz-Minkowski space*, Ann. of Math. **104** (1976), 407–419. MR **55**:4063
- [E-R] F. J. M. Estudillo, A. Romero, *Generalized maximal surfaces in Lorentz-Minkowski space \mathbb{L}^3* , Math. Proc. Camb. Phil. Soc. **111** (1992), 515–524. MR **93b**:53010
- [K] O. Kobayashi, *Maximal surfaces in the 3-dimensional Minkowski space \mathbb{L}^3* , Tokyo J. Math. **6** (1983), 297–309. MR **85d**:53003

DEPARTAMENTO DE GEOMETRIA Y TOPOLOGIA, FACULTAD DE CIENCIAS, UNIVERSIDAD DE GRANADA, 18071-GRANADA
E-mail address: `aromero@ugr.es`