

## A COUNTEREXAMPLE TO THE DIFFERENTIABILITY OF THE BERGMAN KERNEL FUNCTION

SO-CHIN CHEN

(Communicated by Eric Bedford)

ABSTRACT. In this paper we prove the following main result. Let  $D$  be a smoothly bounded pseudoconvex domain in  $\mathbf{C}^n$  with  $n \geq 2$ . Suppose that there exists a complex variety sitting in the boundary  $bD$ ; then we have

$$K_D(z, w) \notin C^\infty(\overline{D} \times \overline{D} - \Delta(bD)).$$

In particular, the Bergman kernel function associated with the Diederich-Fornaess worm domain is not smooth up to the boundary in joint variables off the diagonal of the boundary.

### 1. INTRODUCTION

In several complex variables it has been shown that the Bergman kernel function is closely related to the boundary regularity of holomorphic mappings between two domains. Therefore, it is always a fundamental question to investigate the boundary regularity of the Bergman kernel function associated with a smoothly bounded pseudoconvex domain.

Historically, it was first proved by Kerzman [10], based on Kohn's work, that the Bergman kernel function associated to a smoothly bounded strictly pseudoconvex domain  $D$  can be extended smoothly to  $\overline{D} \times \overline{D} - \Delta(bD)$ , where  $\Delta(bD) = \{(z, z) | z \in bD\}$ . Later it was generalized independently by Bell [2] and Boas [3] to the following.

**Theorem 1.1.** *Let  $D$  be a smoothly bounded pseudoconvex domain in  $\mathbf{C}^n$ . If  $\Gamma_1$  and  $\Gamma_2$  are disjoint open subsets of  $bD$  consisting of points of finite type in the sense of D'Angelo [8], then the Bergman kernel function associated to  $D$  extends smoothly to  $\Gamma_1 \times \Gamma_2$ .*

**Theorem 1.2.** *Let  $D$  be a smoothly bounded pseudoconvex domain in  $\mathbf{C}^n$ . Suppose that condition  $R$  holds on  $D$ . Let  $\Gamma_1$  and  $\Gamma_2$  be two disjoint open subsets of  $bD$  such that  $\Gamma_1$  consists of points of finite type. Then the Bergman kernel function associated to  $D$  extends smoothly to  $\Gamma_1 \times \Gamma_2$ .*

Here condition  $R$  means that the Bergman projection  $P$ , the orthogonal projection from  $L^2(D)$  onto the closed subspace  $H^2(D)$  of square-integrable holomorphic functions, maps  $C^\infty(\overline{D})$  continuously into itself.

Based on the above theorems it is natural to conjecture that a similar extension phenomenon might probably hold on any smoothly bounded pseudoconvex domain in  $\mathbf{C}^n$ . However, in this article we are going to prove the following main result.

---

Received by the editors December 1, 1994.

1991 *Mathematics Subject Classification.* Primary 32H10.

**Main Theorem.** *Let  $D$  be a smoothly bounded pseudoconvex domain in  $\mathbf{C}^n$  with  $n \geq 2$ . Suppose that there exists a complex variety sitting in the boundary  $bD$ ; then we have*

$$K_D(z, w) \notin C^\infty(\overline{D} \times \overline{D} - \Delta(bD)),$$

where  $\Delta(bD) = \{(z, z) | z \in bD\}$ .

## 2. TWO EXAMPLES

In this section we will present two examples. First we fix a smooth real-valued function  $\lambda : \mathbf{R} \rightarrow \mathbf{R}$  as in [9] with the following properties:

- (i)  $\lambda(x) = 0$  if  $x \leq 0$ .
- (ii)  $\lambda(x) > 0$  if  $x > 1$ .
- (iii)  $\lambda''(x) \geq 100\lambda'(x)$  for all  $x$ .
- (iv)  $\lambda''(x) > 0$  if  $x > 0$ .
- (v)  $\lambda'(x) > 100$  if  $\lambda(x) > \frac{1}{2}$ .

**Example A.** For any  $r > 1$ , define the domain  $\Omega_r$  by

$$\Omega_r = \{(z, w) \in \mathbf{C}^2 | \rho_r(z, w) < 0\},$$

where  $\rho_r(z, w) = |z|^2 - 1 + \lambda(|w|^2 - r^2)$ .

Then we have

**Lemma 2.1.** *The domain  $\Omega_r$  defined as above is a smoothly bounded pseudoconvex domain in  $\mathbf{C}^2$  satisfying that*

- (a)  $\Omega_r$  is convex.
- (b) Condition  $R$  holds on  $\Omega_r$ .
- (c)  $\Omega_r$  is strictly pseudoconvex everywhere except on the set

$$M_r = \{(z, w) \in \mathbf{C}^2 | |z| = 1 \text{ and } 0 \leq |w| \leq r\}.$$

Condition  $R$  on  $\Omega_r$  follows from either (a) or the fact that  $\Omega_r$  is Reinhardt. For instance, see Chen [6], [7], Boas and Straube [4] or Straube [12]. Obviously, the boundary contains a family of complex discs parametrized by the unit circle. Hence by our main theorem we have

$$K_{\Omega_r}(z, w) \notin C^\infty(\overline{\Omega_r} \times \overline{\Omega_r} - \Delta(b\Omega_r)).$$

But we would like to provide an elementary proof of the main theorem for  $\Omega_r$  here which gives a more geometric insight into the problem.

Let  $\Delta_r = \{w \in \mathbf{C} | |w| < \sqrt{r^2 + 1}\}$ , and denote by  $\Delta$  the unit disc in the complex plane. Put  $D_r = \Delta \times \Delta_r$ .  $D_r$  is a pseudoconvex domain in  $\mathbf{C}^2$ , and obviously we have  $\Omega_r \subseteq D_r$ . It is also clear that  $\Lambda = \{z^m w^n | m, n \in \mathbf{N} \cup \{0\}\}$  forms a complete orthogonal basis for both  $H^2(\Omega_r)$  and  $H^2(D_r)$ . So we have

$$c_{mn}^r = \int_{\Omega_r} |z^m w^n|^2 d\sigma \leq \int_{D_r} |z^m w^n|^2 d\sigma = d_{mn}^r \text{ for all } m, n.$$

Both equalities are definitions. Now consider any point sequence  $\{z_j\}_{j=1}^\infty$  in  $\Delta$  such that  $\{z_j\}$  approaches a boundary point, say  $z_0$ . Pick any two positive numbers  $p$  and  $q$  such that  $0 < p < q < r$ . So now we have two point sequences  $\{(z_j, p)\}$  and  $\{(z_j, q)\}$  in both  $\Omega_r$  and  $D_r$  which approach the boundary points  $(z_0, p)$  and  $(z_0, q)$  respectively. Clearly we have  $(z_0, p) \neq (z_0, q)$ . Next, by using the explicit formula

for the Bergman kernel function  $K_{\Delta}(z, w)$  on the unit disc in the complex plane, we obtain

$$\begin{aligned} K_{\Omega_r}((z_j, p), (z_j, q)) &= \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{1}{c_{mn}^r} |z_j|^{2m} \cdot (pq)^n \\ &\geq \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{1}{d_{mn}^r} |z_j|^{2m} \cdot (pq)^n \\ &= K_{\Delta}(z_j, z_j) K_{\Delta_r}(p, q) \\ &= \frac{1}{\pi} \cdot \frac{1}{(1 - |z_j|^2)^2} K_{\Delta_r}(p, q). \end{aligned}$$

It follows that

$$\lim_{j \rightarrow \infty} K_{\Omega_r}((z_j, p), (z_j, q)) = \infty.$$

Hence the Bergman kernel function  $K_{\Omega_r}(z, w)$  is not bounded near  $z = (z_0, p)$  and  $w = (z_0, q)$ . In particular, it shows that  $K_{\Omega_r}(z, w) \notin C^{\infty}(\overline{\Omega_r} \times \overline{\Omega_r} - \Delta(b\Omega_r))$ . This completes the proof of the main theorem on  $\Omega_r$ .

**Example B** (Diederich-Fornaess worm domain). Now define the domain  $\Omega_r$ , for any  $r > 1$ , by

$$\Omega_r = \{(z, w) \in \mathbf{C}^2 \mid \rho_r(z, w) < 0\},$$

where  $\rho_r(z, w) = |z + \exp(in|w|^2)|^2 - 1 + \lambda(\frac{1}{|w|^2} - 1) + \lambda(|w|^2 - r^2)$ . Then we obtain the famous Diederich-Fornaess worm domain [9]. These domains  $\Omega_r$ s are smoothly bounded pseudoconvex in  $\mathbf{C}^2$  and are strictly pseudoconvex everywhere except on the annulus

$$A_r = \{(z, w) \in \mathbf{C}^2 \mid z = 0 \text{ and } 1 \leq |w| \leq r\}.$$

Therefore, again by our main theorem we see that

$$K_{\Omega_r}(z, w) \notin C^{\infty}(\overline{\Omega_r} \times \overline{\Omega_r} - \Delta(b\Omega_r)).$$

We should point out that it has been shown in Barrett [1] that the Bergman projection on these domains does not preserve the Sobolev  $k$ -space  $W^k(\Omega_r)$  if  $k$  is large enough. However, it is still not known whether the condition  $R$  holds on  $\Omega_r$  or not.

### 3. PROOF OF THE MAIN THEOREM

Let  $D$  be a smoothly bounded pseudoconvex domain in  $\mathbf{C}^n$  with  $n \geq 2$ , and let  $V$  be a complex variety sitting in the boundary  $bD$ . Let  $p \in V$  be a regular point of  $V$ . Denote by  $\vec{n}$  the unit outward normal at  $p$ . Since the boundary of  $D$  is smooth, there exists small  $\delta, \varepsilon_0 > 0$  such that  $w - \varepsilon \vec{n} \in D$  for all  $w \in bD \cap B(p; \delta)$  and all  $0 < \varepsilon < \varepsilon_0$ . Now let us consider a small complex disc  $\overline{\Delta} \subseteq bD \cap B(p; \delta) \cap V$  centered at  $p$ . More precisely,  $\Delta$  is the holomorphic embedding of the unit disc in the complex plane in the boundary, and the origin is mapped to  $p$ .

Suppose now that  $K_D(z, w) \in C^{\infty}(\overline{D} \times \overline{D} - \Delta(bD))$ . Then we have

$$\sup_{w \in b\Delta} |K_D(p, w)| \leq M < +\infty,$$

for some  $M > 0$ . On the other hand, it was proved in P. Pflug [11] that

$$\lim_{\varepsilon \rightarrow 0} K_D(p - \varepsilon \vec{n}, p - \varepsilon \vec{n}) = +\infty.$$

Then by the maximum modulus principle we get

$$\sup_{w \in b\Delta_\varepsilon} |K_D(p - \varepsilon \vec{n}, w)| \geq K_D(p - \varepsilon \vec{n}, p - \varepsilon \vec{n}),$$

where  $\Delta_\varepsilon = \Delta - \varepsilon \vec{n} \subseteq D$ . It follows that

$$\sup_{w \in b\Delta} |K_D(p, w)| = \lim_{\varepsilon \rightarrow 0} \sup_{w \in b\Delta_\varepsilon} |K_D(p - \varepsilon \vec{n}, w)| = +\infty.$$

This gives the desired contradiction, and the proof of the main theorem is now completed.

#### 4. CONCLUDING REMARK

In view of the main theorem proved above, it is natural to ask: How does the Bergman kernel function  $K_D(z, w)$  behave in joint variables near the boundary, if the domain  $D$  satisfies property  $(P)$  introduced in Catlin [5]? So far we have no clue to answer this question.

#### REFERENCES

1. D. Barrett, *Behavior of the Bergman projection on the Diederich-Fornaess worm*, Acta Math., **168**, (1992), 1–10. MR **93c**:32033
2. S. Bell, *Differentiability of the Bergman kernel and pseudo-local estimates*, Math. Z., **192**, (1986), 467–472. MR **87i**:32034
3. H. P. Boas, *Extension of Kerzman's theorem on differentiability of the Bergman kernel function*, Indiana Univ. Math. J., **36**, (1987), 495–499. MR **88j**:32028
4. H. P. Boas and E. J. Straube, *Sobolev estimates for the  $\bar{\partial}$ -Neumann operator on domains in  $\mathbf{C}^n$  admitting a defining function that is plurisubharmonic on the boundary*, Math. Z., **206**, (1991), 81–88. MR **92b**:32027
5. D. Catlin, *Global regularity for the  $\bar{\partial}$ -Neumann problem*, Proc. Sympos. Pure Math., vol. 41, Amer. Math. Soc., Providence, RI, 1984. MR **85j**:32033
6. S. C. Chen, *Global regularity of the  $\bar{\partial}$ -Neumann problem on circular domains*, Math. Ann., **285**, (1989), 1–12. MR **90i**:32028
7. ———, *Global regularity of the  $\bar{\partial}$ -Neumann problem in dimension two*, Proc. Sympos. Pure Math., vol. 52, Part III, Amer. Math. Soc., Providence, RI, 1991, 55–61. MR **92h**:32027
8. J. D'Angelo, *Real hypersurfaces, order of contact, and applications*, Ann. of Math., **115**, (1982), 615–637. MR **84a**:32027
9. K. Diederich and J. E. Fornaess, *Pseudoconvex domains: an example with nontrivial Neberhülle*, Math. Ann., **225**, (1977), 275–292. MR **55**:3320
10. N. Kerzman, *The Bergman kernel function. Differentiability at the boundary*, Math. Ann., **195**, (1972), 149–158. MR **45**:3762
11. P. Pflug, *Quadratintegrable holomorphe Funktionen und die Serre-Vermutung*, Math. Ann., **216**, (1975), 285–288. MR **52**:3599
12. E. J. Straube, *Exact regularity of Bergman, Szegő and Sobolev space projections in non pseudoconvex domains*, Math. Z., **192**, (1986), 117–128. MR **87k**:32045

INSTITUTE OF APPLIED MATHEMATICS, NATIONAL TSING HUA UNIVERSITY, HSINCHU 30043, TAIWAN, REPUBLIC OF CHINA

*E-mail address*: scchen@am.nthu.edu.tw