

## ON THE WEAK UNIFORM CONVEXITY OF $Q(R)$

SHEN YU-LIANG

(Communicated by Albert Baernstein II)

ABSTRACT. We will discuss the geometry of the unit sphere in the Banach space of integrable holomorphic quadratic differentials on a Riemann surface and answer some questions posed by L.R. Goldberg (Proc. Amer. Math. Soc. **118** (1993), 1179–1185).

### 1

For a given hyperbolic Riemann surface  $R$ , that is, covered by the unit disk  $\Delta$ , we denote by  $Q(R)$  the Banach space of all holomorphic quadratic differentials  $\phi(z) dz^2$  on  $R$  with the  $L^1$  norm

$$\|\phi\| = \iint_R |\phi(z)| dx dy < \infty.$$

It is well known that  $Q(R)$  is of finite dimension if and only if  $R$  is of finite type. Let  $S(R) = \{\phi \in Q(R) : \|\phi\| = 1\}$  denote the unit sphere in  $Q(R)$ . We are going to study the geometry of  $S(R)$ , answering some questions of Goldberg [5].

In general, if  $X$  is a Banach space, then  $X^*$  is the Banach space of bounded real-linear functionals  $l : X \rightarrow \mathbf{R}$ . Now define  $X$  to be smooth if for every unit vector  $x$  in  $X$  there is a unique  $l$  in  $X^*$  with  $l(x) = \|l\| = 1$ . Note that our definition of smoothness is taken from Diestel [1] and differs from the definition in Goldberg [5]; however, Diestel proved that the two definitions are equivalent.

It is well known that  $Q(R)$  is smooth (see [2] or [4]). If  $\psi \in S(R)$ , the unique functional  $\psi_*$  in  $Q(R)^*$  with  $\psi_*(\psi) = \|\psi_*\| = 1$  is

$$\psi_*(\phi) = \operatorname{Re} \iint_R \phi(z) \frac{\overline{\psi(z)}}{|\psi(z)|} dx dy, \quad \phi \in Q(R).$$

It is also well known that  $Q(R)$  is strictly convex, meaning (as usual) that every point of  $S(R)$  is an extreme point of the closed unit ball.

---

Received by the editors July 26, 1994 and, in revised form, December 22, 1994.

1991 *Mathematics Subject Classification*. Primary 30F30, 30C70, 30F60.

*Key words and phrases*. Quadratic differential, weak uniform convexity, Hamilton sequence.

The author was supported in part by Jiangsu Provincial Natural Science Foundation.

A Banach space  $X$  is said to be uniformly convex if for all sequences  $(x_n)$  and  $(y_n)$  in the closed unit ball,  $\|x_n + y_n\| \rightarrow 2$  implies  $\|x_n - y_n\| \rightarrow 0$ . It is obvious that a finite dimensional strictly convex Banach space is uniformly convex (in particular  $Q(R)$  is uniformly convex if  $R$  has finite type), but the infinite dimensional case is more complicated. In her study of the convexity of  $Q(\Delta)$ , Goldberg uses two weaker notions.

Following Goldberg [5] we say that  $X$  is uniformly convex at the unit vector  $x$  if for all sequences  $(x_n)$  in the closed unit ball,  $\|x_n + x\| \rightarrow 2$  implies  $x_n \rightarrow x$ . If  $X$  is smooth,  $x$  is a unit vector, and  $x_*$  is the unique functional of norm one such that  $x_*(x) = 1$ , we say that  $X$  is weakly uniformly convex at  $x$  if for all sequences  $(x_n)$  in the unit sphere,  $x_*(x_n) \rightarrow 1$  implies  $x_n \rightarrow x$ .

If  $x_*(x_n) \rightarrow 1$ , then  $x_*(x_n + x) \rightarrow 2$ , so  $\|x_n + x\| \rightarrow 2$ . Therefore uniform convexity at  $x$  implies weak uniform convexity at  $x$ . Uniform convexity trivially implies uniform convexity at each unit vector  $x$ , so if  $R$  is a finite Riemann surface, then  $Q(R)$  is weakly uniformly convex at every point of  $S(R)$ . That solves Problem 3 of Goldberg [5].

To study the weak uniform convexity of  $Q(R)$  in the infinite dimensional case we use Hamilton sequences. Let  $l$  be a linear functional on  $Q(R)$ . By definition, a Hamilton sequence for  $l$  is a sequence  $(\phi_n)$  in  $S(R)$  such that  $l(\phi_n) \rightarrow \|l\|$ . The Hamilton sequence  $(\phi_n)$  is called degenerate if  $\phi_n \rightarrow 0$  locally uniformly in  $R$ .

Now we can state our main result.

**Theorem 1.** *The Banach space  $Q(R)$  is weakly uniformly convex at  $\psi \in S(R)$  if and only if the linear functional  $\psi_*$  does not possess any degenerate Hamilton sequence.*

*Remark.* Theorem 1 gives a characterization of the flat points in  $S(R)$  where  $Q(R)$  is not weakly uniformly convex; this gives an answer to Goldberg's Problem 2. A solution to her Problem 1 will be given in §3.

The paper is organized as follows: We prove Theorem 1 in §2. In §3, we obtain some consequences by using Strebel's Frame Mapping Condition from the theory of quasiconformal mappings. Finally, in §4, we discuss the uniform convexity of the Banach space  $Q(R)$ .

## 2

We first point out that Hamilton sequences obey a well known principle which is implicit in Strebel [7] and also Harrington-Ortel [6]. The principle is this: If  $l$  is a nontrivial linear functional on  $Q(R)$ , then either  $l$  has a degenerating Hamilton sequence or else every Hamilton sequence converges (in norm) to the unique  $\phi$  in  $S(R)$  such that  $l(\phi) = \|l\|$ . These two possibilities are obviously mutually exclusive.

Now we give the proof of Theorem 1.

Assume first that  $Q(R)$  is weakly uniformly convex at  $\psi \in S(R)$ . Let  $(\phi_n)$  be any Hamilton sequence for  $\psi_*$ . Since  $\psi_*(\phi_n) \rightarrow 1$ , the weak uniform convexity implies that  $\phi_n \rightarrow \psi$ , so every Hamilton sequence for  $\psi_*$  converges in norm to  $\psi$ , and no Hamilton sequence for  $\psi_*$  is degenerate.

Conversely, suppose  $\psi \in S(R)$  and no Hamilton sequence for  $\psi_*$  is degenerate. We shall prove that  $Q(R)$  is weakly uniformly convex at  $\psi$ . Let  $(\phi_n)$  be a sequence in  $S(R)$  such that  $\psi_*(\phi_n) \rightarrow 1$ . Since  $(\phi_n)$  is a Hamilton sequence for  $\psi_*$ , the above-stated principle implies that  $\phi_n \rightarrow \psi$  as required.

## 3

For an extremal quasiconformal mapping  $f : R \rightarrow R'$ , we denote by  $H(f)$  the boundary dilatation of  $f$  (see Strebel [8] or Gardiner [4, §6.8] for the definition) and by  $K_0(f)$  its maximal dilatation. For a given  $\psi \in S(R)$ , choose  $k$  in the open interval  $(0, 1)$  and denote by  $f_\psi$  the extremal quasiconformal mapping on  $R$  whose Beltrami coefficient is  $k\bar{\psi}/|\psi|$ .

**Theorem 2.** *If  $\psi \in S(R)$ , then  $Q(R)$  is weakly uniformly convex at  $\psi$  if and only if  $H(f_\psi) < K_0(f_\psi)$ .*

*Proof.* By Strebel's Frame Mapping Condition (see [4] or [8]) and a recent result of Earle-Li [3], if  $\psi \in S(R)$ , then  $H(f_\psi) < K_0(f_\psi)$  if and only if the linear functional  $k\psi_*$  defined by the Beltrami differential of  $f_\psi$  has no degenerate Hamilton sequence, that is,  $\psi_*$  has no degenerate Hamilton sequence. Therefore,  $Q(R)$  is weakly uniformly convex at  $\psi$  if and only if  $H(f_\psi) < K_0(f_\psi)$ , by Theorem 1.

**Corollary.** *Let  $\psi \in S(\Delta)$  have a holomorphic extension to a neighbourhood of the closed disk with no zeroes on the unit circle. Then  $Q(\Delta)$  is weakly uniformly convex at  $\psi$ .*

*Proof.* Under the hypothesis, the restriction of  $f_\psi$  to a neighbourhood of the unit circle is a real analytic diffeomorphism, so  $H(f_\psi) = 1 < K_0(f_\psi)$ .

*Remark.* The Corollary applies in particular to  $\psi = \frac{n+2}{2\pi} z^n dz^2 \in S(\Delta)$  and solves Goldberg's Problem 1.

## 4

Goldberg [5] shows that  $Q(\Delta)$  is nowhere uniformly convex. For any Riemann surface  $R$ , we have

**Theorem 3.** (i) *If  $\dim Q(R) < \infty$ ,  $Q(R)$  is uniformly convex.*

(ii) *If  $\dim Q(R) = \infty$ ,  $Q(R)$  is nowhere uniformly convex.*

*Proof.* We have already remarked in §1 that since  $Q(R)$  is strictly convex, it is uniformly convex whenever it is finite dimensional.

Now we suppose that  $\dim Q(R) = \infty$ . We modify Goldberg's discussion (see [5]). Fix  $\psi \in S(R)$  and  $\varepsilon > 0$ . Choose some compact set  $K \subset R$  such that  $\iint_K |\psi(z)| dx dy > 1 - \varepsilon$ . Since  $\dim Q(R) = \infty$ , there must exist a degenerate sequence  $(\phi_n)$  in  $S(R)$ , that is,  $\phi_n \rightarrow 0$  locally uniformly in  $R$ . When  $n$  is sufficiently large,  $\iint_{S-K} |\phi_n(z)| dx dy > 1 - \varepsilon$ . Therefore,

$$\begin{aligned} \|\phi_n \pm \psi\| &= \iint_K |\phi_n \pm \psi| dx dy + \iint_{S-K} |\phi_n \pm \psi| dx dy \\ &\geq \iint_K |\psi| dx dy - \iint_K |\phi_n| dx dy + \iint_{S-K} |\phi_n| dx dy - \iint_{S-K} |\psi| dx dy \geq 2 - 4\varepsilon. \end{aligned}$$

Thus,  $Q(R)$  is not uniformly convex at  $\psi \in S(R)$ .

## ACKNOWLEDGEMENT

I wish to thank Prof. Li Zhong for his encouragement and help. I also wish to express my gratitude to the referee for his useful advice and for calling my attention to the result of C.J. Earle and Li Zhong.

## REFERENCES

1. J. Diestel, *Geometry of Banach spaces—selected topics*, Springer-Verlag, New York, 1975. MR **57**:1079
2. C.J. Earle, *On holomorphic cross-sections in Teichmüller spaces*, Duke Math. J. **36** (1969), 409–416. MR **40**:7442
3. C.J. Earle and Li Zhong, *Isometrically embedded polydisks in infinite dimensional Teichmüller spaces*, to appear.
4. F.P. Gardiner, *Teichmüller theory and quadratic differentials*, Wiley-Interscience, New York, 1987. MR **88m**:32044
5. L.R. Goldberg, *On the shape of the unit sphere in  $Q(\Delta)$* , Proc. Amer. Math. Soc. **118** (1993), 1179–1185. MR **93m**:46019
6. A. Harrington and M. Ortel, *The dilatation of an extremal quasiconformal mapping*, Duke Math. J. **43** (1976), 533–544. MR **54**:13074
7. K. Strebel, *On quadratic differentials and extremal quasiconformal mappings*, Proceedings Int. Congr. Math. Vancouver (1974), 223–227. MR **58**:22549
8. K. Strebel, *On the existence of extremal Teichmüller mappings*, Journal d'Analyse Math. **30** (1976), 464–480. MR **55**:12912
9. K. Strebel, *Extremal quasiconformal mappings*, Resulate Math. **10** (1986), 168–210. MR **88a**:30048

DEPARTMENT OF MATHEMATICS, SUZHOU UNIVERSITY, SUZHOU 215006, PEOPLE'S REPUBLIC OF CHINA