

THE LU QI-KENG CONJECTURE FAILS GENERICALLY

HAROLD P. BOAS

(Communicated by Eric Bedford)

ABSTRACT. The bounded domains of holomorphy in \mathbf{C}^n whose Bergman kernel functions are zero-free form a nowhere dense subset (with respect to a variant of the Hausdorff distance) of all bounded domains of holomorphy.

A domain in \mathbf{C}^n is called a *Lu Qi-Keng domain* if its Bergman kernel function has no zeroes. Lu Qi-Keng [11] raised the question of which domains, besides the ball and the polydisc, have this property. A motivation for the question is that the vanishing of the Bergman kernel function obstructs the global definition of Bergman representative coordinates. Over the years since Lu Qi-Keng's paper appeared, various versions of a *Lu Qi-Keng conjecture* have been mooted to the effect that all domains, or most domains, or all domains satisfying some geometrical hypothesis, are Lu Qi-Keng domains.

In the complex plane \mathbf{C}^1 , a bounded domain with smooth boundary is a Lu Qi-Keng domain if and only if it is simply connected [16] (and thus biholomorphically equivalent to the disc). I have given a counterexample [1] showing that no analogous topological characterization of Lu Qi-Keng domains can hold in higher dimensions: there exists (in \mathbf{C}^2 , and similarly in \mathbf{C}^n for $n > 2$) a bounded, strongly pseudoconvex, contractible domain with C^∞ regular boundary whose Bergman kernel function does have zeroes.

In this note, I show that the Lu Qi-Keng domains of holomorphy may be viewed as exceptional: they form a nowhere dense set with respect to a suitable topology. Thus, contrary to former expectations, it is the normal situation for the Bergman kernel function of a domain to have zeroes.

To formulate the result precisely, I need a metric on bounded open sets. Since I impose no restriction on the regularity of the boundaries of the sets, some variant of the Hausdorff metric will be appropriate. The Hausdorff distance \mathcal{H} is normally defined for nonempty, bounded, closed sets by the property that $\mathcal{H}(A, B) < \epsilon$ if and only if each point of A has Euclidean distance less than ϵ from some point of B , and vice versa.

After the seminal paper of Ramadanov [12], it is clear in the context of the Bergman kernel function that if a sequence of open sets $\{\Omega_j\}$ is going to be said to converge to an open set Ω , then every compact subset of Ω should eventually be contained in Ω_j . It is less clear what requirement should be imposed if the Ω_j contain points outside of Ω . The example [15, p. 39], [13, p. 280] of decreasing

Received by the editors December 10, 1994.

1991 *Mathematics Subject Classification*. Primary 32H10.

This research was partially supported by NSF grant number DMS-9203514.

concentric disks in the complex plane converging to a disk with a slit removed shows that it is inadequate to require merely that for every open neighborhood of the closure $\overline{\Omega}$, eventually Ω_j is contained in the neighborhood.

I shall consider two different notions of convergence of open sets in \mathbf{C}^n . Both have the property that if $\Omega_j \rightarrow \Omega$, then the Ω_j eventually swallow every compact subset of Ω . However, they differ in what they require about the sets $\Omega_j \setminus \Omega$.

First I define a metric ρ_1 on bounded, nonempty, open sets via $\rho_1(U, V) = \mathcal{H}(\overline{U}, \overline{V}) + \mathcal{H}(\partial U, \partial V)$. If $\rho_1(\Omega_j, \Omega) \rightarrow 0$, then the Ω_j eventually swallow every compact subset of Ω and are eventually swallowed by every open neighborhood of $\overline{\Omega}$. The converse holds when Ω equals the interior of its closure, but not in general. By requiring that both the closures and the boundaries converge, convergence in the metric ρ_1 eliminates examples like the one above involving slits or punctures in the limit domain.

The metric ρ_1 can also be thought of in terms of functions. Define the distance function d_U of an open set U via $d_U(z) = \text{dist}(z, \mathbf{C}^n \setminus U)$, where dist denotes the Euclidean distance. Then $\Omega_j \rightarrow \Omega$ according to the metric ρ_1 if and only if the continuous functions d_{Ω_j} converge uniformly on \mathbf{C}^n to d_Ω and the $d_{\mathbf{C}^n \setminus \overline{\Omega_j}}$ converge uniformly to $d_{\mathbf{C}^n \setminus \overline{\Omega}}$.

In some contexts—one will appear below—it is useful to relax the hypothesis on how the sets $\Omega_j \setminus \Omega$ behave. They could be required to shrink in volume (Lebesgue $2n$ -dimensional measure), but not necessarily in terms of Euclidean distance from Ω . I therefore introduce a second metric ρ_2 on bounded open sets via $\rho_2(U, V) = \text{vol}(U \setminus V) + \text{vol}(V \setminus U) + \sup_{z \in \mathbf{C}^n} |d_U(z) - d_V(z)|$. Convergence of Ω_j to Ω in this metric allows Ω_j to have a long thin tail whose width shrinks to zero but whose length does not shrink.

So far I have not assumed that the open sets in question are connected. It is easy to see that the Bergman kernel function $K(w, z)$ for a disconnected open set is identically equal to zero if w and z are in different connected components, while if the points are in the same connected component, then $K(w, z)$ equals the Bergman kernel function of that component. I will say that a (possibly) disconnected, bounded, nonempty, open set is a *Lu Qi-Keng open set* if its Bergman kernel function has no zeroes when the two variables are in the same connected component.

Theorem. *The Lu Qi-Keng open sets are nowhere dense in each of the following metric spaces, where the metric is ρ_1 :*

- (1) *the bounded pseudoconvex open sets;*
- (2) *the bounded connected pseudoconvex open sets (domains of holomorphy);*
- (3) *the bounded strongly pseudoconvex open sets;*
- (4) *the bounded connected strongly pseudoconvex open sets.*

If one considers only open sets of Euclidean diameter less than some fixed constant M , then the same assertion holds when the metric is taken to be ρ_2 .

I will base the proof of the theorem on the following two folklore lemmas. The ideas of the proofs are all in the literature, but since I do not know a reference for precisely these formulations, I will indicate proofs of the lemmas after the proof of the theorem.

Stability lemma for the Bergman kernel function. *Let $\{\Omega_j\}$ be a sequence of bounded pseudoconvex open sets that converges, in the sense of either ρ_1 or ρ_2 , to a nonempty bounded open set Ω ; in the case of ρ_2 , assume also that the Ω_j have*

uniformly bounded diameters (this is automatic in the case of ρ_1). Suppose U is a connected component of Ω that has C^∞ regular boundary and that is separated from the rest of Ω (that is, the closure of U is disjoint from the closure of $\Omega \setminus U$). Then the Bergman kernel functions of the Ω_j converge to the Bergman kernel function of U uniformly on compact subsets of $U \times U$.

In the statement of the stability lemma, pseudoconvexity of the limit set Ω is automatic: the limit function $-\log d_\Omega$ inherits plurisubharmonicity from the functions $-\log d_{\Omega_j}$, which converge uniformly on compact subsets of Ω .

A special case of considerable interest is when the Ω_j and Ω are all bounded, connected, pseudoconvex domains with C^∞ regular boundaries. The lemma then says that if the Ω_j converge to Ω , in the sense that Ω_j eventually swallows every compact subset of Ω and the volume of $\Omega_j \setminus \Omega$ tends to zero, then the Bergman kernel functions of the Ω_j converge uniformly on compact subsets of $\Omega \times \Omega$ to the Bergman kernel function of Ω .

The C^∞ regularity hypothesis in the lemma can be reduced to C^2 regularity, but I shall not prove this here.

I take the name of the second lemma from [10, Chap. 5, Exercise 21].

Barbell lemma. *Suppose G_1 and G_2 are bounded, connected, strongly pseudoconvex domains in \mathbf{C}^n with C^∞ regular boundaries and with disjoint closures. Let γ be a smooth curve (that is, a C^∞ embedding of $[0, 1]$ into \mathbf{C}^n) joining a boundary point of G_1 to a boundary point of G_2 , and otherwise disjoint from the closures of G_1 and G_2 , and let V be an arbitrary neighborhood in \mathbf{C}^n of the curve γ . Then there exists a bounded, connected, strongly pseudoconvex domain Ω with C^∞ regular boundary such that Ω is contained in $G_1 \cup G_2 \cup V$, and Ω coincides with $G_1 \cup G_2$ outside V .*

When G_1 and G_2 are balls of equal size, and γ is the shortest line segment joining them, then the domain Ω is a “barbell”, or dumbbell.

The C^∞ regularity can be changed everywhere in the statement of the lemma to C^k regularity, where k is any integer greater than or equal to 2.

Proof of the theorem. I have not claimed that the bounded pseudoconvex open sets which fail to be Lu Qi-Keng form an open set in either of the metrics ρ_1 or ρ_2 , and I do not know whether or not this is the case for sets with irregular boundaries. However, if Ω is, for example, a bounded strongly pseudoconvex open set with C^∞ regular boundary, and the Bergman kernel function of Ω has zeroes on some connected component Ω_0 , then there is a ρ_1 neighborhood of Ω containing no pseudoconvex Lu Qi-Keng open set. Indeed, if a sequence of pseudoconvex open sets converges to Ω in the metric ρ_1 , then the corresponding Bergman kernel functions converge on Ω_0 to the Bergman kernel function of Ω_0 by the stability lemma, and by Hurwitz’s theorem these approximating Bergman kernel functions cannot all be zero-free on Ω_0 . The analogous statement holds for the metric ρ_2 if one restricts attention to sets of uniformly bounded Euclidean diameter.

Accordingly, to prove the theorem it will suffice to construct, arbitrarily close (according to either ρ_1 or ρ_2) to a given bounded pseudoconvex open set G , a bounded strongly pseudoconvex open set Ω with C^∞ regular boundary whose Bergman kernel function does have zeroes on some connected component; if G is connected, then Ω should be connected too.

It is standard that the pseudoconvex open set G can be exhausted from inside by strongly pseudoconvex open sets with C^∞ regular boundaries: namely, by sublevel sets of a smooth, strictly plurisubharmonic exhaustion function. It is evident that these interior approximating sets converge to G in both of the metrics ρ_1 and ρ_2 . Consequently, there is no loss of generality in supposing from the start that G is a bounded strongly pseudoconvex open set with C^∞ regular boundary.

Place close to G a strongly pseudoconvex domain D with C^∞ regular boundary and small diameter, the Bergman kernel function of D having zeroes. (In \mathbf{C}^1 , the domain D could be an annulus; in higher dimensions, D could be a small homothetic copy of the counterexample domain that I constructed in [1].) Then $G \cup D$ will be a disconnected strongly pseudoconvex open set that is close to G in both of the metrics ρ_1 and ρ_2 . This open set $G \cup D$ will serve as the required Ω to prove parts (1) and (3) of the theorem.

To prove parts (2) and (4) of the theorem, I need to produce a connected Ω when G is connected. To do this, join G to D with a closed line segment L , and use the barbell lemma to construct a sequence of bounded, connected, strongly pseudoconvex open sets Ω_k with C^∞ regular boundaries, the Ω_k being contained in $G \cup D \cup V_k$, where the V_k are shrinking neighborhoods of the line segment L . The Ω_k converge to $G \cup D$ in the metric ρ_2 , so the stability lemma and Hurwitz's theorem imply that the Bergman kernel function of Ω_k has zeroes (on D) when k is sufficiently large. Since the Euclidean distance of $D \cup V_k$ from G is small, Ω_k is close to G in the metric ρ_1 as well as in the metric ρ_2 . Thus one of the Ω_k serves as the required Ω . \square

Proof of the stability lemma. The main point is to prove an L^2 approximation theorem for holomorphic functions. I claim that if f is a square-integrable holomorphic function on U , and if a positive ϵ is prescribed, then for all sufficiently large j there exists a square-integrable holomorphic function g_j on Ω_j such that $\|f - g_j\|_{L^2(\Omega_j \cap U)} < \epsilon$ and $\|g_j\|_{L^2(\Omega_j \setminus U)} < \epsilon$.

I first need to show that the holomorphic functions in the Sobolev space $W^1(U)$ of square-integrable functions with square-integrable first derivatives are dense in the space of square-integrable holomorphic functions on U . This is a consequence of Kohn's global regularity theorem [9] for the $\bar{\partial}$ -Neumann problem with weights. Namely, for a suitably large positive number t , the weighted $\bar{\partial}$ -Neumann operator N_t for U is a bounded operator on the Sobolev space $W^2(U)$. Consequently, the corresponding weighted Bergman projection operator P_t , which satisfies the relation $P_t = \text{Id} - \bar{\partial}_t^* N_t \bar{\partial}$, maps $W^3(U)$ into the holomorphic subspace of $W^1(U)$. Now if f is a square-integrable holomorphic function in U , take a sequence $\{v_j\}$ of C^∞ functions converging to f in $L^2(U)$, and project these functions by P_t . The functions $P_t v_j$ are holomorphic functions in $W^1(U)$ that converge to f in $L^2(U)$.

Therefore, there is no loss of generality in assuming from the start that the holomorphic function f lies in $W^1(U)$. Consequently, f is the restriction to U of a function $F \in W^1(\mathbf{C}^n)$.

It follows from the hypothesis of the lemma that there is an open neighborhood V of the closure of U such that the $2n$ -dimensional Lebesgue measure of $V \cap (\Omega_j \setminus U)$ tends to zero as $j \rightarrow \infty$. There is no harm in cutting off the function F so that its support lies inside V .

The one-form $\bar{\partial}F$ is then defined on all of \mathbf{C}^n , zero on U , zero outside V , and square-integrable. Since the measure of $V \cap (\Omega_j \setminus U)$ shrinks to zero, the $L^2(\Omega_j)$ norm

of $\bar{\partial}F$ tends to zero as $j \rightarrow \infty$. Use Hörmander's L^2 theory [7] to solve the equation $\bar{\partial}u_j = \bar{\partial}F$ on Ω_j for a square-integrable function u_j whose $L^2(\Omega_j)$ norm is bounded by a constant (depending only on the uniform bound on the diameters of the Ω_j) times the $L^2(\Omega_j)$ norm of $\bar{\partial}F$. Thus the norm of u_j on Ω_j tends to zero as $j \rightarrow \infty$. Consequently, the function $g_j := F - u_j$, which is holomorphic and square-integrable on Ω_j , has norm on $\Omega_j \cap U$ close to the norm of f when j is large. Also, the norm of g_j on $\Omega_j \setminus U$ tends to zero with the measure of $V \cap (\Omega_j \setminus U)$. This confirms the claimed approximation property.

The remainder of the proof of the stability lemma follows standard lines. However, I mention that I am dispensing with the hypothesis of monotonicity of the domains that is typically assumed [8, pp. 180–182], [15, pp. 36–39].

Fix a point z in U . The Bergman kernel function $K(\cdot, z)$ (for U , or equivalently for Ω when the free variable is in U) is the unique square-integrable holomorphic function f on U that maximizes $f(z)$ subject to the nonlinear constraint $f(z) \geq \|f\|_{L^2(U)}^2$. Let f_j denote the corresponding extremal function for the approximating domain Ω_j . By the mean-value property of holomorphic functions, $f_j(z)$ is bounded by a constant times $\|f_j\|_{L^2(\Omega_j)}$ times an inverse power of the distance from z to the boundary of Ω_j ; the extremal property of f_j then implies that $\|f_j\|_{L^2(\Omega_j)}$ too is bounded by a constant times an inverse power of the distance from z to the boundary of Ω_j . Therefore the $\|f_j\|_{L^2(\Omega_j)}$ are uniformly bounded, and so the f_j form a normal family on U . Consequently, the f_j have a subsequence that converges uniformly on compact subsets of U to a holomorphic limit function f_∞ . (Once I show that the limit f_∞ actually is f , it will follow that the original sequence $\{f_j\}$, not just a subsequence, converges to f .)

By Fatou's lemma, it follows that the limit function f_∞ satisfies $f_\infty(z) \geq \|f_\infty\|_{L^2(U)}^2$. By the approximation property proved above, there exists a square-integrable holomorphic function g_j on Ω_j such that $g_j(z) \geq \|g_j\|_{L^2(\Omega_j)}^2$, and $g_j(z) \geq (1 - \delta_j)f(z)$, where the positive numbers δ_j tend to zero as $j \rightarrow \infty$. The extremal function f_j therefore has the property that $f_j(z) \geq (1 - \delta_j)f(z)$. Consequently, $f_\infty(z) \geq f(z)$. The uniqueness of the extremal function implies that $f_\infty = f$. This proves that the Bergman kernel functions $K_j(w, z)$ for the Ω_j converge pointwise to $K(w, z)$ on $U \times U$.

Since $|K_j(w, z)|^2 \leq K_j(w, w)K_j(z, z)$ by the Cauchy-Schwarz inequality, and the right-hand side is bounded by a constant depending only on the distances of z and w from the boundary of Ω_j , the functions $K_j(\cdot, \cdot)$ form a normal family in $U \times U$. From the normality and the pointwise convergence just proved, it is immediate that the convergence is uniform on compact subsets of $U \times U$. □

Proof of the barbell lemma. In the complex plane \mathbf{C}^1 , there is nothing to prove, for every planar domain is strongly pseudoconvex. In higher dimensions, there is no loss of generality in supposing that the curve γ meets the boundaries of G_1 and G_2 transversely, since the barbell Ω is not prescribed inside the neighborhood V . By [4, Theorem 4] (a result that the authors attribute to [5]), the set $\bar{G}_1 \cup \bar{G}_2 \cup \gamma$ has a basis of Stein neighborhoods, so there exists a connected, strongly pseudoconvex domain with C^∞ regular boundary that outside V is a small perturbation of $G_1 \cup G_2$. This conclusion is already enough for the application to the proof of the main theorem.

The stronger statement that one can find a barbell that actually matches $G_1 \cup G_2$ outside a neighborhood of the curve γ was demonstrated by Shcherbina for the

case when G_1 and G_2 are balls [14, Lemma 1.2 and its Corollary]. The general case follows from this special one because any strongly pseudoconvex domain can be perturbed in an arbitrarily small neighborhood of a boundary point to obtain a new strongly pseudoconvex domain whose boundary near that point is a piece of the boundary of a ball. This can be seen from the patching lemma for strictly plurisubharmonic functions in [3, Lemma 3.2.2] by taking the totally real set there to be a single point. \square

OPEN QUESTIONS

(1) In the stability lemma, the C^∞ regularity hypothesis can be reduced to C^2 regularity by inspecting Kohn's proof [9] to see that C^{k+1} boundary regularity suffices for W^k regularity of the weighted $\bar{\partial}$ -Neumann operator; one also needs techniques as in [2] to see that the weighted Bergman projection has the same regularity as the weighted $\bar{\partial}$ -Neumann operator. Can the hypothesis in the stability lemma be reduced to C^1 boundary regularity?

(2) The conclusion of the theorem—that most pseudoconvex domains are not Lu Qi-Keng domains—changes if the topology on domains is changed. For example, any small C^∞ perturbation of the unit ball is a Lu Qi-Keng domain [6]. Does the set of bounded pseudoconvex Lu Qi-Keng domains with C^1 regular boundary have nonempty interior in the C^1 topology on pseudoconvex domains? This is the case for domains in the complex plane \mathbf{C}^1 .

(3) My proof of the stability lemma for the Bergman kernel function uses pseudoconvexity. Can the word “pseudoconvex” be removed from the statement of the main theorem?

(4) Is every bounded *convex* domain a Lu Qi-Keng domain?

REFERENCES

- [1] H. P. Boas, Counterexample to the Lu Qi-Keng conjecture, *Proc. Amer. Math. Soc.* **97** (1986), 374–375. MR **87i**:32035
- [2] H. P. Boas and E. J. Straube, Equivalence of regularity for the Bergman projection and the $\bar{\partial}$ -Neumann operator, *manuscripta math.* **67** (1990), 25–33. MR **90k**:32057
- [3] Yakov Eliashberg, Topological characterization of Stein manifolds of dimension > 2 , *International J. Math.* **1** (1990), no. 1, 29–46. MR **91k**:32012
- [4] John E. Fornæss and Bill Zame, Runge exhaustions of domains in \mathbf{C}^n , *Math. Z.* **194** (1987), 1–5. MR **88c**:32026
- [5] John Erik Fornæss and Edgar Lee Stout, Spreading polydiscs on complex manifolds, *Amer. J. Math.* **99** (1977), no. 5, 933–960. MR **57**:10009
- [6] R. E. Greene and Steven G. Krantz, Stability properties of the Bergman kernel and curvature properties of bounded domains, in *Recent Developments in Several Complex Variables*, Princeton Univ. Press, Princeton, NJ, 1981, 179–198. MR **83d**:32023
- [7] Lars Hörmander, L^2 estimates and existence theorems for the $\bar{\partial}$ -operator, *Acta Math.* **113** (1965) 89–152. MR **31**:3691
- [8] M. Jarnicki and P. Pflug, *Invariant Distances and Metrics in Complex Analysis*, de Gruyter, Berlin, New York, 1993. MR **94k**:32039
- [9] J. J. Kohn, Global regularity for $\bar{\partial}$ on weakly pseudo-convex manifolds, *Trans. Amer. Math. Soc.* **181** (1973), 273–292. MR **49**:9442
- [10] Steven G. Krantz, *Function Theory of Several Complex Variables*, second edition, Wadsworth & Brooks/Cole, Pacific Grove, CA, 1992. MR **93c**:32001
- [11] Lu Qi-Keng, On Kaehler manifolds with constant curvature, *Chinese Math.* **8** (1966), 283–298; English translation of *Acta Math. Sinica* **16** (1966), 269–281. MR **34**:6806
- [12] I. Ramadanov, Sur une propriété de la fonction de Bergman, *C. R. Acad. Bulgare Sci.* **20** (1967), 759–762. MR **37**:1632

- [13] I.-P. Ramadanov, Some applications of the Bergman kernel to geometrical theory of functions, *Complex Analysis*, Banach Center Publications, Vol. 11, PWN—Polish Scientific Publishers, Warsaw, 1983, 275–286. MR **85h**:32040
- [14] N. V. Shcherbina, On fibering into analytic curves of the common boundary of two domains of holomorphy, *Math. USSR Izvestiya* **21** (1983), no. 2, 399–413; English translation of *Izv. Akad. Nauk SSSR Ser. Mat.* **46** (1982), no. 5, 1106–1123. MR **84b**:32018
- [15] M. Skwarczyński, Biholomorphic invariants related to the Bergman function, *Dissertationes Math.* **173** (1980). MR **82e**:32038
- [16] N. Suita and A. Yamada, On the Lu Qi-Keng conjecture, *Proc. Amer. Math. Soc.* **59** (1976), 222–224. MR **54**:13142

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TEXAS 77843–3368

E-mail address: boas@math.tamu.edu