

$C(K, A)$  AND  $C(K, H^\infty)$  HAVE  
THE DUNFORD-PETTIS PROPERTY

MANUEL D. CONTRERAS AND SANTIAGO DÍAZ

(Communicated by Theodore W. Gamelin)

ABSTRACT. Denote by  $X$  either the disc algebra  $A$ , or the space  $H^\infty$  of bounded analytic functions on the disc, or any of their even duals. Then  $C(K, X)$  has the Dunford-Pettis property.

An open question in the theory of Banach spaces is when the space  $C(K, X)$  of all continuous functions defined over a Hausdorff compact space  $K$  with values in a Banach space  $X$  has the Dunford-Pettis property. The natural conjecture, that this occurs *if and only if*  $X$  does, was disproved by Talagrand [13]. On the other hand, Bourgain [1] showed that this is the case when  $X$  is an abstract  $L^1(\mu)$ -space and this was extended to  $\mathcal{L}^1$ -spaces by Emmanuele [8]. Moreover, it is easy to see that the conjecture is true for  $\mathcal{L}^\infty$ -spaces. It is known that some classical Banach spaces of analytic functions have the Dunford-Pettis property (see Bourgain [2], Chaumat [3]). This led E. Saab and P. Saab to pose the following problem [12, Question 14]: if  $A$  is the disc algebra, does  $C(K, A)$  have the Dunford-Pettis property? In this paper, we give a positive answer to this question and also show that  $C(K, H^\infty)$  has this property, where  $H^\infty$  is the space of bounded analytic functions on the disc. Note that  $A$  and  $H^\infty$  are not  $\mathcal{L}^p$ -spaces for any  $1 \leq p \leq \infty$ , since they do not have even local unconditional structure [11, Theorem 4.2].

A Banach space  $X$  is said to have the Dunford-Pettis property if every weakly compact operator from  $X$  into an arbitrary Banach space is completely continuous, or equivalently, if given sequences  $(x_n)$  in  $X$  and  $(x_n^*)$  in  $X^*$ , both weakly convergent to zero, then  $\langle x_n^*, x_n \rangle$  tends to zero. A detailed exposition of this property can be found in [5]. For the rest of the notation and terminology used and not explained here, we refer the reader to the monographs of Diestel and Uhl [6], Lindenstrauss and Tzafriri [10], and Wojtaszczyk [14]. We begin by proving a technical result, which is of independent interest.

**Theorem 1.** *If  $\ell^\infty(X)$  has the Dunford-Pettis property, then  $C(K, X)$  has the same property for every compact Hausdorff space  $K$ .*

*Proof.* Let  $\Sigma$  be the Borel  $\sigma$ -field of  $K$  and  $\mathcal{B}(\Sigma, X)$  be the Banach space of all totally  $\Sigma$ -measurable functions with values in  $X$  [7, Chapter II. §9]. It is well-known [6, p. 182] that  $C(K, X)^*$  is isometrically isomorphic to the Banach space

---

Received by the editors January 6, 1995 and, in revised form, May 9, 1995.

1991 *Mathematics Subject Classification.* Primary 46E15, 46E40; Secondary 46B03, 46B25.

*Key words and phrases.* Dunford-Pettis property, disc algebra, bounded analytic functions .

This research has been partially supported by La Consejería de Educación y Ciencia de la Junta de Andalucía.

$M(K, X^*)$  of all weak\*-regular  $X^*$ -valued vector measures defined on  $\Sigma$  that are of bounded variation. Then, using the representation of  $\mathcal{B}(\Sigma, X)$  as the injective tensor product of  $\mathcal{B}(\Sigma)$  and  $X$ , and [8, Lemma 1], we have that the canonical injection  $J_1$  of  $C(K, X)$  into its bidual can be extended to an isometric operator, which we still denote  $J_1$ , defined over  $\mathcal{B}(\Sigma, X)$  and with values in  $C(K, X)^{**}$ . Namely,

$$\langle \mu, J_1(f) \rangle = \int_K f d\mu, \quad \mu \in M(K, X^*), \quad f \in \mathcal{B}(\Sigma, X).$$

Suppose that  $(f_n)$  and  $(\mu_n)$  are sequences in  $C(K, X)$  and  $C(K, X)^*$  respectively, both weakly convergent to zero. We have to prove that  $\langle \mu_n, f_n \rangle$  tends to zero. For each  $n \in \mathbb{N}$ ,  $f_n(K)$  is a norm-compact subset of  $X$ . Therefore, there is a simple  $\Sigma$ -measurable  $X$ -valued function  $g_n(\cdot) = \sum_{i=1}^{p_n} x_n(i) \chi_{A_n(i)}(\cdot)$  such that  $\|f_n - g_n\| \leq 1/n$ , for the  $\mathcal{B}(\Sigma, X)$ -norm. Note that

$$|\langle \mu_n, f_n \rangle| = \left| \int_K f_n d\mu_n \right| \leq \frac{1}{n} \|\mu_n\| + \left| \int_K g_n d\mu_n \right|.$$

Hence, it is enough to prove that  $\langle \mu_n, J_1(g_n) \rangle$  tends to zero.

For every  $n \in \mathbb{N}$ , consider the following family of pairwise disjoint  $\Sigma$ -measurable subsets of  $K$ :

$$F_n = \{A_1(m_1) \cap A_2(m_2) \cap \dots \cap A_n(m_n) : m_i = 1, \dots, p_i, \quad i = 1, \dots, n\}.$$

Since  $F_n$  has a finite number of elements, by removing those with empty intersections, we can write  $F_n = \{B_1^n, B_2^n, \dots, B_{q_n}^n\}$  for some  $q_n \in \mathbb{N}$ . Now, define the closed subspace of  $\mathcal{B}(\Sigma, X)$

$$Y_n = \left\{ \sum_{i=1}^{q_n} x_i \chi_{B_i^n} : x_i \in X \right\},$$

which is clearly isometrically isomorphic to  $\ell_{q_n}^\infty(X)$ . Since  $(Y_n)$  is an increasing sequence of subspaces, we can consider the space  $Y$  defined as the closure in  $\mathcal{B}(\Sigma, X)$  of their union. We observe that  $g_n$  belongs to  $Y$ , for all  $n \in \mathbb{N}$ . On the other hand, let  $J_2$  be the canonical injection of  $C(K, X)^*$  into  $C(K, X)^{***}$  and denote by  $J_2^Y(\mu_n)$  the restriction of  $J_2(\mu_n)$  to  $J_1(Y)$ . It is easy to check that  $(J_1(g_n))$  and  $(J_2^Y(\mu_n))$  are weakly null sequences in  $J_1(Y)$  and  $J_1(Y)^*$ , respectively. Moreover,

$$\langle J_2^Y(\mu_n), J_1(g_n) \rangle = \langle \mu_n, J_1(g_n) \rangle = \int_K g_n d\mu_n.$$

Therefore, the theorem will we proved if we show that  $Y$  has the Dunford-Pettis property. According to a result due to Bourgain [1, Proposition 2], we only have to prove that  $(\oplus_n Y_n)_\infty$  has the Dunford-Pettis property. But, this follows on combining the hypothesis that  $\ell^\infty(X)$  has the Dunford-Pettis property with the following topological isomorphisms:

$$(\oplus_n Y_n)_\infty \cong (\oplus_n \ell_{q_n}^\infty(X))_\infty \cong (\oplus_n X)_\infty = \ell^\infty(X). \quad \square$$

In general, the converse of the above theorem is not true. The example goes as follows: Let  $M_n$  be the algebra of  $n \times n$  complex matrices. Then, by [4, p. 61], we know that  $Z = (\oplus_n M_n)_0$  is a  $C^*$ -algebra which has the Dunford-Pettis property; thus  $C(K, Z)$  has the Dunford-Pettis property for every compact Hausdorff space  $K$  [12, Theorem 48]. On the other hand,  $\ell^\infty(Z)$  contains a complemented copy of  $(\oplus_n M_n)_\infty$ , thus a complemented copy of  $(\oplus_n \ell_n^2)_\infty$ . Finally, note that this last space has a complemented copy of the reflexive space  $\ell^2$  [5, p. 22].

Our next theorem is the main result of this paper. As we pointed out, it answers Question 14 of [12].

**Theorem 2.**  $C(K, A)$  and  $C(K, H^\infty)$  have the Dunford-Pettis property for every compact Hausdorff space  $K$ .

*Proof.* In both cases, we shall apply Theorem 1. Since  $H^\infty$  has the Dunford-Pettis property [2, Corollary 5.4], the case of  $H^\infty$  follows directly from the topological isomorphism  $\ell^\infty(H^\infty) \cong H^\infty$  [14, Part III.E.13].

For the case of the disc algebra  $A$ , recall that  $A$  can be described as the smallest closed subspace of  $C(\mathbb{T})$  containing all complex polynomials. Therefore, it is easy to see that

$$\ell^\infty(A) = \|\cdot\|_\infty\text{-closure} \left\{ \ell^\infty \left( \bigcup_{k=1}^\infty A_k \right) \right\},$$

where  $A_k = \text{span}\{1, z, \dots, z^{2k}\}$ .

Suppose that  $(x_n)$  and  $(x_n^*)$  are sequences in  $\ell^\infty(A)$  and  $\ell^\infty(A)^*$ , respectively, that converge weakly to zero. Using the above expression for  $\ell^\infty(A)$ , we can find  $y_n \in \ell^\infty(\bigcup_{k=1}^\infty A_k)$  such that  $\|x_n - y_n\| \leq \frac{1}{n}$ , for all  $n \in \mathbb{N}$ . Moreover, for each  $m \in \mathbb{N}$ , we can find  $k = k(n, m) \in \mathbb{N}$  such that  $y_n(m) \in A_{k(n, m)}$  and  $k(n + 1, m) > k(n, m)$ , for  $n, m \in \mathbb{N}$ . Of course, we only have to show that  $\langle x_n^*, y_n \rangle$  tends to zero.

Consider the subspace  $Z_n = (\oplus_m A_{k(n, m)})_\infty \subset \ell^\infty(A)$ . For every  $n \in \mathbb{N}$ ,  $Z_n$  is contained in  $Z_{n+1}$ , so we can define  $Z$  as the subspace of  $\ell^\infty(A)$  given by the closure of  $\bigcup_{n=1}^\infty Z_n$ . We observe that  $(y_n)$  is a weakly null sequence in  $Z$ , and the restriction of  $x_n^*$  to  $Z$  is also a weakly null sequence in  $Z^*$ . Hence, to finish the proof, it is enough to show that  $Z$  has the Dunford-Pettis property or, according again to [1, proposition 2], that  $(\oplus_n Z_n)_\infty$  has the Dunford-Pettis property. It is clear that  $(\oplus_n Z_n)_\infty = (\oplus_n (\oplus_m A_{k(n, m)})_\infty)_\infty$  is isomorphic to a complemented subspace of  $(\oplus_n (\oplus_m A_m)_\infty)_\infty$ . Finally, by [14, Part III.E.18], this last space is isomorphic to  $\ell^\infty(H^\infty)$ , and we reduce to the former situation.  $\square$

In our next result, we extend Theorem 2 to other Banach spaces, namely, the even duals of  $A$  and  $H^\infty$ . To fix the notation, we denote by  $X^{(n)}$  the  $n^{\text{th}}$  dual of the Banach space  $X$  and  $X^{(0)} = X$ .

**Theorem 3.**  $C(K, A^{(2n)})$  and  $C(K, (H^\infty)^{(2n)})$  have the Dunford-Pettis property for every compact Hausdorff space  $K$  and for all  $n \in \mathbb{N}$ .

*Proof.* In the same way as in the scalar case, it can be shown that  $\ell^\infty(X)^*$  is the topological direct sum of (a subspace isomorphic to)  $\ell^1(X^*)$  and the space of those functionals vanishing over  $c_0(X)$ . Then,  $\ell^\infty(X)^{**}$  contains a complemented subspace isomorphic to  $\ell^\infty(X^{**})$  and, by induction, we obtain that  $\ell^\infty(X)^{(2n)}$  contains a complemented subspace isomorphic to  $\ell^\infty(X^{(2n)})$ , for all  $n \in \mathbb{N}$ .

Since any of the duals of  $H^\infty$  have the Dunford-Pettis property [2, Corollary 5.4], we have that  $\ell^\infty((H^\infty)^{(2n)})$  has the Dunford-Pettis property, and the case of  $H^\infty$  follows from Theorem 1.

On the other hand, since  $A^{**}$  is isomorphic to the  $\ell^\infty$ -sum of  $H^\infty$  and a certain  $\mathcal{L}^\infty$ -space  $L$  [11, Chapter 1], we have the following topological isomorphisms:

$$\ell^\infty(A^{(2n)}) \cong \ell^\infty((H^\infty)^{(2n-2)}) \oplus_\infty \ell^\infty(L^{(2n-2)}), \quad n \in \mathbb{N}.$$

Since  $L^{(2n-2)}$  are again  $\mathcal{L}^\infty$ -spaces, in order to prove that  $\ell^\infty(A^{(2n)})$  has the Dunford-Pettis property, we only need to show that  $\ell^\infty(X)$  has the Dunford-Pettis property whenever  $X$  is an arbitrary  $\mathcal{L}^\infty$ -space. Now we recall the following characterization: a Banach space  $X$  is an  $\mathcal{L}^\infty$ -space if and only if there is a real number  $\lambda$  such that for all Banach spaces  $Z \supset Y$ , every compact operator  $T : Y \rightarrow X$  has an extension  $\tilde{T} : Z \rightarrow X$  with  $\|\tilde{T}\| \leq \lambda\|T\|$  [9, Theorem 4.1 and its proof]. Hence, dealing with the projections from  $\ell^\infty(X)$  onto its different coordinates, we see that  $\ell^\infty(X)$  is an  $\mathcal{L}^\infty$ -space and, therefore, its bidual is isomorphic to a complemented subspace of some  $C(K)$ .  $\square$

**Addendum.** After this paper was submitted, N. Randrianantoanina has proved, with completely different techniques, that  $C(K, A)^*$  has the Dunford-Pettis property in a paper posted at the Banach bulletin board entitled “Some results on the Dunford-Pettis property”. The authors want to thank the referee for pointing out to them Randrianantoanina’s result.

#### REFERENCES

1. J. Bourgain, *On the Dunford-Pettis property*, Proc. Amer. Math. Soc. **81** (1981), 265–272. MR **83g**:46038
2. ———, *New Banach space properties of the disc algebra and  $H^\infty$* , Acta Math. **152** (1984), 1–48. MR **85j**:46091
3. J. Chaumat, *Une généralisation d’un théorème de Dunford-Pettis*, Université de Paris XI, Orsay, U.E.R. Mathématique no. 85, 1974.
4. C.-H. Chu and B. Iochum, *The Dunford-Pettis property in  $C^*$ -algebras*, Studia Math. **97** (1990), 59–64. MR **92b**:46091
5. J. Diestel, *A survey of results related to the Dunford-Pettis property*, Contemporary Math., vol. 2, Proc. of the Conf. on Integration, Topology and Geometry in Linear Spaces, Amer. Math. Soc., Providence, RI, 1980, pp. 15–60. MR **82i**:46023
6. J. Diestel and J. Uhl, *Vector Measures*, Math. Surveys, vol. 15, Amer. Math. Soc., Providence, RI, 1977. MR **56**:12216
7. N. Dinculeanu, *Vector Measures*, Pergamon Press, New York, 1967. MR **34**:6011
8. G. Emmanuele, *Remarks on weak compactness of operators defined on certain injective tensor products*, Proc. Amer. Math. Soc. **116** (1992), 473–476. MR **92m**:46109; MR **93f**:47022
9. J. Lindenstrauss and H. P. Rosenthal, *The  $\mathcal{L}_p$ -spaces*, Israel J. Math. **7** (1969), 325–349. MR **42**:5012
10. J. Lindenstrauss and L. Tzafriri, *Classical Banach Spaces*, Lecture Notes in Mathematics, vol. 338, Springer-Verlag, Berlin-Heidelberg-New York, 1973. MR **54**:3344
11. A. Pełczyński, *Banach spaces of analytic functions and absolutely summing operators*, CBMS, Regional Conference Series, no. 30, Amer. Math. Soc., Providence, RI, 1977. MR **58**:23526
12. E. Saab and P. Saab, *On stability problems of some properties in Banach spaces*, K. Jarosz (Ed.), Lecture Notes in Pure and Appl. Math., vol. 136, Marcel Dekker, 1992, pp. 367–394. MR **92m**:46021
13. M. Talagrand, *La propriété de Dunford-Pettis dans  $C(K, E)$  et  $L_1(E)$* , Israel J. Math. **44** (1983), 317–321. MR **84j**:46065
14. P. Wojtaszczyk, *Banach Spaces for Analysts*, Cambridge Studies in Advanced Mathematics, vol. 25, Cambridge University Press, Cambridge, 1991. MR **93d**:46001

E. S. INGENIEROS INDUSTRIALES, AVDA. REINA MERCEDES S/N, 41012-SEVILLA, SPAIN  
E-mail address: [contreras@cica.es](mailto:contreras@cica.es)

E-mail address: [madrigal@cica.es](mailto:madrigal@cica.es)