

CLASSIFYING SPACES AND HOMOTOPY SETS OF AXES OF PAIRINGS

KENSHI ISHIGURO

(Communicated by Thomas Goodwillie)

Dedicated to Professor Teiichi Kobayashi on his 60th birthday

ABSTRACT. We consider the maps between classifying spaces of the form $BK \times BL \rightarrow BG$. The main theorem shows that if the restriction map on BL is a weak epimorphism, then the restriction on BK should factor through the classifying spaces of the center of the compact Lie group G . An application implies that BG is an H-space (Hopf space) if and only if G is abelian.

For a map $f : Y \rightarrow Z$, the set of the homotopy classes of axes, denoted by $f^\perp(X, Z)$, consists of all homotopy classes of maps $\alpha : X \rightarrow Z$ such that there is a map (called a pairing) $\mu : X \times Y \rightarrow Z$ with restrictions (axes) $\mu|_X \simeq \alpha$ and $\mu|_Y \simeq f$, [16]. If Z is an H-space, then $f^\perp(X, Z) = [X, Z]$: for the H-multiplication $m : Z \times Z \rightarrow Z$, a pairing of f and α is given by the composite map $m \circ (\alpha \times f)$. It follows that, for example, if G, K , and L are compact Lie groups, then $f^\perp(K, G) = [K, G]$ with $f : L \rightarrow G$. In this paper we will study the classifying space version, namely $f^\perp(BK, BG)$ with $f : BL \rightarrow BG$. As group theoretical analog indicates, our results will show that few maps in $[BK, BG]$ belong to $f^\perp(BK, BG)$, in general. Other forms of axial maps (H-pairing) are studied in [6].

The main theorem deals with the case that the map $f : BL \rightarrow BG$ is a weak epimorphism studied in [8]. (We recall the definition in §2.) An obvious example of a weak epimorphism is given by a map $f = B\rho$ induced by a group epimorphism ρ . The unstable Adams operations $\{\psi^k\}$ are also weak epimorphisms.

Theorem 1. *Let L and G be connected compact Lie groups and let K be a compact (not necessarily connected) Lie group. If $f : BL \rightarrow BG$ is a weak epimorphism, the following hold:*

- (1) *If $\alpha \in f^\perp(BK, BG)$, then the map α factors through $BZ(G)$ up to homotopy, where $Z(G)$ denotes the center of G .*
- (2) *Moreover, we have $f^\perp(BK, BG) = \text{Hom}(K, Z(G))$.*

Taking $L = G$ and $f = id$ (the identity map), our problem asks about possible BK -actions on BG . In fact, this work was motivated by the following result of G. Dula and D. Gottlieb.

Received by the editors April 10, 1995 and, in revised form, June 15, 1995.

1991 *Mathematics Subject Classification.* Primary 55R35; Secondary 55P15, 55P60.

Key words and phrases. Classifying space, mapping spaces, p -completion, Lie groups, weak epimorphism, pairing.

Theorem 2 ([2]). *Let $\alpha : X \rightarrow Z$ be a map, and X an H -space. Then the following are equivalent.*

- (a) *There exists a space Y and a homotopy equivalence $X \times Y \rightarrow Z$ such that the orbit map $X \rightarrow X \times Y \rightarrow Z$ is homotopy equivalent to α .*
- (b) *The map α is the orbit of an action of X on Z and $\alpha^\# : [Z, X] \rightarrow [X, X]$ is onto.*

For instance, let $X = BS^1$ and $Z = BU(n)$. A map $\alpha : BS^1 \rightarrow BU(n)$ is induced by a homomorphism $\rho : S^1 \rightarrow U(n)$ so that $\alpha = B\rho$. Since $BU(n)$ is indecomposable, [7], Theorem 2 implies that the BS^1 -action on $BU(n)$ under the map α does not exist if α is induced by the inclusion $S^1 = U(1) \hookrightarrow U(n)$. Theorem 1 says that the BS^1 -action on $BU(n)$ under the map α exists if and only if $\rho(S^1)$ is central in $U(n)$. Generally, a consequence of Theorem 1 shows that BK -actions on BG under α exist if and only if the map $\alpha : BK \rightarrow BG$ is induced by a homomorphism $\rho : K \rightarrow G$ such that $\rho(K)$ is central in G . In particular, if K is connected and G is semi-simple, there are no non-trivial BK -actions on BG .

Furthermore, if we take $K = L = G$ and $f = \alpha = id$, the problem now asks whether BG is an H -space. Corollary 2.4 implies that BG is an H -space if and only if G is abelian. (G need not be connected.) Related results were obtained in [12] and [1].

I would like to thank Giora Dula for bringing this problem to my attention. I would also like to thank my colleague, Nobuyuki Oda for his help.

1. MAPPING SPACES AND CENTRALIZERS OF p -TORAL GROUPS

We give a necessary and sufficient condition that a map $\alpha : X \rightarrow Z$ be contained in $f^\perp(X, Z)$ in terms of mapping spaces. Then a special case of classifying space version is discussed.

Proposition 1.1. *Suppose X, Y and Z are pointed connected spaces. For a map $f : Y \rightarrow Z$, a map $\alpha : X \rightarrow Z$ is contained in $f^\perp(X, Z)$ if and only if the map f factors through $map(X, Z)_\alpha$, the connected component of the mapping space containing α , under the evaluation map $map(X, Z)_\alpha \xrightarrow{ev} Z$.*

Remark. It is easy to see that $\alpha \in f^\perp(X, Z)$ if and only if $f \in \alpha^\perp(Y, Z)$. Consequently α must factor through $map(Y, Z)_f$.

Proof of Proposition 1.1. If $\alpha \in f^\perp(X, Z)$, there is a pairing $\mu : X \times Y \rightarrow Z$ with $\mu|_X \simeq \alpha$ and $\mu|_Y \simeq f$. Let $\bar{\mu} : Y \rightarrow map(X, Z)_\alpha$ be the adjoint map of μ . Then we see that the map f is expressed as the composite of the adjoint map and the evaluation map :

$$f : Y \xrightarrow{\bar{\mu}} map(X, Z)_\alpha \xrightarrow{ev} Z$$

since $ev \circ \bar{\mu}(y) = \bar{\mu}(y)(*) = \mu(*, y) = f(y)$.

Conversely, suppose f factors through $map(X, Z)_\alpha$ so that $f \simeq ev \circ \bar{\mu}$ for some map $\bar{\mu} : Y \rightarrow map(X, Z)_\alpha$. Let $\epsilon : X \times map(X, Z)_\alpha \rightarrow Z$ be the canonical map with $\epsilon(x, \beta) = \beta(x)$ for $x \in X$ and $\beta \in map(X, Z)_\alpha$. Then a pairing is constructed as the composite

$$\mu : X \times Y \xrightarrow{1_X \times \bar{\mu}} X \times map(X, Z)_\alpha \xrightarrow{\epsilon} Z,$$

where 1_X is the identity map of X , since $\mu(x, y) = \epsilon \circ (1_X \times \bar{\mu})(x, y) = \epsilon(x, \bar{\mu}(y)) = \bar{\mu}(y)(x)$ so that $\mu(x, *) = \bar{\mu}(*)(x) = \alpha(x)$ and $\mu(*, y) = \epsilon(*, \bar{\mu}(y)) = ev \circ \bar{\mu}(y) \simeq f(y)$. \square

We next consider the “BG”-version. It is worth to recall some property of homomorphisms. Suppose $\rho : L \rightarrow G$ and $\alpha : K \rightarrow G$ are homomorphisms. If there is a pairing homomorphism $\mu : K \times L \rightarrow G$ with $\mu|_K = \alpha$ and $\mu|_L = \rho$, then the image $\rho(L)$ must be contained in the centralizer of α in G , denoted by $C_G(\alpha)$. The following is a “BG”-analog at a prime p . If a map $\alpha : BK \rightarrow BG$ is induced by a homomorphism, let $C_G(\alpha)$ denote the centralizer of the homomorphism. For a p -toral group K (a group extension of a torus by a finite p -group), it is known, [5] and [14], that any map $\alpha : BK \rightarrow BG$ (at p) has the form $\alpha = B\eta$ ($\alpha = (B\eta)_p^\wedge$) for some homomorphism η . Let BG_p^\wedge denote the p -completion of BG . Since $map(BK, BG)_\alpha$ is p -equivalent to $BC_G(\alpha)$, the following is immediate from Proposition 1.1.

Corollary 1.2. *Suppose K is p -toral. Then $\alpha \in f^\perp(BK_p^\wedge, BG_p^\wedge)$ if and only if the map f factors through $BC_G(\alpha)_p^\wedge$ up to homotopy, under the map $BC_G(\alpha)_p^\wedge \rightarrow BG_p^\wedge$ induced by the inclusion.*

As the above indicates, if the mapping space is computable, then the set of the homotopy classes of axes $f^\perp(X, Z)$ would be determined. It is, however, hard to compute $map(X, Z)_\alpha$ or $map(Y, Z)_f$ in general. Thus our work is to consider the problem, with or without calculation of mapping spaces.

2. WEAK EPIMORPHISMS AND THE PROOF OF THEOREM 1

We will begin with some lemmas to prove Theorem 1.

Lemma 2.1. *Let R and S be rings of polynomials in n indeterminates over a field. If there is an epimorphism $\varphi : R \rightarrow S$, then φ is an isomorphism.*

Proof. Let $\mathfrak{p} = Ker \varphi$ so that $R/\mathfrak{p} \cong S$. We need to show that $\mathfrak{p} = 0$. The ideal \mathfrak{p} is prime, since R/\mathfrak{p} is an integral domain. Consequently, if $dim(R)$ denotes the Krull dimension of R , then

$$height(\mathfrak{p}) + dim(R/\mathfrak{p}) = dim(R).$$

It follows that

$$\begin{aligned} dim(S) &= dim(R/\mathfrak{p}) \\ &= dim(R) - height(\mathfrak{p}). \end{aligned}$$

Since $dim(S) = dim(R) = n$, we see that $height(\mathfrak{p}) = 0$. In a domain, this means $\mathfrak{p} = 0$. \square

Recall that, for any compact connected Lie group G , there is a covering $\gamma_G \rightarrow \tilde{G} \rightarrow G$ such that $\tilde{G} = G_s \times T$ is a product of a simply-connected Lie group and a torus, and that γ_G is a finite central subgroup of \tilde{G} . Such a covering is called a *universal finite covering*.

Lemma 2.2. *Suppose G is a connected compact Lie group and H is a connected closed subgroup with inclusion $\iota : H \rightarrow G$. If the p -completed map $(B\iota)_p^\wedge : (BH)_p^\wedge \rightarrow (BG)_p^\wedge$ is rationally equivalent, then $H = G$.*

Proof. Since H and G are connected, we can find universal finite coverings \tilde{H} and \tilde{G} so that we have the commutative diagram

$$\begin{CD} \tilde{H} @>\tilde{\iota}>> \tilde{G} \\ @VVV @VVV \\ H @>\iota>> G \end{CD}$$

Note that \tilde{H} and \tilde{G} are products of simply-connected simple Lie groups and a torus, $\tilde{H} = \prod_i \tilde{H}_i$ and $\tilde{G} = \prod_j \tilde{G}_j$ respectively. Since the map $(B\iota)_p^\wedge : (BH)_p^\wedge \rightarrow (BG)_p^\wedge$ is rationally equivalent, for each \tilde{G}_j , by [8], we can find \tilde{H}_i such that the restricted map $(B\tilde{H}_i)_p^\wedge \rightarrow (B\tilde{G}_j)_p^\wedge$ is a rational equivalence, and the map $(B\tilde{\iota})_p^\wedge$ is diagonal. Using the classification of compact simply-connected simple Lie groups and their maximal rank subgroups, one can show that $\tilde{\iota}(\tilde{H}_i) = \tilde{G}_j$. Hence $\tilde{\iota} : \tilde{H} \rightarrow \tilde{G}$ is onto. Consequently the inclusion $\iota : H \rightarrow G$ is an epimorphism, and therefore $H = G$. \square

Lemma 2.3. *Let G be a compact Lie group with center $Z(G)$. Then the map induced by the inclusion $i : Z(G) \hookrightarrow G$*

$$(Bi)_\# : [BK, BZ(G)] \rightarrow [BK, BG]$$

is one-to-one for any compact Lie group K .

Proof. Notice that $[BK, BZ(G)] = Hom(K, Z(G))$, since $Z(G)$ is abelian, [11]. For $\rho_1, \rho_2 \in Hom(K, Z(G))$, suppose $(Bi)_\#(B\rho_1) = (Bi)_\#(B\rho_2)$. Thus, for each p -toral subgroup K_q of K , we have $\rho_1(x) = \rho_2(x)$ for any $x \in K_q$, since the images of ρ_1 and ρ_2 are contained in the center $Z(G)$. Define $\rho \in Hom(K, Z(G))$ by $\rho(k) = \rho_1(k)\rho_2(k)^{-1}$ for any $k \in K$ so that $\rho|_{K_q} = 0$ for any p -toral subgroup K_q . Since $[BK_p^\wedge, BG_p^\wedge] = [\text{holim} \rightarrow (BK_q)_p^\wedge, BG_p^\wedge] = \text{holim} \leftarrow [(BK_q)_p^\wedge, BG_p^\wedge]$, it follows that $\rho = 0$. Consequently $\rho_1 = \rho_2$, and therefore $B\rho_1 = B\rho_2$. \square

Remark. For $\iota : H \hookrightarrow G$ the map $[BK, BH] \rightarrow [BK, BG]$ need not be one-to-one. A counterexample is given by $[B\mathbb{Z}/2, BT^2] \not\hookrightarrow [B\mathbb{Z}/2, BU(2)]$ where T^2 is a maximal torus of $U(2)$.

We consider $f^\perp(BK, BG)$ when $f : BL \rightarrow BG$ is a weak epimorphism, defined as follows. Suppose L and G are connected. A map $BL \rightarrow BG$ or $BL_p^\wedge \rightarrow BG_p^\wedge$ is called a *weak epimorphism*, [8], if there exists a fibration $Z \rightarrow BL \rightarrow BG$ or $Z \rightarrow BL_p^\wedge \rightarrow BG_p^\wedge$ such that $H^*(\Omega Z; \mathbb{Q})$ is a finite dimensional \mathbb{Q} -module or that $H^*(\Omega Z; \mathbb{Z}_p^\wedge) \otimes \mathbb{Q}$ is a finite dimensional \mathbb{Q}_p^\wedge -module, respectively. Some examples were given in the introduction.

Proof of Theorem 1. First we show Part (1). Fix a prime p . Suppose K_q is a p -toral subgroup of K and $\alpha_q = \alpha_p^\wedge|_{(BK_q)_p^\wedge}$. Corollary 1.2 shows that the p -completed map $f_p^\wedge : BL_p^\wedge \rightarrow BG_p^\wedge$ factors through $BC_G(\alpha_q)_p^\wedge$ up to homotopy. Since K_q is p -toral, $\pi_0(C_G(\alpha_q))$ is a finite p -group. Since BL_p^\wedge is 1-connected, the map f_p^\wedge factors through the p -completed classifying space of the connected component of the centralizer $C_G(\alpha_q)$. Since f is a weak epimorphism, the map f_p^\wedge is also a weak epimorphism. Consequently there is a fibration $Z \rightarrow BL_p^\wedge \rightarrow BC_G(\alpha_q)_p^\wedge$ satisfying

certain finiteness conditions mentioned above. Let H be the connected component of $C_G(\alpha_q)$ with inclusion $\iota : H \rightarrow G$. Then we have the commutative diagram

$$\begin{CD} Y @>>> BL_p^\wedge @>\bar{f}_p^\wedge>> BH_p^\wedge \\ @VVV @| @VV(B\iota)_p^\wedge V \\ Z @>>> BL_p^\wedge @>f_p^\wedge>> BG_p^\wedge \end{CD}$$

Here Y is a homotopy fibre. Since f_p^\wedge is a weak epimorphism, we see that $H^*(BL_p^\wedge; \mathbb{Q}_p^\wedge) \cong H^*(BG_p^\wedge; \mathbb{Q}_p^\wedge) \otimes H^*(Z; \mathbb{Q}_p^\wedge)$. Thus, for the induced homomorphism $(f_p^\wedge)^* : H^*(BG_p^\wedge; \mathbb{Q}_p^\wedge) \rightarrow H^*(BL_p^\wedge; \mathbb{Q}_p^\wedge)$ there is a homomorphism (left inverse) $r : H^*(BL_p^\wedge; \mathbb{Q}_p^\wedge) \rightarrow H^*(BG_p^\wedge; \mathbb{Q}_p^\wedge)$ with $r \circ (f_p^\wedge)^* = 1$. Consequently $r \circ (\bar{f}_p^\wedge)^* \circ ((B\iota)_p^\wedge)^* = 1$. Let φ be the endomorphism $((B\iota)_p^\wedge)^* \circ r \circ (\bar{f}_p^\wedge)^*$ of the polynomial ring $H^*(BH_p^\wedge; \mathbb{Q}_p^\wedge)$. Since $r \circ (\bar{f}_p^\wedge)^*$ is onto and $((B\iota)_p^\wedge)^*$ is one-to-one, the Krull dimension of the image of φ is equal to $rank(G)$. Hence the endomorphism φ is an isomorphism and therefore $((B\iota)_p^\wedge)^*$ is onto. Lemma 2.1 implies that BH_p^\wedge is rationally equivalent to BG_p^\wedge . From Lemma 2.2, we see that $C_G(\alpha_q) = G$. This implies that the map α_q is induced by a homomorphism into the center $Z(G)$. Hence $\alpha|_{(BK_q)_p^\wedge}$ factors through $BZ(G)_p^\wedge$ for any p -toral subgroup K_q .

Since $BK_p^\wedge \simeq \varinjlim (BK_q)_p^\wedge$, [9], the map $\alpha_p^\wedge : BK_p^\wedge \rightarrow BG_p^\wedge$ factors through $BZ(G)_p^\wedge$ for any p . Let K_0 denote the connected component of K containing the identity so that there is an exact sequence $K_0 \rightarrow K \xrightarrow{\pi} \pi_0 K$. Consider the canonical fibration $BZ(G) \rightarrow BG \xrightarrow{Bq} B(G/Z(G))$. Since $(Bq \cdot \alpha|_{BK_0})_p^\wedge$ is null homotopic, the map $Bq \cdot \alpha|_{BK_0}$ is rationally null homotopic. Since K_0 is connected, then $Bq \cdot \alpha|_{BK_0} = 0$. This induces a map $\bar{\alpha} : B\pi_0 K \rightarrow B(G/Z(G))$ with $Bq \cdot \alpha = \bar{\alpha} \cdot B\pi$. Since $(Bq \cdot \alpha)_p^\wedge = 0$ for any p and the homomorphism π is onto, the restriction of $\bar{\alpha}$ on each p -Sylow subgroup of the finite group $\pi_0 K$ is null homotopic. This implies $\bar{\alpha} = 0$. Consequently α factors through $BZ(G)$.

Next we show Part (2). The canonical map $(Bi)_\# : [BK, BZ(G)] \rightarrow [BK, BG]$ factors through $f^\perp(BK, BG)$. Let Φ be the map $[BK, BZ(G)] \rightarrow f^\perp(BK, BG)$ and let Ψ be the map $f^\perp(BK, BG) \rightarrow [BK, BG]$ so that $(Bi)_\# = \Psi \circ \Phi$. Part (1) shows Φ is onto. Lemma 2.3 shows $(Bi)_\#$ is one-to-one. This implies Φ is one-to-one. Thus Φ is bijective. Since $[BK, BZ(G)] = Hom(K, Z(G))$, this completes the proof. \square

Remark. As shown in §1, there is a strong relationship between the set of the homotopy classes of axes $f^\perp(BK, BG)$ and $map(BL, BG)_f$. Theorem 1 seems to indicate that $map(BL, BG)_f$ can be homotopy equivalent to $BZ(G)$ when $f : BL \rightarrow BG$ is a weak epimorphism. One can show, however, that $map(BS^3, BS^3)_{\psi^k} \not\cong BZ(S^3)$ (without p -completion), from the following result of Dwyer–Mislin [3]: The space of pointed maps $map_*(BS^3, BS^3)_{\psi^k}$ is homotopy equivalent to Sullivan’s profinite completion $SO(3)^\wedge$. What to ask is the p -equivalence between $map(BL, BG)_f$ and $BZ(G)$. A result of Jackowski–McClure–Oliver [9] and Notbohm [15] shows that if $f : BG \rightarrow BG$ is a self-equivalence, the map $BZ(G) \rightarrow map(BG, BG)_f$ is a mod p equivalence for any prime p . A related result can be found in [4].

Corollary 2.4. *Suppose G is a compact (not necessarily connected) Lie group. If BG is an H -space, then G is abelian.*

Proof. If BG is an H-space, then $(id)^\perp(BG, BG) = [BG, BG]$. Suppose first that G is connected. Taking $\alpha = id$ in Theorem 1, we see that the identity map of BG factors through $BZ(G)$. Consequently $G = Z(G)$, and therefore G is a torus. In general, there is the exact sequence $G_0 \rightarrow G \rightarrow \pi_0 G$. If BG is an H-space, from the fibration $BG_0 \rightarrow BG \rightarrow B\pi_0 G$ it follows that both BG_0 and $B\pi_0 G$ are also H-spaces. This implies that the groups G_0 and $\pi_0 G$ are abelian, since $\pi_0 G$ is a finite group so that $B\pi_0 G = K(\pi_0 G, 1)$. Let $\pi = \pi_0 G$ and let π_p denote a p -Sylow subgroup of the finite group π . Suppose G_p is the subgroup of G over π_p :

$$\begin{array}{ccccc} G_0 & \longrightarrow & G & \longrightarrow & \pi_0 G = \pi \\ \parallel & & \uparrow & & \uparrow \\ G_0 & \longrightarrow & G_p & \longrightarrow & \pi_p \end{array}$$

Since BG is an H-space, the space BG_p is also an H-space. Notice that G_p is p -toral. Consequently the H-multiplication on BG_p is induced by a homomorphism $G_p \times G_p \rightarrow G_p$. This implies that G_p is abelian for any prime p , and therefore $G (= G_0 \times \pi_0 G)$ is abelian. \square

Remark. The p -completed version of Corollary 2.4 does not hold. For instance, the symmetric group Σ_3 is non-abelian, but the 2-completed classifying space $(B\Sigma_3)_2^\wedge = B\mathbb{Z}/2$ is an H-space.

REFERENCES

1. J. Aguadé and L. Smith, *On the mod p torus theorem of John Hubbuck*, Math. Z. **191** (1986), 325-326. MR **87e**:57044
2. G. Dula and D. Gottlieb, *Splitting off H-spaces and Conner-Raymond Splitting Theorem*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **37** (1990), 321-334. MR **91i**:55018
3. W.G. Dwyer and G. Mislin, *On the homotopy type of $map_*(BS^3, BS^3)$* , Proc. 1986 Barcelona conference, LNM **1298** (1987), 82-89. MR **89e**:55018
4. W.G. Dwyer and C. Wilkerson, *The center of a p -compact group*, Preprint.
5. W.G. Dwyer and A. Zabrodsky, *Maps between classifying spaces*, Proc. 1986 Barcelona conference, LNM **1298** (1987), 106-119. MR **89b**:55018
6. Y. Hemmi, *The projective plane of an H-pairing*, J.P.A.A. **75** (1991), 277-296. MR **92m**:55011
7. K. Ishiguro, *A p -local splitting of $BU(n)$* , Proc. of AMS **95**(2) (1985), 307-311. MR **86m**:55023
8. K. Ishiguro and D. Notbohm, *Fibrations of classifying spaces*, Transaction of AMS **343**(1) (1994), 391-415. MR **94h**:55028
9. S. Jackowski, J.E. McClure and B. Oliver, *Homotopy classification of self-maps of BG via G -actions Part I and Part II*, Ann. of Math. **135** (1992), 183-226, 227-270. MR **93e**:55019
10. S. Jackowski, J.E. McClure and B. Oliver, *Self homotopy equivalences of classifying spaces of compact connected Lie groups*, Preprint.
11. R. Lashof, P. May and G. Segal, *Equivariant bundles with abelian structural group*, Contemporary Math. **19** (1983), 167-176. MR **85b**:55023
12. J. Lin, *A cohomological proof of the torus theorem*, Math. Z. **190** (1985), 469-476. MR **87c**:55009
13. G. Mislin and J. Thévenaz, *The Z^* -Theorem for compact Lie groups*, Math. Ann. **291** (1991), 103-111. MR **92i**:22006
14. D. Notbohm, *Maps between classifying spaces*, Math. Z. **207** (1991), 153-168. MR **92b**:55017
15. D. Notbohm, *Maps between classifying spaces and applications*, J.P.A.A. **89** (1993), 273-294. MR **95c**:55019

16. N. Oda, *The homotopy set of the axes of pairings*, *Canad. J. Math.* **42** (1990), 856–868. MR **91i:55019**
17. N. Oda, *Localization of the homotopy set of the axes of pairings*, Adams memorial symposium on algebraic topology volume 1, *London Math. Soc. Lecture Notes* 175, Cambridge Univ. Press (1992), 163–177. MR **93i:55014**

DEPARTMENT OF MATHEMATICS, FUKUOKA UNIVERSITY, FUKUOKA 814-80, JAPAN
E-mail address: kenshi@ssat.fukuoka-u.ac.jp