

ON AN OPTIMALITY PROPERTY OF RAMANUJAN SUMS

GENNADY BACHMAN

(Communicated by William W. Adams)

ABSTRACT. We evaluate $\inf_{b_n} \sum_{a=1}^q \left| \sum_{\substack{n=1 \\ (n,q)=1}}^q b_n e^{2\pi i a n/q} \right|$, where the inf is taken over sequences b_n satisfying $b_n \geq 1$. In particular we show that it is attained by taking $b_n = 1$ for all n , which reduces the summation over n to a Ramanujan sum $c_q(a) = \sum_{\substack{n=1 \\ (n,q)=1}}^q e^{2\pi i a n/q}$.

Let a positive integer q be fixed. In this note we consider the problem of determining

$$(1) \quad \inf_{b_n} \sum_{a=1}^q \left| \sum'_n b_n e\left(\frac{an}{q}\right) \right|,$$

where $e(\alpha)$ stands for $e^{2\pi i \alpha}$, \sum'_n denotes the summation over n in the range $1 \leq n \leq q$, $(n, q) = 1$, and the infimum is taken over all sequences b_n satisfying

$$(2) \quad b_n \geq 1.$$

We note that if $b_n = 1$ for all n , the innermost sum in (1) is the Ramanujan sum

$$c_q(a) = \sum'_n e\left(\frac{an}{q}\right).$$

Using the well-known identity

$$(3) \quad c_q(a) = \frac{\varphi(q) \mu(q/(a, q))}{\varphi(q/(a, q))},$$

where φ and μ are Euler's and Möbius functions respectively (see, for example, [HW, Theorem 272]), and letting $q_0 = \prod_{p|q} p$ be the square-free kernel of q , we obtain

$$(4) \quad \begin{aligned} \sum_{a=1}^q |c_q(a)| &= \sum_{a=1}^q \frac{\varphi(q) \mu^2(q/(a, q))}{\varphi(q/(a, q))} \\ &= \varphi(q) \sum_{d|q_0} \frac{1}{\varphi(d)} \sum_{\substack{1 \leq a \leq q \\ (a, q) = q/d}} 1 = \varphi(q) \sum_{d|q_0} 1 \\ &= \varphi(q) 2^{\omega(q)}, \end{aligned}$$

where $\omega(q)$ is the number of distinct prime divisors of q . Thus (4) gives a “trivial” upper bound for (1). In fact this author was led to consider this problem while

Received by the editors March 28, 1995 and, in revised form, October 19, 1995.
 1991 *Mathematics Subject Classification*. Primary 11L03.

working on estimates for more general exponential sums in an attempt to beat this estimate by introducing a “smoothing factor” b_n which had to satisfy (2). This might seem plausible at first since if $q/(a, q) > 3$, one can easily find b_n satisfying (2) for which

$$\sum'_n b_n e\left(\frac{an}{q}\right) = 0.$$

We will show however that (1) is attained by taking $b_n = 1$ for all n . In fact, the following more general result is no more difficult.

Theorem. *Let r be a real number satisfying $r \geq 1$. Then for any sequence of complex numbers b_n we have*

$$(5) \quad \sum_{a=1}^q \left| \sum'_n b_n e\left(\frac{an}{q}\right) \right|^r \geq \left(\frac{|\sum'_n b_n|}{\varphi(q)} \right)^r \sum_{a=1}^q |c_q(a)|^r.$$

Proof. We may assume that b_n is defined for all integers n and is periodic with period q . We set

$$B = \sum'_n b_n$$

and

$$S = \sum_{a=1}^q \left| \sum'_n b_n e\left(\frac{an}{q}\right) \right|^r.$$

Using n^* to denote the multiplicative inverse of n modulo q for $(n, q) = 1$, we write

$$\begin{aligned} |B|^r \sum_{a=1}^q |c_q(a)|^r &= \sum_{a=1}^q |Bc_q(a)|^r \\ &= \sum_{a=1}^q \left| \sum'_m \left(\sum'_n b_{mn} \right) e\left(\frac{am}{q}\right) \right|^r \\ &= \sum_{a=1}^q \left| \sum'_n \sum'_m b_{mn} e\left(\frac{an^*mn}{q}\right) \right|^r. \end{aligned}$$

By the Hölder inequality, or trivially in the case $r = 1$, the last summation over n is bounded by

$$\leq \varphi(q)^{1/r'} \left(\sum'_n \left| \sum'_m b_{mn} e\left(\frac{an^*mn}{q}\right) \right|^r \right)^{1/r},$$

where r' satisfies $1/r + 1/r' = 1$. Therefore

$$\begin{aligned} |B|^r \sum_{a=1}^q |c_q(a)|^r &\leq \varphi(q)^{r/r'} \sum'_n \sum_{a=1}^q \left| \sum'_m b_{mn} e\left(\frac{an^*mn}{q}\right) \right|^r \\ &= \varphi(q)^{r/r'+1} S \\ &= \varphi(q)^r S, \end{aligned}$$

and the theorem follows. □

We observe that (5) may fail for $r < 1$. For example, taking $q = 5$, $b_1 = b_4 = 1 + 1/(2 \cos(2\pi/5))$ and $b_2 = b_3 = 1$, we obtain, by (3),

$$(6) \quad \sum_{a=1}^5 \left| \sum_{n=1}^4 b_n e\left(\frac{an}{5}\right) \right|^r = \sum_{a=1}^4 \left| \frac{1}{2 \cos(2\pi/5)} \left(e\left(\frac{a}{5}\right) + e\left(\frac{4a}{5}\right) \right) - 1 \right|^r + \left| 4 + \frac{1}{\cos(2\pi/5)} \right|^r \\ = 2 \left| \frac{\cos(4\pi/5)}{\cos(2\pi/5)} - 1 \right|^r + \left| 4 + \frac{1}{\cos(2\pi/5)} \right|^r,$$

and

$$(7) \quad \sum_{a=1}^5 |c_q(a)|^r = 4 + 4^r.$$

Thus for $r > 0$ sufficiently small (6) will be smaller than (7).

REFERENCES

- [HW] G.H. Hardy & E.M. Wright, *An introduction to the theory of numbers*, fifth edition, Oxford University Press, 1979. MR **81i**:10002

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF NEVADA, LAS VEGAS, NEVADA 89154-4020

E-mail address: bachman@nevada.edu