

THE SLICE GENUS AND THE THURSTON-BENNEQUIN INVARIANT OF A KNOT

LEE RUDOLPH

(Communicated by Ronald Stern)

ABSTRACT. For any knot $K \subset S^3$, $g_s(K) \geq (\text{TB}(K) + 1)/2$.

Let $K \subset S^3 = \partial D^4 \subset \mathbf{C}^2 \subset \mathbf{C}\mathbf{P}^2$ be a (smooth) knot. The *slice genus* $g_s(K)$ is the smallest genus of a smooth, oriented surface in D^4 with boundary K . If K is *Legendrian*, that is, everywhere tangent to the field of complex lines tangent to S^3 , then the *Thurston-Bennequin invariant* $\text{tb}(K)$ is the integer associated in the usual way (using linking numbers in S^3 , with the orientation induced by \mathbf{C}^2) to the normal linefield to K which assigns to each point of K the tangent line to K multiplied by $\sqrt{-1}$; cf. [1]. For arbitrary K , the *maximal Thurston-Bennequin invariant* $\text{TB}(K)$ is $\sup\{\text{tb}(K') : K' \subset S^3 \text{ is a Legendrian knot ambient isotopic to } K\}$.

Appealing to a theorem of Donaldson, I showed [4] that if $g_s(K) = 0$, then $\text{TB}(K) \leq -1$. The following inequality is more general; the proof requires more than Donaldson's theorem, namely, the truth of the local Thom Conjecture [2].

Theorem. $g_s(K) \geq (\text{TB}(K) + 1)/2$.

Proof. Using the construction of *entire totally-tangential C-links* given in [4], and elementary rational approximation theory, it is easily seen that, for any integer $t \leq \text{TB}(K)$, there is a rational immersion $F : \mathbf{C}\mathbf{P}^1 \rightarrow \mathbf{C}\mathbf{P}^2$, in general position, such that $A = F^{-1}(D^4) \subset \mathbf{C} \subset \mathbf{C}\mathbf{P}^1$ is a neighborhood of S^1 diffeomorphic to an annulus, $F|_A$ is an embedding, and $\partial F(A) = \partial A(K, t) \subset S^3$, where $\partial A(K, t) \subset S^3$ denotes an (embedded) annulus of type K with t twists (i.e., each component of $\partial A(K, t)$ is ambient isotopic to K , and the linking number of the components is $-t$).

Let $D_i \subset \mathbf{C}\mathbf{P}^1$, $i = 1, 2$, be the two 2-disks complementary to $\text{Int } A$. Let the smoothly immersed 2-disk $F(D_i) \subset \mathbf{C}\mathbf{P}^2 \setminus \text{Int } D^4$ have d_i doublepoints; let $S_i \subset \mathbf{C}\mathbf{P}^2$ be a smoothly immersed surface which is the union along a component of $\partial A(K, t)$ of $F(D_i)$ and a suitable smoothly embedded surface in D^4 of genus $g_s(K)$. Let n_i be the degree of S_i . Then the degree of $F(\mathbf{C}\mathbf{P}^1)$ is $n_1 + n_2$, and, by the local Thom Conjecture, $g_s(K) + d_i \geq (n_i - 1)(n_i - 2)/2$.

Let p be the intersection number of the relative cycles $F(D_1)$ and $F(D_2)$ in $\mathbf{C}\mathbf{P}^2 \setminus \text{Int } D^4$. Then $p - t = n_1 n_2$ is the intersection number of S_1 and S_2 in $\mathbf{C}\mathbf{P}^2$, and $p + d_1 + d_2 = (n_1 + n_2 - 1)(n_1 + n_2 - 2)/2$.

Received by the editors October 12, 1995.

1991 *Mathematics Subject Classification.* Primary 57M25; Secondary 14H99.

Key words and phrases. Slice genus, Thom Conjecture, Thurston-Bennequin invariant.

The author was partially supported by NSF grant DMS-9504832 and CNRS.




FIGURE 1. Left, global signs of the four types of crossings with local sign $+$. Right, changing a local sign from $-$ to $+$.

By combining these equations and inequalities, we find that $2g_s(K) - 1 \geq t$. \square

Remarks. (1) An advantage of the Theorem over the slice-Bennequin inequality is that a lower bound (however crude) for $\text{TB}(K)$, and thus for $g_s(K)$, can be read off from any (smooth or piecewise-linear) knot diagram $\mathbf{D}(K) \subset \mathbf{R}^2$ for K , without a need for preliminary braiding. The *global sign* of a doublepoint \mathbf{P} of $\mathbf{D}(K)$ is its sign ($+$ or $-$) in the usual sense (cf. Figure 1); the *writhe* $w(\mathbf{D}(K))$ is the number of doublepoints of global sign $+$, less the number of doublepoints of global sign $-$. A linear form $\ell : \mathbf{R}^2 \rightarrow \mathbf{R}$ with $2m_\ell(\mathbf{D}(K)) < \infty$ local extrema on $\mathbf{D}(K)$, of which none is a doublepoint, assigns a *local sign* $s_\ell(\mathbf{P}) \in \{+, -\}$ to each doublepoint \mathbf{P} . If $s_\ell(\mathbf{P}) = +$ for all \mathbf{P} , then $\mathbf{D}(K)$ is piecewise-smoothly isotopic to a *front* [4] and, as described in [4], $\text{TB}(K) \geq w(\mathbf{D}(K)) - m_\ell$; in general (cf. Figure 1), $\text{TB}(K) \geq w(\mathbf{D}(K)) - (m_\ell + \text{card } s_\ell^{-1}(-))$.

(2) Of course the genus $g(K)$ is greater than or equal to $g_s(K)$. Is there a direct, elementary, 3-dimensional proof that $g(K) \geq (w(\mathbf{D}(K)) - m_\ell - \text{card } s_\ell^{-1}(-) + 1)/2$?

(3) Since this note was first circulated, Kronheimer and Mrowka [3] have improved the estimate for any Legendrian knot K with $\text{TB}(K) > 0$; by using the adjunction inequality rather than just the local Thom Conjecture, they show that for any such K , $g_s(K) \geq (\text{tb}(K) + |r(K)| + 1)/2$, where $r(K)$ denotes the *rotation number* of K (cf. [1]). (The restriction that $\text{TB}(K) > 0$ may be unnecessary.)

REFERENCES

1. D. Bennequin, *Entrelacements et équations de Pfaff*, Astérisque **107-8** (1983), 87–161. MR **86e**:58070
2. P. Kronheimer & T. Mrowka, *Gauge theory for embedded surfaces, I*, Topology (1993). MR **94k**:57048
3. ———, personal communication, December 11, 1995.
4. Lee Rudolph, *An obstruction to sliceness via contact geometry and “classical” gauge theory*, Invent. Math. **119** (1995), 155–163. MR **95k**:57013
5. ———, *Quasipositivity as an obstruction to sliceness*, Bull. Amer. Math. Soc. **29** (1993), 51–59. MR **94d**:57028

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, CLARK UNIVERSITY, WORCESTER, MASSACHUSETTS 01610

E-mail address: lrudolph@black.clarku.edu