

## DIFFERENTIAL FORMS ON QUOTIENTS BY REDUCTIVE GROUP ACTIONS

MICHEL BRION

(Communicated by Roe Goodman)

ABSTRACT. Let  $X$  be a smooth affine algebraic variety where a reductive algebraic group  $G$  acts with a smooth quotient space  $Y = X//G$ . We show that the algebraic differential forms on  $X$  which are pull-backs of forms on  $Y$  are exactly the  $G$ -invariant horizontal differential forms on  $X$ .

### 1. INTRODUCTION

Let  $G$  be a reductive algebraic group of automorphisms of an affine algebraic variety  $X$  over an algebraically closed field  $k$  of characteristic zero. Denote by  $k[X]$  the algebra of regular functions on  $X$ . The subalgebra  $k[X]^G$  of  $G$ -invariants in  $k[X]$  is finitely generated; let  $Y = X//G$  be the affine algebraic variety such that  $k[Y] = k[X]^G$  and let  $\pi : X \rightarrow Y$  be the quotient map defined by the inclusion  $k[Y] \subset k[X]$ . Then we have a pull-back map  $\pi^* : \Omega^*(Y) \rightarrow \Omega^*(X)$ , where  $\Omega^*$  denotes the algebra of Kähler differential forms, see e.g. [K].

In the present note, we describe the image of  $\pi^*$  in the case where  $X$  and  $Y$  are smooth. In other words, we characterize the differential forms on  $X$  which can be written as sums of products of invariant functions and of their differentials. Clearly, such forms are  $G$ -invariant and *horizontal*, that is, their interior product with any vector field induced by the  $G$ -action is zero. Let  $\Omega_{\text{hor}}^*(X)^G$  be the space of  $G$ -invariant horizontal forms; such forms are also called *basic*.

Is any basic differential form in the image of  $\pi^*$ ? This question was raised by P. Michor in the setting of a representation of a compact Lie group on a real vector space  $V$ , and of differential forms on  $V$  with smooth coefficients (which do not coincide with Kähler differentials for the ring of smooth functions on  $V$ ; but differential forms with polynomial coefficients coincide with Kähler differentials for the ring of polynomial functions). Michor showed that the answer is negative in general, but positive for polar representations; see [M1] and [M2].

Here we obtain the following

**Theorem 1.** *Let  $G$  be a reductive group of automorphisms of a smooth, affine variety  $X$ , and let  $\pi : X \rightarrow Y$  be the quotient map. If  $Y$  is smooth, then  $\pi^* : \Omega^*(Y) \rightarrow \Omega_{\text{hor}}^*(X)^G$  is an isomorphism.*

This proves a conjecture of V. L. Popov formulated in his talk at the problem session of the conference “Algebraic Groups”, Kazimierz (Poland), May 25-June 1,

---

Received by the editors October 25, 1996 and, in revised form, January 29, 1997.

1991 *Mathematics Subject Classification.* Primary 14L30, 22E99.

1996. Some applications, and a generalization to the case of a singular quotient space, are given in §3 below. The author thanks V. L. Popov, G. Schwarz (who gave another proof of Theorem 1, in the case of a representation of a connected group) and D. Zaitsev (who gave another proof of Theorem 1, in the case where  $G$  is finite) for useful discussions. Thanks are also due to the referee for suggesting simplifications of several points in the proof.

## 2. PROOF OF THEOREM 1

Our approach is similar to the one in [B], §2. We begin with an observation concerning reflexive modules; this notion of commutative algebra can be defined as follows. Let  $A$  be a Noetherian domain, and let  $M$  be a finite  $A$ -module. Denote by  $M^\vee = \text{Hom}_A(M, A)$  the dual module. Then we have a natural map from  $M$  to its double dual  $M^{\vee\vee}$ , and  $M$  is called reflexive if this map is an isomorphism.

**Lemma 1.** *Let  $A$  be a normal Noetherian ring, let  $M, N$  be finite  $A$ -modules with  $M$  reflexive and  $N$  torsion-free, and let  $u : M \rightarrow N$  be an  $A$ -linear map. If  $u$  is an isomorphism in codimension one, then  $u$  is an isomorphism.*

*Proof.* Set  $Z = \text{Spec}(A)$  and let  $U \subset Z$  be the open subset where  $u$  is an isomorphism. Denote by  $\mathcal{M}, \mathcal{N}$  the sheaves on  $Z$  associated to  $M, N$ . Then we have a commutative square

$$\begin{array}{ccc} M & \rightarrow & \Gamma(U, \mathcal{M}) \\ \downarrow & & \downarrow \\ N & \rightarrow & \Gamma(U, \mathcal{N}) \end{array}$$

By assumption, the map  $\Gamma(U, \mathcal{M}) \rightarrow \Gamma(U, \mathcal{N})$  is an isomorphism. Moreover, the codimension of  $Z \setminus U$  in  $Z$  is at least two, and hence the map  $M \rightarrow \Gamma(U, \mathcal{M})$  is an isomorphism; see [H], Proposition 1.6. Finally, the map  $N \rightarrow \Gamma(U, \mathcal{N})$  is injective because  $N$  is torsion-free. These three facts imply our statement.  $\square$

We will check that Lemma 1 applies to the ring  $k[Y]$  and to the map  $\pi^* : \Omega^*(Y) \rightarrow \Omega_{\text{hor}}^*(X)^G$ . Observe that both  $\Omega^*(Y)$  and  $\Omega_{\text{hor}}^*(X)^G$  are  $k[Y]$ -modules, and that  $\pi^*$  is  $k[Y]$ -linear. Because  $Y$  is smooth,  $\Omega^*(Y)$  is locally free over  $k[Y]$ , and hence reflexive. Moreover, the  $k[X]$ -module  $\Omega^*(X)$  is finitely generated and torsion-free. Therefore, the same holds for the  $k[Y]$ -module  $\Omega^*(X)^G$  (see [P-V], Theorem 3.24) and for its submodule  $\Omega_{\text{hor}}^*(X)^G$ .

So it suffices to check that  $\pi^*$  is an isomorphism in codimension one. For this, we will use Luna's slice theorem, see [L1]. First recall a

**Definition.** An equivariant morphism  $f : X' \rightarrow X$  of affine  $G$ -varieties is *excellent* if

- (i)  $f$  is étale,
- (ii) the induced morphism  $f//G : X'//G := Y' \rightarrow Y$  is étale, and
- (iii) the morphism  $(f, \pi) : X' \rightarrow X \times_Y Y'$  is an isomorphism.

Let  $y \in Y$ . The fiber  $\pi^{-1}(y)$  contains a unique closed orbit, say  $G \cdot x$ , and the isotropy group  $G_x := H$  is reductive. By Luna's slice theorem, we have a locally closed smooth, affine,  $H$ -stable subvariety  $S \subset X$  which contains  $x$ , with the following properties:

- (1)  $N := T_x S$  is an  $H$ -complement to  $T_x(G \cdot x)$  in  $T_x X$ .
- (2) The canonical map  $G \times^H S \rightarrow X$  is excellent.

(3) There is an excellent  $H$ -morphism  $S \rightarrow N$  sending  $x$  to the origin in  $N$ . (Hence the induced map  $G \times^H S \rightarrow G \times^H N$  is excellent.)

We will also need the following

**Lemma 2.** *Let  $f : X' \rightarrow X$  be excellent, and set  $Y' := X'//G$ . Then both maps  $k[Y'] \otimes_{k[Y]} \Omega_{\text{hor}}^*(X)^G \rightarrow \Omega_{\text{hor}}^*(X')^G$  and  $k[Y'] \otimes_{k[Y]} \Omega^*(Y) \rightarrow \Omega^*(Y')$  are isomorphisms of  $k[Y']$ -modules.*

*Proof.* Because  $f//G$  is etale, the natural map  $u : k[Y'] \otimes_{k[Y]} \Omega^1(Y) \rightarrow \Omega^1(Y')$  is surjective (indeed, the cokernel of  $u$  is the space of relative differentials  $\Omega^1(Y'/Y)$  which vanishes because  $f//G$  is unramified). Moreover, because  $Y$  is smooth and  $f//G$  is etale,  $Y'$  is smooth, too. Therefore, the  $k[Y']$ -modules  $k[Y'] \otimes_{k[Y]} \Omega^1(Y)$  and  $\Omega^1(Y')$  are locally free of rank  $\dim(Y') = \dim(Y)$ . It follows that  $u$  is an isomorphism. This implies the second statement.

Because  $k[X'] = k[X] \otimes_{k[Y]} k[Y']$  and  $f$  is etale, there are equivariant isomorphisms  $k[Y'] \otimes_{k[Y]} \Omega^*(X) \rightarrow k[X'] \otimes_{k[X]} \Omega^*(X) \rightarrow \Omega^*(X')$ . This implies the first statement.  $\square$

Using the above and Lemma 2 twice, one reduces Theorem 1 to spaces of the form  $G \times^H N$ . Now, by Lemma 4.1 in [M1], restriction to  $N \subset G \times^H N$  induces an isomorphism  $\Omega_{\text{hor}}^*(G \times^H N)^G \simeq \Omega_{\text{hor}}^*(N)^H$ , so we can reduce to the case of representations. Moreover, writing  $N = N^H \oplus V$  as an  $H$ -representation, we can reduce to considering the action of  $H$  on  $V$ , since the extra trivial factor just tensors the algebras  $k[V]$  and  $k[V]^H$  with  $k[N^H]$  and the spaces  $\Omega_{\text{hor}}^*(V)^H$  and  $\Omega^*(V//H)$  with  $\Omega^*(N^H)$ .

Moreover, by Lemma 1, we may replace the quotient  $Y$  by any open subset  $U$  such that  $Y \setminus U$  has codimension at least two in  $Y$ . Recall that  $Y$  is stratified by slice types  $(H, N)$ ; see [L1]. If  $y \in Y$  is in the open stratum (resp. in a stratum of codimension one), then  $V//H$  is a point (resp. is one-dimensional). So we can finally reduce to the case of a reductive group  $H \subset \text{GL}(V)$ , where  $V^H = 0$  and  $\dim(V//H) \leq 1$ . Then the proof is concluded by the following lemmas.

**Lemma 3.** *In the case where  $V//H$  is a point, the spaces  $\Omega_{\text{hor}}^p(V)^H$  vanish for all  $p \geq 1$ .*

*Proof.* We may assume that  $H$  is connected. First consider the case where  $H$  is a torus. Then, because  $V//H$  is a point, all weights of  $H$  in  $V$  lie in an open half-space. So the weights of  $\wedge^p V^*$  lie in the opposite half-space. Observing that  $\Omega^p(V) \simeq S^\bullet(V^*) \otimes \wedge^p V^*$ , where  $S^\bullet(V^*)$  is the symmetric algebra of  $V^*$ , we obtain  $\Omega^p(V)^H = 0$  for  $p \geq 1$ , which implies our statement.

In the general case of reductive  $H$ , let  $T \subset H$  be a maximal torus. By the Hilbert-Mumford criterion, there exists a  $T$ -submodule  $W \subset V$  such that  $V = HW$  and that all weights of  $T$  in  $W$  lie in an open half-space. Because the map

$$\begin{aligned} f : H \times W &\rightarrow V \\ (g, w) &\mapsto gw \end{aligned}$$

is surjective, we can find  $w \in W$  such that the differential

$$\begin{aligned} df_{e,w} : \text{Lie}(H) \times W &\rightarrow V \\ (\xi, z) &\rightarrow \xi w + z \end{aligned}$$

is surjective, where  $\text{Lie}(H)$  denotes the Lie algebra of  $H$ . In other words, we have  $V = \text{Lie}(H)w + W$ . It follows that  $\wedge^p V = \sum_{q=0}^p (\wedge^{p-q} \text{Lie}(H)w) \wedge (\wedge^q W)$ . Let  $\omega \in$

$\Omega_{\text{hor}}^p(V)^H$ . Because  $\omega$  is horizontal, it vanishes on  $\sum_{q=0}^{p-1} (\wedge^{p-q} \text{Lie}(H)\omega) \wedge (\wedge^q W)$ . Moreover, restriction of  $\omega \in S^\bullet(V^*) \otimes \wedge^p V^*$  to  $W$  is a  $T$ -equivariant morphism from  $W$  to  $\wedge^p V^*$ . By the first step of the proof, this morphism vanishes on the subspace  $\wedge^p W$  of  $\wedge^p V$ . So restriction of  $\omega$  to  $W$  is trivial. Because  $V = HW$  and  $\omega$  is  $H$ -invariant, we conclude that  $\omega = 0$ .  $\square$

**Lemma 4.** *In the case where  $V//H$  is one-dimensional, the map  $\Omega^1(V//H) \rightarrow \Omega_{\text{hor}}^1(V)^H$  is an isomorphism, and moreover  $\Omega_{\text{hor}}^p(V)^H = 0$  for all  $p \geq 2$ .*

*Proof.* The algebra  $k[V]^H$  is generated by a non-constant homogeneous function  $f$ . We have  $\Omega^1(V//H) = k[f]df$  and  $\Omega^p(V//H) = 0$  for  $p \geq 2$ . By Lemma 3 and our reductions, the  $k[f]$ -linear map  $\Omega^*(V//H) \rightarrow \Omega_{\text{hor}}^*(V)^H$  is an isomorphism over the field  $k(f)$ . Because the  $k[f]$ -module  $\Omega_{\text{hor}}^*(V)^H$  is torsion-free, it follows that  $\Omega_{\text{hor}}^p(V)^H = 0$  for  $p \geq 2$  and that  $\Omega_{\text{hor}}^1(V)^H = k[f]\omega$  for some homogeneous form  $\omega$ . Let  $m$  be the degree of  $f$ , and let  $n$  be the degree of  $\omega$ . Because  $df$  is in  $k[f]\omega$ , we have  $m - 1 = rm + n$  for some integer  $r \geq 0$ . This forces  $r = 0$  and then  $\omega$  is a scalar multiple of  $df$ .  $\square$

3. SOME APPLICATIONS AND A GENERALIZATION

Here are two applications of Theorem 1 which were pointed out by V. L. Popov. First, we recover the following classical result of L. Solomon [So].

**Corollary 1.** *Let  $G \subset \text{GL}(V)$  be a finite group generated by pseudo-reflections. Then any  $G$ -invariant differential form on  $V$  is the pull-back of a differential form on  $V/G$ .*

Indeed, the quotient space  $V/G$  is smooth, and Theorem 1 applies.

Another application is an algebraic version of a recent result of P. Michor: let  $G$  be a Lie group of isometries of a smooth Riemannian manifold  $M$ . Assume that the  $G$ -action on  $M$  admits a section  $\Sigma$ , and denote by  $W(\Sigma) = N_G(\Sigma)/Z_G(\Sigma)$  the corresponding “generalized Weyl group”. Then restriction of differential forms with smooth coefficients induces an isomorphism  $\Omega_{\text{hor}}^*(M)^G \rightarrow \Omega^*(\Sigma)^{W(\Sigma)}$ ; see [M1], Theorem 3.7, and [M2], Theorem 2.

**Corollary 2.** *Let  $X$  be a smooth irreducible affine variety with an action of a reductive group  $G$ . Denote by  $H \subset G$  a principal isotropy group of  $X$ , by  $W$  the quotient group  $N_G(H)/H$ , and by  $\Sigma \subset X^H$  the closure of the subset of principal points. If the quotient  $X//G$  is smooth, then restriction to  $\Sigma$  induces an isomorphism  $\Omega_{\text{hor}}^*(X)^G \simeq \Omega_{\text{hor}}^*(\Sigma)^W$ .*

Indeed, we have a commutative diagram

$$\begin{array}{ccc} \Omega^*(X//G) & \rightarrow & \Omega^*(\Sigma//W) \\ \downarrow & & \downarrow \\ \Omega_{\text{hor}}^*(X)^G & \rightarrow & \Omega_{\text{hor}}^*(\Sigma)^W \end{array}$$

where the top horizontal map is pull-back by the morphism  $\Sigma//W \rightarrow X//G$ . But the latter is an isomorphism by the Luna-Richardson restriction theorem; see [L2], Corollaire 4, and [L-R]. So, by Theorem 1, both vertical arrows are isomorphisms.

Finally, Theorem 1 can be extended to the case where the quotient space  $Y$  may be singular. Then we have to replace the  $k[Y]$ -module of Kähler differentials  $\Omega^*(Y)$  by its double dual  $\Omega^*(Y)^{\vee\vee}$ , the module of Zariski-Lipman differential forms. By adapting the previous argument, we obtain the following

**Theorem 2.** *Let  $G$  be a reductive group acting on a smooth affine variety  $X$ , and let  $\pi : X \rightarrow Y$  be the quotient map. If no divisor in  $X$  is mapped by  $\pi$  to a subvariety of codimension at least two in  $Y$ , then  $\pi^* : \Omega^*(Y) \rightarrow \Omega_{\text{hor}}^*(X)^G$  factors through an isomorphism  $\Omega^*(Y)^{\vee\vee} \rightarrow \Omega_{\text{hor}}^*(X)^G$ .*

In particular, the Zariski-Lipman differential forms on a quotient of a smooth variety  $X$  by a finite group are exactly the invariant differential forms on  $X$ .

*Proof.* The assumption that no divisor in  $X$  is contracted by  $\pi$  implies that the  $k[Y]$ -module  $\Omega_{\text{hor}}^*(X)^G$  is reflexive, using [H] Proposition 1.6. Therefore,  $\pi^*$  factors through a map  $\Omega^*(Y)^{\vee\vee} \rightarrow \Omega_{\text{hor}}^*(X)^G$ . Furthermore, this map is an isomorphism, by Lemma 1, smoothness in codimension one of  $Y$ , and Theorem 1.  $\square$

We ignore whether the assumption that  $\pi$  contracts no divisor is necessary for Theorem 2 to hold. Observe that the Zariski-Lipman differentials cannot be replaced by Kähler differentials in this statement, as shown by the following example.

Let  $G$  be a cyclic group of order  $n$  acting on  $X = k^2$  by scalar multiplication by  $n$ -th roots of unity. Then the algebra  $k[X]^G$  is generated by all monomials of total degree  $n$  in the coordinates  $x_1, x_2$ . Therefore, the differentials of these monomials are the minimal system of homogeneous generators of the  $k[Y]$ -module  $\Omega^1(Y)$ ; this system consists of  $n + 1$  elements. But the  $k[Y]$ -module  $\Omega_{\text{hor}}^1(X)^G = \Omega^1(X)^G$  is minimally generated by  $2n$  homogeneous elements, products of monomials of degree  $n - 1$  by  $dx_1, dx_2$ . So the quotient  $\Omega^1(X)^G / \pi^* \Omega^1(Y)$  has dimension  $n - 1$ .

## REFERENCES

- [B] M. Brion: *Sur les modules de covariants*, Ann. scient. Éc. Norm. Sup. **26** (1993), p. 1-21. MR **95c**:14062
- [H] R. Hartshorne: *Stable reflexive sheaves*, Math. Ann. **254** (1980), p. 121-176. MR **82b**:14011
- [K] E. Kunz: *Kähler Differentials*, Vieweg, Braunschweig-Wiesbaden, 1986. MR **88e**:14025
- [L1] D. Luna: *Slices étales*, Bull. Soc. math. France, Mémoire **33** (1973), p. 81-105. MR **49**:7269
- [L2] D. Luna: *Adhérences d'orbites et invariants*, Invent. Math. **29** (1975), 231-238. MR **51**:12879
- [L-R] D. Luna and R. W. Richardson: *A generalization of the Chevalley restriction theorem*, Duke Math. J. **46** (1979), p. 487-496. MR **80k**:14049
- [M1] Peter W. Michor: *Basic differential forms for actions of Lie groups*, Proc. Amer. Math. Soc. **124** (1996), p. 1633-1642. MR **96g**:57041
- [M2] Peter W. Michor: *Basic differential forms for actions of Lie groups II*, Proc. Amer. Math. Soc. **125** (1997), 2175-2177. CMP 97:10
- [P-V] V. L. Popov and E. B. Vinberg: *Invariant Theory*, Encyclopaedia of Mathematical Sciences, Algebraic Geometry IV, vol. **55**, Springer-Verlag, Berlin 1994, p. 123-284. MR **92d**:14010
- [So] L. Solomon: *Invariants of finite reflection groups*, Nagoya Math. J. **22** (1963), p. 57-64. MR **27**:4872

INSTITUT FOURIER, B. P. 74, 38402 SAINT-MARTIN D'HÈRES CEDEX, FRANCE  
*E-mail address:* mbrion@fourier.ujf-grenoble.fr