

THE SHARP SOBOLEV INEQUALITY AND THE BANCHOFF-POHL INEQUALITY ON SURFACES

RALPH HOWARD

(Communicated by Christopher Croke)

ABSTRACT. Let (M, g) be a complete two dimensional simply connected Riemannian manifold with Gaussian curvature $K \leq -1$. If f is a compactly supported function of bounded variation on M , then f satisfies the Sobolev inequality

$$4\pi \int_M f^2 dA + \left(\int_M |f| dA \right)^2 \leq \left(\int_M \|\nabla f\| dA \right)^2.$$

Conversely, letting f be the characteristic function of a domain $D \subset M$ recovers the sharp form $4\pi A(D) + A(D)^2 \leq L(\partial D)^2$ of the isoperimetric inequality for simply connected surfaces with $K \leq -1$. Therefore this is the Sobolev inequality “equivalent” to the isoperimetric inequality for this class of surfaces. This is a special case of a result that gives the equivalence of more general isoperimetric inequalities and Sobolev inequalities on surfaces.

Under the same assumptions on (M, g) , if $c: [a, b] \rightarrow M$ is a closed curve and $w_c(x)$ is the winding number of c about x , then the Sobolev inequality implies

$$4\pi \int_M w_c^2 dA + \left(\int_M |w_c| dA \right)^2 \leq L(c)^2,$$

which is an extension of the Banchoff-Pohl inequality to simply connected surfaces with curvature ≤ -1 .

1. INTRODUCTION

Let (M, g) be a two dimensional Riemannian manifold, and for any domain D with compact closure in M (write this as $D \Subset M$) let $A(D)$ be the area of D and $L(\partial D)$ be the length of the boundary ∂D of D . Then it is well known that the isoperimetric inequality

$$4\pi A(D) \leq L(\partial D)^2 \quad \text{for all } D \Subset M$$

holds if and only if the Sobolev inequality

$$(1.1) \quad 4\pi \int_M f^2 dA \leq \left(\int_M \|\nabla f\| dA \right)^2$$

holds for all compactly supported real valued functions of bounded variation on M (see §2.1 below for a short discussion of functions of bounded variation). For (M, g)

Received by the editors June 21, 1996 and, in revised form, February 6, 1997.

1991 *Mathematics Subject Classification*. Primary 53C42; Secondary 53A04, 53C65.

Key words and phrases. Isoperimetric inequalities, Sobolev inequalities, Banchoff-Pohl inequality.

Euclidean space this is due to Federer and Fleming [5], and Yau [15] extended their proof to Riemannian manifolds.

Moreover, in the case of $(M, g) = (\mathbf{R}^2, dx^2 + dy^2)$ the standard plane, if c is a closed curve in \mathbf{R}^2 , $w_c(x, y)$ is the winding number of c about the point (x, y) , and $L(c)$ the length of c , then Osserman [8, p. 1194] observed that the Sobolev inequality (1.1) can be used to prove the wonderful inequality

$$4\pi \int_{\mathbf{R}^2} w_c^2 dA \leq L(c)^2$$

of Banchoff and Pohl [1].

In the hyperbolic plane with constant Gaussian curvature -1 the sharp isoperimetric inequality is

$$(1.2) \quad 4\pi A(D) + A(D)^2 \leq L(\partial D)^2$$

for all domains D with compact closure. In this note we find the Sobolev inequality equivalent to this isoperimetric inequality and use it to give the form of the Banchoff-Pohl inequality in the class of simply connected complete surfaces that have a negative upper bound on the curvature.

Theorem 1 (Sharp Sobolev Inequality). *Let (M, g) be a noncompact two dimensional Riemannian manifold (which need not be complete) and assume there are constants $a > 0$ and b so that for every domain $D \Subset M$ the isoperimetric inequality*

$$(1.3) \quad aA(D) + bA(D)^2 \leq L(\partial D)^2$$

holds. If $b < 0$ also assume

$$A(M) \leq \frac{a}{2|b|}.$$

Then, for every compactly supported f of bounded variation on M ,

$$(1.4) \quad a \int f^2 dA + b \left(\int |f| dA \right)^2 \leq \left(\int \|\nabla f\| dA \right)^2.$$

If equality holds then, up to a set of measure zero, f is a constant multiple of the characteristic function of a domain $D \Subset M$, and D makes equality hold in the isoperimetric inequality (1.3). Conversely, if the inequality (1.4) holds for all compactly supported functions of bounded variation, then the isoperimetric inequality (1.3) holds for all D with compact closure in M .

Theorem 2 (Generalized Banchoff-Pohl Inequality). *Let (M, g) be a noncompact two dimensional simply connected Riemannian (which is not assumed to be complete) and K_0 a constant. Assume the Gaussian curvature of (M, g) satisfies*

$$K \leq K_0, \quad \text{and if } K_0 > 0 \text{ then } A(M) \leq \frac{2\pi}{K_0}.$$

If $c: [a, b] \rightarrow M$ is a closed curve and $w_c(P)$ the winding number of c about $P \in M$, then

$$(1.5) \quad 4\pi \int_M w_c^2 dA - K_0 \left(\int_M |w_c| dA \right)^2 \leq L(c)^2.$$

Equality holds if and only if c is the boundary (possibly transversed more than once) of a domain in M isometric to a geodesic disk in the simply connected space of constant curvature K_0 .

A simply connected noncompact surface is diffeomorphic to the plane \mathbf{R}^2 , so the winding number w_c can be defined in the usual manner. These results apply to simply connected domains in the sphere S^2 with area $\leq 2\pi$. As these domains are not complete, assuming completeness is not natural in Theorem 2.

Our reason for working with functions of bounded variation is that it simplifies the proofs of when equality holds in the inequalities. In many proofs that a Sobolev inequality like (1.4) implies an isoperimetric inequality like (1.3) it is usual to approximate a characteristic function χ_D by smooth (or Lipschitz) functions f in (1.3) and then take limits (cf. [5, rmk. 6.6, p.487], [15], [8, p. 1194], [3, p. 97], [16, p. 81]). As with most proofs of inequalities by approximation, this makes understanding the case of equality difficult. The advantage of working with functions of bounded variation in this setting is that if D is a domain with compact closure in M and so that the boundary ∂D has finite length, then the characteristic function of D is of bounded variation and its total variation is given by $\int_M \|\chi_D\| dA = L(\partial D)$. Thus in the class of functions of bounded variation the isoperimetric inequality (1.3) can be proven by directly putting $f = \chi_D$ in the Sobolev inequality (1.4). This makes understanding the case of equality more or less straightforward. While using functions of bounded variation in problems of this type is certainly not a new idea, it deserves to be better known.

Under the assumptions that $K_0 \leq 0$ and (M, g) is simply connected and complete, B. Süßmann has independently given a proof of the inequality (1.5). His proof uses the very ingenious idea of studying the effect of the flow of the curve shortening equation on the inequality. When (M, g) is the hyperbolic plane Teufel [10] has given another generalization of the Banchoff-Pohl inequality: $4\pi \int_M w_c^2 dA + (\int_M w_c dA)^2 \leq L(c)^2$. While this inequality is sharp in that equality holds exactly when c is the boundary of a geodesic disk (possibly transversed more than once), if w_c changes sign on M , then the inequality (1.5) gives a better lower bound on $L(\partial D)^2$. For other extensions of the Banchoff-Pohl inequality to curved surfaces, see [4], [7], [10], [11], [12], [13], [14].

Notation and terminology. By smooth we mean of class C^∞ . A domain in a manifold is an open set which we do not assume is connected. If D is a domain in M then $D \Subset M$ means that the closure of D in M is compact. By convention we assume simply connected domains are connected.

2. PROOFS

2.1. Functions of bounded variation and the coarea formula. Let (M, g) be an oriented n dimensional Riemannian manifold and let dV the the volume form on M . Then for a smooth function $f: M \rightarrow \mathbf{R}$ let ∇f be the gradient of f ; that is, ∇f is the vector field so that for all tangent vectors V one has $df(V) = \langle \nabla f, V \rangle$. Let $C_0^\infty(M, T(M))$ be the space of compactly supported smooth vector fields on M with the usual inductive limit topology (that is, $\Phi_\ell \rightarrow \Phi$ iff there is a compact set that contains the supports of all the Φ_ℓ and moreover the sequence $\{\Phi_\ell\}_{\ell=1}^\infty$ and all its partial derivatives converge uniformly to the corresponding partial derivatives of Φ). If f is a locally integrable function, define a continuous linear functional on $C_0^\infty(M, T(M))$ by

$$\Lambda_f(\Phi) := - \int_M f \operatorname{div}(\Phi) dV.$$

If f is C^1 , then by the divergence theorem

$$\Lambda_f(\Phi) = \int_M \langle \nabla f, \Phi \rangle dV$$

and so when f is sufficiently smooth the linear functional Λ_f is represented by integration against the classical gradient ∇f of f . In general Λ_f can be viewed as the distributional gradient of f . A function is of **bounded variation** iff the linear functional Λ_f is represented by measures of finite total variation—that is, if and only if in a local coordinate system (x^1, \dots, x^n) on M defined on an open set U of M there are Borel measures of finite total variation μ_1, \dots, μ_n so that for any smooth vector field $\Phi = \sum \varphi^i \partial/\partial x^i$ supported in U

$$\Lambda_f(\Phi) = \int_U \sum_{i,j} g^{ij} \varphi_i d\mu_j.$$

In this case μ_j is the distributional derivative $\partial f/\partial x^j$. A function of bounded variation need not be continuous. If $D \subseteq M$ with Lipschitz boundary, then the characteristic function χ_D is of bounded variation (cf. [16, p. 229]). More generally, a set $E \subseteq M$ is of **finite perimeter** iff the characteristic function χ_E is of bounded variation. For our purposes all that matters about sets of finite perimeter is that a set E of finite perimeter has a generalized boundary $\partial^* E$ (cf. [16, p. 240]) (which agrees with the usual topological boundary when E is a domain with C^1 boundary) and $\mathcal{H}^{n-1}(\partial^* E) < \infty$, where \mathcal{H}^{n-1} is $n - 1$ dimensional Hausdorff measure.

If Φ is a vector field on M , let $|\Phi(x)| = \sqrt{\langle \Phi(x), \Phi(x) \rangle}$. The **total variation measure** $\|\nabla f\| dV$ of a function of bounded variation is defined first on non-negative real valued continuous functions u by

$$\int_M u \|\nabla f\| dV := \sup\{\Lambda_f(\Phi) : \Phi \in C_0^\infty(M, T(M)), |\Phi(x)| \leq u(x)\}$$

and then extended to arbitrary continuous functions by linearity (cf. [16, p. 221]). There is another characterization of the total variation measure of f by the version of the coarea formula due to Fleming and Rishel [6] (or cf. [16, thm. 5.4.4, p. 231, and thm. 5.8.1, p. 247]), which gives an integral formula for the total variation of f :

$$(2.1) \quad \int_M \|\nabla f\| dV = \int_0^\infty \mathcal{H}^{n-1}(\partial^*\{x : |f(x)| \geq t\}) dt.$$

(One of the conclusions of [16, thm. 5.4.4, p. 231] is that $\{x : |f(x)| \geq t\}$ is of finite perimeter for almost all $t \in \mathbf{R}$, so the integral on the right makes sense.)

In what follows we will only be interested in the two dimensional case. Then the volume measure dV will be replaced by the area measure dA , and we will denote the one dimensional Hausdorff measure of E by $L(E)$, as in the case when E is a curve $\mathcal{H}^1(E)$ is just the length of E . We also simplify the notation in (2.1) and use ∂ for ∂^* . This should not lead to any confusion. With this notation the coarea formula for functions of bounded variation on a surface becomes

$$\int_M \|\nabla f\| dA = \int_0^\infty L(\partial\{f(x) : |f(x)| \geq t\}) dt.$$

Note that this form of the coarea formula makes it clear that if $D \Subset M$ has a rectifiable boundary, then

$$\int_M \|\nabla \chi_D\| dA = L(\partial D).$$

Finally, if f is in $W^{1,1}(M)$ (that is, the distributional first derivatives of f exist and are Lebesgue integrable), then f is of bounded variation; the total variation measure is absolutely continuous with respect to the area measure on M and is given by

$$\|\nabla f\| dA = |\nabla f| dA,$$

where $|\nabla f(x)| = \sqrt{\langle \nabla f(x), \nabla f(x) \rangle}$.

Proof of Theorem 1. Let (M, g) be a noncompact two dimensional Riemannian manifold as in the statement of Theorem 1 and so that the isoperimetric inequality (1.3) holds. We use the notation

$$A(t) := A\{x \in M : |f(x)| \geq t\}, \quad L(t) := L(\partial\{x \in M : |f(x)| \geq t\}).$$

(That is, $A(t)$ is the Lebesgue measure of the set $\{x \in M : |f(x)| \geq t\}$ and $L(t)$ is the one dimensional Hausdorff measure of $\partial\{x \in M : |f(x)| \geq t\}$.) By a standard result from real analysis, for any measurable function u on M

$$\int_M |u| dA = \int_0^\infty A\{x : |u(x)| \geq t\} dt.$$

Applying this to f and f^2 gives $\int_M |f| dA = \int_0^\infty A(t) dt$ and

$$\begin{aligned} \int_M |f|^2 dA &= \int_0^\infty A\{x : f(x)^2 \geq s\} ds \\ &= \int_0^\infty A\{x : f(x)^2 \geq t^2\} 2t dt = 2 \int_0^\infty A(t)t dt. \end{aligned}$$

By the coarea formula and the isometric inequality (1.3)

$$\int_M \|\nabla f\| dA = \int_0^\infty L(t) dt \geq \int_0^\infty \sqrt{aA(t) + bA(t)^2} dt.$$

So it is enough to prove that

$$(2.2) \quad 2a \int_0^\infty A(t)t dt + b \left(\int_0^\infty A(t) dt \right)^2 \leq \left(\int_0^\infty \sqrt{aA(t) + bA(t)^2} dt \right)^2.$$

The proof now splits into two cases.

Case 1. $b \leq 0$. This case follows closely the ideas in the papers of Federer and Fleming [5] and Yau [15]. Set

$$\begin{aligned} F(s) &:= 2a \int_0^s A(t)t dt + b \left(\int_0^s A(t) dt \right)^2, \\ G(s) &:= \left(\int_0^s \sqrt{aA(t) + bA(t)^2} dt \right)^2. \end{aligned}$$

Then

$$F'(s) = 2aA(s)s + 2b \int_0^s A(t) dt A(s),$$

$$G'(s) = 2 \int_0^s \sqrt{aA(t) + bA(t)^2} dy \sqrt{aA(s) + bA(s)^2}.$$

As $A(\cdot)$ is a decreasing function, $\int_0^s A(t) dt \geq sA(s)$. Also $b \leq 0$, so

$$(2.3) \quad F'(s) \leq 2asA(s) + 2bsA(s)^2.$$

Again, since $A(\cdot)$ is decreasing and the function $A \mapsto \sqrt{aA + bA^2}$ is increasing on $[0, a/(2|b|)]$ (and by one of our assumptions $A(t) \leq a/(2|b|)$),

$$G'(s) \geq 2s(aA(s) + bA(s)^2).$$

Therefore $F'(s) \leq G'(s)$ and $F(0) = G(0)$, so $F(s) \leq G(s)$. Letting $s \rightarrow \infty$ completes the proof that the required inequality (2.2) holds and completes the proof that (1.3) implies (1.4) when $b \leq 0$. If equality holds in (1.4), then equality must hold in (2.3) for almost all $s > 0$. If s_0 is a point where equality holds and $A(s_0) > 0$, then $A(s) = A(s_0)$ for all $s \in [0, s_0]$. If s_0 is a point where $A(s_0) = 0$, then A is non-negative and monotone decreasing, so $A(s) = 0$ for $s > s_0$. Thus for some constants $c_1, c_2 > 0$ the function $A(\cdot)$ is given by $A(s) = c_1 \chi_{[0, c_2]}(s)$. Then a further chase through the definitions shows for some domain $D \Subset M$ that $f = \pm c_2 \chi_D$, where D is a domain with $A(D) = c_1$. As equality holds in (1.4), it follows that $aA(D) + bA(D)^2 = L(\partial D)^2$.

Case 2. $b \geq 0$. Set

$$H(\lambda) := \left(\int_0^\infty \sqrt{aA(t) + \lambda A(t)^2} dt \right)^2.$$

Then using the Cauchy-Schwarz inequality we estimate the derivative of $H(\cdot)$ from below:

$$H'(\lambda) = \left(\int_0^\infty \sqrt{aA(t) + \lambda A(t)^2} dt \right) \left(\int_0^\infty \frac{A(t)^2}{\sqrt{aA(t) + \lambda A(t)^2}} dt \right) \geq \left(\int_0^\infty A(t) dt \right)^2.$$

Note that the argument we used in proving (2.2) in the case $b \leq 0$ only used the fact that $A(\cdot)$ was decreasing, so we can let $b = 0$ in that inequality to get

$$(2.4) \quad H(0) = \left(\int_0^\infty \sqrt{aA(t)} dt \right)^2 \geq 2a \int_0^\infty A(t)t dt.$$

This implies for all $\lambda \geq 0$ that

$$H(\lambda) = \left(\int_0^\infty \sqrt{aA(t) + \lambda A(t)^2} dt \right)^2 \geq 2a \int_0^\infty A(t)t dt + \lambda \left(\int_0^\infty A(t) dt \right)^2.$$

Letting $\lambda = b$ in this inequality gives that (2.2) holds and completes the proof of the inequality in the case $b \geq 0$. If equality holds, then equality must hold in (2.4). But as this was proven by the same method that was used in the case $b \leq 0$, the same analysis shows that equality in (1.4) implies $f = c \chi_D$, where $D \Subset M$ makes equality hold in (1.3).

Conversely, if (M, g) is so that the Sobolev inequality (1.4) holds for all compactly supported f of bounded variation, then for a $D \Subset M$ with ∂D rectifiable the characteristic function χ_D will have bounded variation, and so letting $f = \chi_D$ in (1.4) gives the isoperimetric inequality (1.3) and completes the proof. \square

Proof of Theorem 2. Recall that by our convention a simply connected domain is also connected. The full force of the following lemma is not needed in the proof of Theorem 2, but it is of interest for its own sake. It is not hard to give examples of complete simply connected surfaces where the domain of least perimeter for a given area is either disconnected or connected but not simply connected. Thus the conclusion of the lemma that in some cases the “isoperimetric” domains must be simply connected is not vacuous.

Lemma. *Let (M, g) be a compact simply connected two dimensional Riemannian manifold such that every simply connected domain $D \Subset M$ satisfies the isoperimetric inequality (1.3), and if $b < 0$ also assume $A(M) \leq a/2|b|$. Then every $D \Subset M$ satisfies this inequality. If D is a domain so that equality holds in the inequality, then D is simply connected.*

Proof. By the classification of surfaces M is diffeomorphic to the plane \mathbf{R}^2 . Let $D_1 \Subset M$ be a connected domain in M . Let D be the domain obtained from D_1 by filling in the holes of D_1 . To be precise, a point x of M is in D if and only if there is a closed curve c in D_1 so that the winding number of c about x is non-zero. (As M is diffeomorphic to \mathbf{R}^2 , the winding number can be defined in the usual manner.) Then D is also a bounded domain in M , and it is simply connected. This D satisfies the given isoperimetric inequality. But $D_1 \subseteq D$ and $\partial D \subseteq \partial D_1$, so $A(D_1) \leq A(D)$ and $L(\partial D) \leq L(\partial D_1)$. Therefore

$$aA(D_1) + bA(D_1)^2 \leq aA(D) + bA(D)^2 \leq L(D)^2 \leq L(D_1)^2$$

as the function $A \mapsto aA + bA^2$ is increasing on the interval $[0, A(M)]$ (this is where the assumption $A(M) \leq a/2|b|$ for $b < 0$ is used). This shows that any connected domain $D_1 \Subset M$ satisfies the required inequality. Moreover as $A(D_1) = A(D)$ if and only if $D_1 = D$, we see that equality holds for a connected domain D_1 if and only if $D_1 = D$, that is, if and only if D_1 is simply connected.

It is an elementary exercise to show that, for positive real numbers A_1, A_2, L_1, L_2 with both $aA_1 + bA_1^2$ and $aA_2 + bA_2^2$ nonnegative, the implication

$$(2.5) \quad aA_1 + bA_1^2 \leq L_1^2 \quad \text{and} \quad aA_2 + bA_2^2 \leq L_2^2 \\ \text{imply} \quad a(A_1 + A_2) + b(A_1 + A_2)^2 < (L_1 + L_2)^2$$

holds. This and induction shows that the required inequality holds for all domains $D_2 \Subset M$ that are finite unions of connected domains. As any domain $D_3 \Subset M$ is a countable union of connected domains, the general case follows by an easy limit argument.

If D is so that the equality $aA(D) + bA(D)^2 = L(D)^2$ holds, then D must be connected, as otherwise D would be the disjoint union of two subdomains D' and D'' each of which satisfies the inequality (1.3). But then the implication (2.5) would imply $aA(D) + bA(D)^2 < L(D)^2$, contrary to the assumption that equality holds. But if D is connected, then, as remarked above in the “filling in the holes” argument, equality in the isoperimetric inequality implies D is simply connected. This completes the proof. \square

To prove Theorem 2 we first note if (M, g) satisfies the hypothesis of the theorem and $D \Subset M$ is simply connected, then the Euler characteristic of D is $\chi(D) = 1$. By the form of the isoperimetric inequality in the book of Burago and Zalgaller [2, thm. 2.2.1, p. 11] the domain D satisfies

$$4\pi\chi(D)A(D) - K_0A(D)^2 = 4\pi A(D) - K_0A(D)^2 \leq L(\partial D)^2.$$

Therefore by the lemma this inequality holds for all $D \Subset M$. Now let $c: [a, b] \rightarrow M$ be a rectifiable curve. Then the function $x \mapsto w_c(x)$ is of bounded variation on M and, as in [8, pp. 1194–1195],

$$\int_M \|\nabla w_c\| dA = L(c).$$

The inequality (1.5) of Theorem 2 now follows by letting $f = w_c$ and using Theorem 1.

If equality holds in (1.5), then by the assertion on when equality holds in Theorem 1 there exist a constant C and a domain $D \Subset M$ so that $w_c = C\chi_D$ and D makes equality hold in the isoperimetric inequality (1.2). By the lemma this implies D is simply connected, and therefore ∂D is connected. But then c must be ∂D , transversed one or more times in the same direction. But equality holds in the isoperimetric inequality for a simply connected domain $D \Subset M$ if and only if D is isometric to a disk in the simply connected complete surface of constant curvature K_0 (cf. [2, thm. 4.3.1, p. 33]). This completes the proof. \square

3. REMARKS AND AN OPEN PROBLEM

Let H^n be the n dimensional hyperbolic space. Then it would be interesting to find an analytic inequality “equivalent” to the isoperimetric inequality in H^n . Let ω_n be the surface area of the unit sphere S^{n-1} in \mathbf{R}^n . Let $V(r)$ be the volume of a geodesic ball of radius r in H^n and let $A(r)$ be the surface area measure of a geodesic sphere of radius r . As the geodesic balls in H^n solve the isoperimetric problem for H^n , the isoperimetric inequality in H^n is given by the relationship between $A(r)$ and $V(r)$. They are given by

$$A(r) = \omega_n \sinh^{n-1}(r), \quad V(r) = \omega_n \int_0^r \sinh^{n-1}(t) dt.$$

When $n = 3$, $A(r) = 4\pi \sinh^2(r)$ and $V(r) = 2\pi(\cosh(r) \sinh(r) - r)$. But $\cosh(r)$ and $\sinh(r)$ are rational functions in e^r , and e^r is transcendental over the field of rational functions in r . Thus in this case there is no algebraic relationship between $A(r)$ and $V(r)$. A similar argument shows there is no algebraic relationship between $A(r)$ and $V(r)$ whenever n is odd. If n is even, then both $V(r)$ and $A(r)$ are rational functions in e^r , and thus there is a polynomial relation between $V(r)$ and $A(r)$, but for $n \geq 4$ this polynomial is rather complicated as can be seen by computing it for $n = 4$. Thus it seems that the results here do not have a straightforward generalization to higher dimensions.

Problem. Find a Sobolev type inequality for functions of bounded variation on the n dimensional hyperbolic H^n space that is equivalent to the sharp isoperimetric inequality in H^n .

ACKNOWLEDGMENTS

The idea of introducing the parameter λ and taking the derivative used in Case 2 of the proof of Theorem 1 is the result of several very enjoyable conversations with Harold Shaperio. Eberhard Teufel read a version of the manuscript and made several useful comments. I also benefitted from conversations and/or correspondence with Lars Andersson, Mike Gage, and Bernd Süßmann.

REFERENCES

- [1] T. F. Banchoff and W. F. Pohl, *A generalization of the isoperimetric inequality*, Jour. Diff. Geo. **6** (1971), 175–213. MR **46**:4449
- [2] J. D. Burago and V. A. Zalgaller, *Geometric inequalities*, Grundlehren, vol. 285, Springer, Berlin, 1980. MR **89b**:52020
- [3] I. Chavel, *Eigenvalues in Riemannian geometry*, Pure and Applied Mathematics, vol. 115, Academic Press, New York, 1984. MR **86g**:58140
- [4] K. Enomoto, *A generalization of the isoperimetric inequality on S^2 and flat tori in S^3* , Proc. Amer. Math. Soc. **120** (1994), no. 2., 553–558. MR **94d**:53001
- [5] H. Federer and W. H. Fleming, *Noraml and integral currents*, Ann. Math. **72** (1960), 458–520. MR **23**:A588
- [6] W. Fleming and R. Rishel, *An integral formula for the total gradient variation*, Arch. Math. **11** (1960), 218–222. MR **22**:5710
- [7] L. Gysin, *The isoperimetric inequality for nonsimple closed curves*, Proc. Amer. Math. Soc. **118** (1993), no. 1, 197–203. MR **93f**:53063
- [8] R. Osserman, *The isoperimetric inequality*, Bull. Amer. Math. Soc. **84** (1978), no. 6, 1182–1238. MR **58**:18161
- [9] B. Süßmann, *Curve shorting and the Banchoff-Pohl inequality on surfaces of nonpositive curvature*, Preprint (1996).
- [10] E. Teufel, *A generalization of the isoperimetric inequality in the hyperbolic plane*, Arch. Math. **57** (1991), no. 5, 508–513. MR **92j**:52015
- [11] ———, *Isoperimetric inequalities for closed curves in spaces of constant curvature*, Results Math. **22** (1992), 622–630. MR **93h**:53073
- [12] ———, *On integral geometry in Riemannian spaces*, Abh. Math. Sem. Univ. Hamburg **63** (1993), no. 2, 17–27. MR **94f**:53127
- [13] J. L. Weiner, *A generalization of the isoperimetric inequality on the 2-sphere*, Indiana Univ. Math. Jour. **24** (1974), 243–248. MR **52**:1584
- [14] ———, *Isoperimetric inequalities for immersed closed spherical curves*, Proc. Amer. Math. Soc. **120** (1994), no. 2, 501–506. MR **94d**:53003
- [15] S. T. Yau, *Isoperimetric constants and the first eigenvalue of a compact manifold*, Ann. Sci. Ec. Norm. Super., Paris (4) **8** (1975), 487–507. MR **53**:1478
- [16] W. P. Ziemer, *Weakly differentiable functions*, Graduate Texts in Mathematics, vol. 120, Springer-Verlag, New York, 1989. MR **91e**:46046

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTH CAROLINA, COLUMBIA, SOUTH CAROLINA 29208

E-mail address: howard@math.sc.edu