

## A VOLUME COMPARISON THEOREM FOR FINSLER MANIFOLDS

CARLOS E. DURÁN

(Communicated by Christopher B. Croke)

ABSTRACT. Let  $(M^n, F)$  be a symmetric Finsler manifold, endowed with the Busemann volume form, and let  $D$  be its unit disk bundle endowed with the canonical symplectic volume form. It is shown that  $Vol(D) \leq C(n)Vol(M^n)$ , where  $C(n)$  is the volume of the unit disk in  $\mathbb{R}^n$ . Moreover, equality holds if and only if  $(M^n, F)$  is Riemannian.

### 1. INTRODUCTION

A *Finsler structure* on a manifold  $M^n$  is a function  $F : TM \rightarrow \mathbb{R}$  which is homogenous of degree 1, strictly convex and smooth off the zero section. The function  $F$  is to be thought of as a “norm” in each tangent space  $T_pM$ , which is not necessarily Euclidean. The Finsler manifold  $(M, F)$  is said to be *symmetric* if  $F(v) = F(-v)$ .

There has been a revived interest in the study of Finsler manifolds recently, sparked by S.S. Chern and others, (e.g., [2], [3], [9]). There are also some older, beautiful geometric studies of Finsler manifolds; for example, [1], [5], [6], [7]. A basic problem is to find computable invariants that distinguish Finsler manifolds from Riemannian manifolds; the basic invariant is the Cartan 3-tensor  $A$ , which is, in essence, simply three derivatives of the energy function along the fibers. Riemannian manifolds are characterized by  $A \equiv 0$ . (see [2]).

In this note we will find a global integral invariant of Finsler metrics that attains its maximum exactly at the set of Riemannian metrics: the ratio of the symplectic volume of the unit disk bundle to the volume of the manifold. The philosophy behind the construction is that of Busemann: Finsler geometry should be thought of as the geometry of families of convex sets in  $\mathbb{R}^n$ , parametrized by a manifold. These geometries can be put together via the associated calculus of variations, i.e. the Hamiltonian geometry of  $(TM, \omega, F)$ , where  $\omega$  is the pullback of the canonical symplectic form of  $T^*M$  via the Legendre transformation.

### 2. THE BUSEMANN VOLUME FORM

Let  $(M, F)$  be a symmetric Finsler manifold. The *Busemann volume form* of  $(M, F)$  is defined as follows: let  $e^1, \dots, e^n$  be a basis for  $T_xM$ , with dual basis

---

Received by the editors March 6, 1997.

1991 *Mathematics Subject Classification*. Primary 53C60, 53C15.

*Key words and phrases*. Riemannian geometry, Finsler geometry.

Supported by CNPq, Brasil.

$\eta_1, \dots, \eta_n$ . Given  $x \in M$ , let  $D(x) \subset \mathbb{R}^n$  be given by

$$D(x) = \{y \in \mathbb{R}^n : F(y_i e^i) \leq 1\}$$

(summation convention will be used throughout). Denote by  $V(x)$  the volume of  $D(x)$ , with respect to the standard Euclidean structure of  $\mathbb{R}^n$ . Then the Busemann volume form  $B_F$  is given by

$$B_F = \frac{C(n)}{V(x)} \eta_1 \wedge \dots \wedge \eta_n,$$

where  $C(n)$  is the volume of the unit disk in  $\mathbb{R}^n$ . It is easy to see that  $B$  does not depend on the choice of the (positively oriented) basis. The induced measure on  $M$  coincides with the Hausdorff measure given by the metric; thus  $B_F$  is in the metric sense the “right” volume form for a Finsler manifold (see [4], [9]). As usual, we define the volume of  $M$  by

$$\text{vol}(M) = \int_M B_F.$$

On the other hand, the punctured tangent bundle  $T_0M$  has a symplectic form  $\omega$ , given by the pullback of the canonical symplectic form of  $T^*M$  by the Lagrange transformation  $v \mapsto L_F(v)$ . The top form  $\frac{(-1)^n}{(2n)!} \omega^n$  is the canonical volume form on  $T_0M$ . In local coordinates  $v = (x^i, y_i \frac{\partial}{\partial x^i})$  of  $TM$ , we can write

$$L(x, y) = \frac{1}{2} \frac{\partial F^2}{\partial y_i} dx^i, \quad \frac{(-1)^n}{(2n)!} \omega^n = \det(g_{ij}) dx^1 \wedge \dots \wedge dx^n \wedge dy_1 \wedge \dots \wedge dy_n,$$

where  $g_{ij} = \frac{1}{2} \frac{\partial^2 F^2}{\partial y_i \partial y_j}$  is called the fundamental tensor of the Finsler metric  $F$ .

Let  $C(n)$  and  $c(n)$  be the volume of the unit disk and the unit sphere in  $\mathbb{R}^n$ , respectively. Then we have

**Theorem 1.** *Let  $D$  be the unit disk in  $T_0M$ , and let  $\text{Vol}(D)$  be its volume with respect to the symplectic form. Then*

$$\text{Vol}(D) \leq C(n) \text{vol}(M),$$

*with equality if and only if  $(M, F)$  is Riemannian.*

Recall that  $\omega = -d\alpha$  where  $\alpha$  is the pullback by the Lagrange transformation of the canonical 1-form of  $T^*M$ . Then the unit tangent bundle  $\{F = 1\}$  has an exact contact structure given by  $\alpha$ , and corresponding volume form  $\frac{1}{(n-1)!} \alpha \wedge (d\alpha)^{n-1}$ . We have then the corresponding theorem for the unit tangent bundle:

**Corollary.** *Let  $S$  be the unit tangent bundle  $\{F = 1\}$ , and let  $\text{Vol}(S)$  be its volume with respect to the contact structure. Then*

$$\text{Vol}(S) \leq c(n) \text{vol}(M),$$

*with equality if and only if  $(M, F)$  is Riemannian.*

### 3. PROOF OF THEOREM 1

Cover  $M$  with a full measure coordinate patch giving us coordinates  $(x, y)$  on  $TM$ . Then

$$\omega^n = \det(g_{ij}) dx^1 \wedge \dots \wedge dx^n \wedge dy_1 \wedge \dots \wedge dy_n,$$

which we write in the more compact notation  $\det(g_{ij}) dx \wedge dy$ .

Integrating over the fiber, we have

$$\int_D \omega = \int_{x \in M} \int_{D_x} \omega = \int_{x \in M} dx \int_{D(x)} \det(g_{ij})(x, y) dy.$$

Let  $D^*(x)$  be the dual of the convex set  $D(x)$ . Recall that given a convex set  $K$  in a vector space  $V$ , the *dual set*  $K^* \subset V^*$  is defined by

$$K^* = \{ \xi \in V^* : \xi(v) \leq 1 \forall v \in K \}$$

(see the beautiful book [10] for more details on duality and convex geometry in general).

Note that the map

$$y_i \partial_{x^i} \mapsto \frac{1}{2} \frac{\partial F^2}{\partial y_i}(y) dx^i$$

is a diffeomorphism between  $D(x)$  and  $D^*(x)$ , with Jacobian  $\det(g_{ij})$ . Therefore,

$$\int_{D(x)} \det(g_{ij})(x, y) dy = \int_{D^*(x)} dy = Vol(D^*(x)).$$

Integrating over  $M$ , we have

$$Vol(D) = \int_M Vol(D^*(x)) dx = \frac{1}{C(n)} \int_M C(n) Vol(D^*(x)) dx.$$

Let  $\mu(x) = Vol(D(x))Vol(D^*(x))$ . The volume product  $\mu$  is an affine invariant of the convex body  $D(x)$ , and we have Santaló's inequality (see [8] and the references therein)

$$\mu(x) \leq Vol(E)Vol(E^*) = C(n)^2,$$

where  $E$  is any ellipsoid. Moreover, equality occurs if and only if  $D(x)$  is an ellipsoid.

Then we have

$$\begin{aligned} Vol(D) &= \int_M Vol(D^*(x)) dx \\ &= \frac{1}{C(n)} \int_M \frac{C(n)\mu(x)}{Vol(D(x))} dx \\ &= \frac{1}{C(n)} \int_M \mu(x) B_F \\ &\leq C(n) vol(M), \end{aligned}$$

which proves the inequality in Theorem 1. Equality occurs if and only if  $\mu(x) \equiv C(n)^2$  which only happens if for each  $x$ ,  $F(x, y) \leq 1$  is the interior of an ellipsoid symmetric with respect to the origin, which means that the metric is Riemannian.

The corollary follows from Theorem 1 and Stokes's Theorem.

Note that the proof of Theorem 1 and the non-symmetric version of Santaló's inequality also show the following for non-symmetric Finsler manifolds:

Let  $(M^n, F)$  be a non-symmetric Finsler manifold such that the centroid of the unit disk coincides with zero. Then  $Vol(D) \leq C(n)Vol(M)$ , with equality if and only if the unit disk is an ellipsoid. Therefore such a Finsler metric is actually symmetric and Riemannian.

## REFERENCES

1. H. Akbar-Zadeh, *Sur les espaces de Finsler a courbures sectionnelles constantes*, Acad. Roy. Belg. Bull. Cl. Sci **74** (1988), 281-322. MR **91f**:53069
2. D. Bao, S.-S. Chern and Z. Shen, *An Introduction to Riemann-Finsler Geometry*, Springer-Verlag, Berlin.
3. R. Bryant, *Finsler structures on the 2-sphere satisfying  $K=1$* , Contemporary Mathematics, vol. 196, 1996, pp. 27-41. MR **97e**:53128
4. H. Busemann, *Intrinsic Area*, Ann. Math. **48** (1947), 234-267. MR **8**:573a
5. H. Busemann, *The geometry of Finsler spaces*, Bull. Amer. Math. Society **56** (1950), 5-16. MR **11**:400e
6. H. Busemann, *Geometry of Geodesics*, Academic Press, New York, 1955. MR **17**:779a
7. E. Cartan, *Sur un problème d'équivalence et la théorie des espaces métriques généralisés*, Ouvres Complètes, Vol. II, Part. III, Gauthier-Villars, Paris, pp. 1131-1153.
8. M. Meyer and A. Pajor, *On Santaló's inequality*, Geometric aspects of functional analysis, Springer Lecture Notes in Mathematics 1376, pp. 261-263. MR **90h**:52012
9. Z. Shen, *On Finsler geometry of submanifolds*, Preprint (1996).
10. A.C. Thompson, *Minkowski Geometry*, Cambridge University Press, Cambridge, 1996. MR **97f**:52001

IMPA, ESTRADA DONA CASTORINA 110, JARDIM BOTÂNICO, RIO DE JANEIRO RJ 22460-320,  
BRASIL

*Current address:* IVIC-Matematicas, Apartado 21827, Caracas 1020-A, Venezuela

*E-mail address:* [cduran@impa.br](mailto:cduran@impa.br), [cduran@cauchy.ivic.ve](mailto:cduran@cauchy.ivic.ve)