

ON THE PROJECTIVITY OF MODULE COALGEBRAS

SIU-HUNG NG

(Communicated by Ken Goodearl)

ABSTRACT. In this paper, we derive some criteria for the projectivity of a module coalgebra over a finite dimensional Hopf algebra. In particular, we show that any Hopf algebra over a field of characteristic zero is faithfully flat over its group-like subHopf algebra. Finally, we prove that if B is a finite dimensional subHopf algebra of a Hopf algebra A , then B is normal in A if and only if $AB^+ = B^+A$. This improves a result by S. Montgomery (1993).

1. PRELIMINARY

Let B be a Hopf algebra over the field k . For any left B -module M , let M^B denote the space of invariants of M , that is

$$M^B = \{x \in M \mid bx = \varepsilon(b)x \text{ for } b \in B\},$$

where ε is the counit of B . If B is finite dimensional, B^B is just the space of left integrals \int_B^l . Similarly, we write \int_B^r for the space of right integrals of B . We let ${}_M I$ denote the left B -submodule which consists of the elements $x \in M$ such that $\int_B^r x = 0$. Similarly, if M is a right B -module, I_M denotes the B -submodule which annihilates the left integrals of B .

For any left B -modules M and N , the space $\text{Hom}(M, N)$ of k -linear homomorphisms admits a natural left B -module structure given by

$$(h \cdot f)(x) = \sum_{(h)} h_1 f(S(h_2)x)$$

for $f \in \text{Hom}(M, N)$, $x \in M$, $h \in B$, where S is the antipode of B . In particular, if N is the trivial B -module k , the B -module action on $M^* = \text{Hom}(M, k)$ is given by

$$(b \rightarrow f)(x) = f(S(b)x)$$

for $f \in M^*$, $x \in M$ and $b \in B$.

A left (right) B -module C is called a left (right) B -module coalgebra if C is a coalgebra such that the diagonal map $\Delta_C : C \rightarrow C \otimes C$ and the counit $\varepsilon_C : C \rightarrow k$ are left (right) B -module maps, where k is considered as a trivial B -module. We write C^+ for the coideal $C \cap \ker \varepsilon_C$. We will call $D \subseteq C$ a B -submodule coalgebra if D is a subcoalgebra of C and is invariant under the B -action. If $X, Y \subseteq C$, recall [8] that the “wedge” $X \wedge Y$ is defined as

$$X \wedge Y = \Delta_C^{-1}(X \otimes C + C \otimes Y).$$

Received by the editors September 27, 1996 and, in revised form, April 3, 1997.
1991 *Mathematics Subject Classification*. Primary 16W30.

We define $\bigwedge^1 X = X$ and $\bigwedge^{n+1} X = X \wedge (\bigwedge^n X)$. In particular, if $C_0 \subseteq X$,

$$\sum_{n \geq 1} \bigwedge^n X = C,$$

where C_0 is the coradical of C (cf. [4],[8]).

Let C be a B -module coalgebra. M is called a left (C, B) -Hopf module if M is a left C -comodule and a left B -module such that the comodule structure map $\rho_M : M \rightarrow C \otimes M$ is a left B -module map. If C is also projective as a B -module, we simply call C a projective B -module coalgebra. The category ${}^C_B\mathcal{M}$ of all left (C, B) -Hopf modules is an abelian category.

2. PROJECTIVITY OF MODULE COALGEBRA

Proposition 1. *Let B be a finite dimensional Hopf algebra.*

- (i) *If C is a left B -module coalgebra, then C is a projective B -module if and only if ${}_C I \subseteq \ker \varepsilon_C$.*
- (ii) *If C is a right B -module coalgebra, then C is a projective B -module if and only if $I_C \subseteq \ker \varepsilon_C$.*

Proof. (i) Let us consider the B -module $\text{Hom}(C, B)$ of linear homomorphisms from C to B . As a B -module, $\text{Hom}(C, B)$ is isomorphic to $B \otimes C^*$ under the identification

$$(b \otimes f)(x) = f(x)b$$

for $f \in C^*$, $x \in C$ and $b \in B$. Let C_0^* be the trivial B -module with the same underlying space as C^* . It is easy to see that $B \otimes C_0^*$ is isomorphic to $B \otimes C^*$ under the B -module isomorphism $\phi : B \otimes C_0^* \rightarrow B \otimes C^*$ given by

$$\phi(b \otimes f) = \sum_{(b)} b_1 \otimes (b_2 \rightarrow f)$$

for $b \in B$ and $f \in C^*$. Therefore, $\phi((B \otimes C_0^*)^B) = (B \otimes C^*)^B = \text{Hom}_B(C, B)$. Since $(B \otimes C_0^*)^B = \int_B^l \otimes C^*$,

$$\text{Hom}_B(C, B) = \sum_{(\Lambda)} \Lambda_1 \otimes \Lambda_2 \rightarrow C^*.$$

where Λ is a non-zero left integral of B . Let $f \in C^*$ and $x \in C$.

$$\begin{aligned} \varepsilon_B \circ \sum_{(\Lambda)} (\Lambda_1 \otimes \Lambda_2 \rightarrow f)(x) &= \varepsilon_B(\Lambda_1) f(S(\Lambda_2)x) \\ &= f(S(\Lambda)x). \end{aligned}$$

Note that $S(\Lambda)$ is a right non-zero integral of B . Therefore, ${}_C I \subseteq \ker \varepsilon_C$ iff there exists $f \in C^*$ such that $f(S(\Lambda)x) = \varepsilon_C(x)$ for $x \in C$ which is equivalent to $\varepsilon_B \circ \sum_{(\Lambda)} (\Lambda_1 \otimes \Lambda_2 \rightarrow f) = \varepsilon_C$. By virtue of Doi's Theorem ([2], Corollary 1), the proof is completed. (ii) can be proved similarly. \square

A particular case of a result of Takeuchi ([9], Corollary 3.5) is then an immediate consequence of the above proposition.

Corollary 2. *Let $B \subseteq A$ be a Hopf algebras, with B finite dimensional. Then A is a left projective B -module iff A is a right projective B -module.*

Proof. If A is not projective as a left B -module, then by Proposition 1 there exists $a \in A$ such that $\int_B^r a = 0$ and $\varepsilon_A(a) \neq 0$. Let S be the antipode of A . Then $S(a)S(\int_B^r) = 0$ and $\varepsilon_A(S(a)) = \varepsilon_A(a) \neq 0$. Since $S(\int_B^r) = \int_B^l$, A is not a right projective B -module by Proposition 1. \square

3. PROJECTIVITY FOR HOPF ALGEBRAS OVER GROUP-LIKE SUBALGEBRAS

Let A be a Hopf algebra and B a subHopf algebra of A . Following [9], we use “ B -projective” to mean “projective B -module”.

Lemma 3. *If the antipode of B is bijective, then the following statements are equivalent :*

- (a) A is left B faithfully flat;
- (b) A is right B faithfully flat;
- (c) A is left B -projective;
- (d) A is right B -projective;
- (e) A is a left B -projective generator;
- (f) A is a right B -projective generator;
- (g) for any simple subcoalgebra C of A , BC is a projective B -module;
- (h) if $M \in {}_B^A\mathcal{M}$ and $M = BV$ for some simple left A -comodule V , then M is a projective B -module;
- (i) for $M \in {}_B^A\mathcal{M}$, M is left B -flat.

Proof. By [9], Corollary 3.5, (a) to (f) are all equivalent statements. (e) \Rightarrow (g) and (g) \Rightarrow (h) are consequences of [2], Theorem 4.

(h) \Rightarrow (i) Let $M \in {}_B^A\mathcal{M}$ and \mathcal{S} be the set of all left (A, B) -subcomodules J of M such that J is a flat left B -module. The assumption (h) assures that $\mathcal{S} \neq \emptyset$. Since flatness is preserved under direct limit, by Zorn’s Lemma there is a maximal element $J_0 \in \mathcal{S}$. We claim that $J_0 = M$. If not, there exists a simple left A -subcomodule \overline{V} of M/J_0 . Then $B\overline{V}$ is a flat B -module. Let $V \supset J_0$ be the left (A, B) -submodule of M such that $V/J_0 = B\overline{V}$. Then $V/J_0 = B\overline{V}$ and we have the exact sequence in ${}_B^A\mathcal{M}$

$$0 \longrightarrow J_0 \longrightarrow V \longrightarrow B\overline{V} \longrightarrow 0.$$

As flatness is preserved under extension, V is flat and hence $V \in \mathcal{S}$. This contradicts the maximality of J_0 . Therefore $J_0 = M$, and hence M is left B flat.

(i) \Rightarrow (a) Let N be a non-zero right B -module. By applying the functor $N \otimes_B ?$ to the exact sequence of left B -modules :

$$0 \longrightarrow B \longrightarrow A \longrightarrow A/B \longrightarrow 0$$

we have the long exact sequence

$$\cdots \longrightarrow \text{Tor}_1(N, A/B) \longrightarrow N \otimes_B B \longrightarrow N \otimes_B A \longrightarrow N \otimes_B (A/B) \longrightarrow 0.$$

By assumption (i), A and A/B are left B -flat since they are left (A, B) -modules. Therefore $\text{Tor}_1(N, A/B) = 0$, and so $N \otimes_B A \neq 0$. Hence A is left B faithfully flat. \square

Remark. If B is a Hopf subalgebra of A with bijective antipode, then by virtue of the above lemma, the adjectives “left” and “right” can be dropped. For example, we will simply say A is faithfully flat over B instead of A is left (or right) B faithfully flat.

Let $G(A)$ denote the set of all group-like elements of A . Let $G \subseteq G(A)$ be a subgroup of $G(A)$ and $B = k[G]$.

Lemma 4 ([6], Proposition 2). *If C is a simple subcoalgebra of A , then*

- (i) $G_C = \{g \in G \mid gC = C\}$ is a finite subgroup of G , and
- (ii) $BC = \bigoplus_{g \in S} gC$, where S is a set of left coset representatives of G_C in G .

Proposition 5. *The following statements are equivalent :*

- (i) A is $k[G]$ -projective;
- (ii) for any subgroup H of G , A is projective over $k[H]$;
- (iii) for any finite subgroup H of G , A is projective over $k[H]$.

Proof. (i) \Rightarrow (ii) Suppose A is left $k[G]$ -projective. Then A is a direct summand of a free $k[G]$ -module F . Since $k[G]$ is a free left $k[H]$ -module, F is a free left $k[H]$ -module. Hence A is left $k[H]$ -projective.

(ii) \Rightarrow (iii) is obvious.

(iii) \Rightarrow (i) Assume that A is left $k[H]$ -projective for any finite subgroup H of G . Let C be a simple subcoalgebra of A . By Lemma 4, $H = G_C$ is a finite subgroup of G . Consider the map $\mu : B \otimes_{k[H]} C \rightarrow BC$, $\mu : b \otimes c \mapsto bc$. Clearly, μ is left B -linear and surjective. By Lemma 4 (ii), μ is also injective and hence

$$B \otimes_{k[H]} C \cong BC$$

as B -modules. By Lemma 3, C is a projective $k[H]$ -module. Since the functor $B \otimes_{k[H]} ?$ preserves projective objects, BC is B -projective. It follows from Lemma 3 that A is B -projective. \square

Theorem 6. *Let A be a Hopf algebra over a field k and G be a subgroup of $G(A)$.*

- (i) *If $\text{char } k = 0$, A is faithfully flat over $k[G]$.*
- (ii) *If $\text{char } k = p > 0$, A is faithfully flat over $k[G]$ if and only if A is projective over $k[H]$ for any finite p -subgroup of G .*

Proof. (i) For any finite subgroup H of G , $k[H]$ is semisimple by Maschke's Theorem. Hence, every left $k[H]$ -module is projective and, in particular, A is projective over $k[H]$. By Proposition 5, A is faithfully flat over $k[G]$.

(ii) Let H be a finite subgroup of G and N a p -Sylow subgroup of H . Notice that every $k[H]$ -module M can be embedded into a free $k[H]$ -module F . If M is $k[N]$ -projective, then M is also $k[N]$ -injective and so M is a summand of F as $k[N]$ -module. Therefore, M is a summand of F as $k[H]$ -module (cf. [1], 63.7). Hence, the result follows from Proposition 5. \square

4. THE EXISTENCE OF A UNIQUE MAXIMAL PROJECTIVE MODULE SUBCOALGEBRA

Proposition 7. *Let B be a finite dimensional Hopf algebra and C a left B -module coalgebra.*

- (i) *If C_1 and C_2 are projective B -submodule coalgebras, then $C_1 \wedge C_2$ and $C_1 + C_2$ are B -projective.*
- (ii) *Let $\{C_i\}_{i \in J}$ be a family of projective B -submodule coalgebras of C . Then $\sum_{i \in J} C_i$ is also a projective B -module.*

Proof. Since B is a Frobenius algebra, C_1 is injective as a B -module. Let C'_1 be a B -submodule of C such that $C = C_1 \oplus C'_1$. Then $C_1 \otimes C + C \otimes C_2 = C_1 \otimes C \oplus C'_1 \otimes C_2$ as a B -module. Let $x \in C_1 \wedge C_2$ such that $\int_B^r x = 0$. Then $\Delta(x) = \sum_i a_i \otimes b_i + \sum_i c_i \otimes d_i$ where $\sum_i a_i \otimes b_i \in C_1 \otimes C$ and $\sum_i c_i \otimes d_i \in C'_1 \otimes C_2$. For $\Lambda \in \int_B^r$, $\Delta(\Lambda x) = 0$. Therefore,

$$\sum_{i,j} \Lambda_j a_i \otimes \Lambda'_j b_i + \sum_{i,j} \Lambda_j c_i \otimes \Lambda'_j d_i = 0,$$

where $\Delta(\Lambda) = \sum_j \Lambda_j \otimes \Lambda'_j$. Thus,

$$(1) \quad \sum_{i,j} \Lambda_j a_i \otimes \Lambda'_j b_i = 0,$$

$$(2) \quad \sum_{i,j} \Lambda_j c_i \otimes \Lambda'_j d_i = 0.$$

Applying $id \otimes \varepsilon$ to equation (1) and $\varepsilon \otimes id$ to equation (2), we have

$$\Lambda \left(\sum_i \varepsilon(b_i) a_i \right) = 0,$$

$$\Lambda \left(\sum_i \varepsilon(c_i) d_i \right) = 0.$$

By Proposition 1, $\varepsilon(\sum_i \varepsilon(b_i) a_i) = 0$ and $\varepsilon(\sum_i \varepsilon(c_i) d_i) = 0$. Thus, $\varepsilon(x) = 0$. Since $C_1 \wedge C_2$ is obviously a B -submodule coalgebra, $C_1 \wedge C_2$ is projective by Proposition 1. Obviously, C_1, C_2 are left $(C_1 \wedge C_2, B)$ -Hopf modules and so is $C_1 + C_2$. By [2](Theorem 4), $C_1 + C_2$ is a projective B -module.

(ii) Let $x \in \sum_{i \in I} C_i$ such that $\Lambda x = 0$. Then there exist C_{i_1}, \dots, C_{i_n} such that $x \in \sum_{k=1}^n C_{i_k}$. By (i), $\sum_{k=1}^n C_{i_k}$ is projective and so $\varepsilon(x) = 0$ by Proposition 1. Hence $\sum_{i \in I} C_i$ is B -projective by Proposition 1. \square

Corollary 8. *Let B be a finite dimensional Hopf algebra¹. For any left B -module coalgebra C , there exists a unique maximal projective B -submodule coalgebra $P(C)$. Moreover, $P(C)$ is co-idempotent, i.e. $P(C) \wedge P(C) = P(C)$.*

Proof. Let \mathcal{S} be the set of all projective B -submodule coalgebras of C . By Proposition 7, $P(C) = \sum_{D \in \mathcal{S}} D$ is then the largest projective B -submodule coalgebra of C . $P(C) \wedge P(C) = P(C)$ is a direct consequence of Proposition 7 (i) and the maximality of $P(C)$. \square

Corollary 9. *Let B be a finite dimensional Hopf algebra and C a B -module coalgebra. (i) If D is a B -submodule coalgebra of C , $P(D) = P(C) \cap D$. (ii) If C is a direct sum of B -submodule coalgebras $\{C_i\}$, then $P(C) = \bigoplus_i P(C_i)$.*

Proof. (i) By Corollary 8, $P(D) \subseteq P(C)$ and hence $P(D) \subseteq P(C) \cap D$. Conversely, by Theorem 4 of [2], $P(C) \cap D$ is a projective B -module. Then, we have $P(C) \cap D \subseteq P(D)$.

(ii) If $C = \bigoplus_i C_i$ as B -module coalgebra, it follows by Theorem 3 of [3] that $P(C) = \bigoplus_i (P(C) \cap C_i)$. Hence, by (i), $P(C) = \bigoplus_i P(C_i)$. \square

Corollary 10. *Let B be a finite dimensional Hopf algebra and C a B -module coalgebra. The following statements are equivalent :*

¹After this paper was written, the author was able to eliminate the finite dimension hypothesis on B using different techniques.

- (i) C is a projective B -module,
- (ii) BC_0 is B -projective, where C_0 is the coradical of C ,
- (iii) BD is B -projective for any simple subcoalgebra D of C .

Proof. (i) \Rightarrow (iii) is a direct consequence of Theorem 4 in [2].

(iii) \Rightarrow (ii) follows from Proposition 7 (ii).

(ii) \Rightarrow (i) Since BC_0 is B -projective, by Proposition 7 (i), $\bigwedge^n BC_0$ is B -projective for any $n \geq 1$. Hence, by Proposition 7 (ii), $\sum_{n \geq 1} \bigwedge^n BC_0$ is B -projective. The result follows from the fact that $C = \sum_{n \geq 1} \bigwedge^n BC_0$. \square

5. NORMAL SUBHOPF ALGEBRAS

Definition 11. Let A be any Hopf algebra, B a subHopf algebra of A and S the antipode of A .

- (1) B is *left normal* if

$$a \triangleright b = \sum a_1 b S(a_2) \in B$$

for $a \in A$ and $b \in B$.

- (2) B is *right normal* if

$$b \triangleleft a = \sum S(a_1) b a_2 \in B$$

for $a \in A$ and $b \in B$.

- (3) B is *normal* if B is left normal and right normal.

It is well known that if B is a normal subHopf algebra of A , then $AB^+ = B^+A$ (cf. [4], 3.4.2). However, the converse is open. If A is left or right faithfully flat over B , the converse is known to be true (cf. [4], 3.4.3 and [9], 4.4). In this section, we will show that the converse statement holds if B is finite dimensional which enhances the result in [4], 3.4.4.

Lemma 12. Let B be a finite dimensional Hopf algebra and C a left B -module coalgebra. Let $\eta_C : C \rightarrow C/(B^+C)$ be the canonical B -module coalgebra homomorphism. Then, $\eta_C(cI)$ is a subcoalgebra of $C/(B^+C)$. In particular, if B is a subHopf algebra of a Hopf algebra A , then $\eta_A(AI)$ is a right A -submodule coalgebra of $\eta_A(A)$.

Proof. To simplify, we write \overline{C} for $C/(B^+C)$. Clearly, \overline{C} is a left B -module coalgebra and C admits a natural left and right \overline{C} -comodule structure. Let Λ be a nonzero element in \int_B^r . Consider the map $f_C : \overline{C} \rightarrow C$ defined by

$$f_C(\eta_C(x)) = \Lambda x$$

for $x \in C$ (see [7], p3348). Clearly, the map is well-defined and is a left and right \overline{C} -comodule map. Therefore, $\ker f_C$ is a subcoalgebra of \overline{C} . Notice that $\ker f_C = \eta_C(cI)$ and hence the result follows. If B is a subHopf algebra of A , then η_A is a right A -module map. Since AI is a right A -submodule of A , $\eta_A(AI)$ is a right A -submodule of $\eta_A(A)$. \square

Corollary 13. Let B be a finite dimensional Hopf algebra and C a left B -module coalgebra. If C is a projective B -module, $cI = B^+C$.

Proof. Clearly, $B^+C \subseteq cI$. It suffices to show that $cI \subseteq B^+C$. By Proposition 1, $cI \subseteq \ker \varepsilon_C$. Therefore, $\eta_C(cI) \subseteq \ker \varepsilon_{\overline{C}}$. By Lemma 12 $\eta_C(cI)$ is also a subcoalgebra of \overline{C} . Therefore, $\eta_C(cI) = 0$ and so $cI \subseteq B^+C$. \square

Lemma 14. *Let A be a Hopf algebra and B a finite dimensional subHopf algebra of A . If A is not a left projective B -module, then*

$${}_A I I_A + AB^+ A = A.$$

Proof. By Lemma 12, ${}_A I/(B^+ A)$ is a right A -submodule coalgebra of $A/(B^+ A)$. Note that $AB^+ A$ is a Hopf ideal of A . Thus $AB^+ A/(B^+ A)$ is also a coideal of $A/B^+ A$. Therefore, ${}_A \bar{I} = ({}_A I + AB^+ A)/AB^+ A$ is a right A -submodule coalgebra of $A/AB^+ A$. In particular, ${}_A \bar{I}$ is a right ideal of $A/AB^+ A$. Since A is not a projective B -module, ${}_A I \not\subseteq A^+$ by Proposition 1. Hence ${}_A I \not\subseteq AB^+ A$, and so ${}_A \bar{I}$ is a non-zero right ideal as well as coideal of $A/AB^+ A$. Therefore, ${}_A \bar{I} = A/AB^+ A$ (see [8], p. 108, Exercise 5). Thus, we have

$${}_A I + AB^+ A = A.$$

By Corollary 2, A is not a projective right B -module. By virtue of Proposition 1 (ii), we similarly obtain

$$I_A + AB^+ A = A.$$

Hence, we have ${}_A I I_A + AB^+ A = A$. □

Theorem 15. *Let A be a Hopf algebra and B a finite dimensional subHopf algebra of A . If $AB^+ \subseteq B^+ A$ or $B^+ A \subseteq AB^+$, then A is a free left (right) B -module.*

Proof. By the Nichols-Zoeller Theorem [5], it suffices to consider the case when A is infinite dimensional. By Schneider's Theorem ([7], Theorem 2.4), it suffices to show that A is a left projective B -module. Suppose A is not a projective left B -module. By Lemma 14, we have

$${}_A I I_A + AB^+ A = A.$$

If $AB^+ \subseteq B^+ A$, then $AB^+ A = B^+ A$. Thus, ${}_A I I_A + B^+ A = A$ and so $\int_B^r A = 0$, contradiction! Similarly, if $B^+ A \subseteq AB^+$, then ${}_A I I_A + AB^+ = A$ and hence $A \int_B^l = 0$, contradiction! Therefore, A is a projective B -module. □

Corollary 16. *Let A be a Hopf algebra and B a finite dimensional subHopf algebra of A . Then*

- (i) B is left normal iff $AB^+ \subseteq B^+ A$;
- (ii) B is right normal iff $B^+ A \subseteq AB^+$;
- (iii) B is normal iff $B^+ A = AB^+$;
- (iv) If the antipode of A is bijective, B is left normal iff B is right normal.

Proof. (i) If B is left normal, it is obvious that $AB^+ \subseteq B^+ A$ (see [9], 1.4). Conversely, assume $AB^+ \subseteq B^+ A$. By Theorem 15, A is a free right B -module and hence faithfully flat. By Theorem 4.4 of [9], B is left normal. Similarly, we can prove (ii).

(iii) An immediate consequence of (i) and (ii).

(iv) A direct consequence of Theorem 15 and Corollary 4.5 of [9]. □

ACKNOWLEDGMENT

I would like to express my appreciation to Professors E. Taft and C. Weibel for their helpful suggestions and encouragement while writing this paper. I also wish to thank the referee for his or her valuable comments.

REFERENCES

- [1] C. W. Curtis and I. Reiner, Representation Theory of Finite Groups and Associative Algebras (Interscience, New York, 1962). MR **90g**:16001
- [2] Y. Doi, On the structure of relative Hopf modules, *Comm. Algebra* **11** (1983), 243-255. MR **84a**:16014
- [3] I. Kaplansky, Bialgebras, Chicago Press, 1975. MR **55**:8087
- [4] S. Montgomery, Hopf algebras and their actions on rings, CBMS Lecture Notes, vol. 82, Amer. Math. Soc., Providence, RI, 1993. MR **94i**:16019
- [5] W. D. Nichols and M. B. Zoeller, A Hopf algebra freeness theorem, *Amer. J. Math.* **111** (1989), 381-385. MR **90c**:16008
- [6] D. E. Radford, Freeness (projectivity) criteria for Hopf algebras over Hopf subalgebras, *J. Pure Appl. Algebra* **11** (1977) 15-28. MR **57**:16344
- [7] H.-J. Schneider, Some remarks on exact sequences of quantum groups. *Comm. Algebra* **21** (1993), 3337-3357. MR **94e**:17026
- [8] M. Sweedler, Hopf algebras, Benjamin, New York, 1969. MR **40**:5705
- [9] M. Takeuchi, Quotient spaces for Hopf algebras. *Comm. Algebra* **22** (1994), 2503-2523. MR **95h**:16055

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEW BRUNSWICK, NEW JERSEY 08903
Current address: Department of Mathematics, University of California at Santa Cruz, Santa Cruz, California 95064
E-mail address: shng@math.ucsc.edu