

RESOLUTION OF SINGULARITIES OF CONVOLUTIONS WITH THE GAUSSIAN KERNEL

KATHRIN BERKNER

(Communicated by Frederick W. Gehring)

ABSTRACT. We present a complete classification of the zero set of a function which is a convolution with the Gaussian kernel. In the first part, we calculate the Taylor expansion of the convolution in a critical point. In the second part, we resolve the singularity with the help of the general Newton process which yields the Puiseux expansions for the solutions. Finally, we describe the resolved singularity in terms of Hermite polynomials.

1. INTRODUCTION

We consider the zero set of the function $T_\theta[f] : \mathbb{R} \times \mathbb{R}_+ \rightarrow \mathbb{R}$ which is given by the following convolution integral:

$$(1) \quad T_\theta[f](b, a) = \frac{1}{a} \int_{-\infty}^{\infty} \theta\left(\frac{x-b}{a}\right) f(x) dx,$$

where $f \in L^2(\mathbb{R})$ and $\theta(x) = 1/\sqrt{2\pi} \exp(-\frac{1}{2}x^2)$ denotes the Gaussian function. We call the function $T_\theta[f]$ a *Gauss convolution* (because θ is a symmetric function, the function $T_\theta[f]$ satisfies the standard definition of a convolution). In a regular point of the zero set of $T_\theta[f]$, i.e. $\frac{\partial T_\theta[f]}{\partial b}(b_0, a_0) \neq 0$ or $\frac{\partial T_\theta[f]}{\partial a}(b_0, a_0) \neq 0$, the implicit function theorem gives a description of the solution of $T_\theta[f] = 0$ in a neighborhood of (b_0, a_0) as a smooth curve. This case has attracted a lot of interest in the field of signal processing and has been studied by several authors ([1, 2, 3, 9, 10]). In [7], the zero set of $T_\theta[f]$ near a critical point is studied with restrictions on f (periodic and bandlimited) in filter theory and its applications.

Even though the classification problem of zeros of $T_\theta[f]$ arises from applications in signal processing, it is located in the field of classical real analysis. The analysis of zeros of $T_\theta[f]$ that are critical points is the contents of this paper. A complete classification is given by the following main theorem:

Theorem 1.1. *Let $f \in L^2_\theta(\mathbb{R}) = \{g : \mathbb{R} \rightarrow \mathbb{R} : \int g^2(x)e^{-\frac{1}{2}x^2} dx < \infty\}$ and $(b_0, a_0) \in \mathbb{R} \times \mathbb{R}_+$ with $T_\theta[f](b_0, a_0) = 0$.*

Received by the editors May 15, 1997.

1991 *Mathematics Subject Classification.* Primary 32S45; Secondary 44A35.

Key words and phrases. Resolution of singularities, convolutions, bifurcations, orthogonal polynomials.

Supported in part by Texas ATP under grant number TATP 003604-018 and by Alexander von Humboldt-Stiftung, Germany.

- 1) If $\frac{\partial T_\theta[f]}{\partial b}(b_0, a_0) \neq 0$, then there exist neighborhoods U of b_0 and V of a_0 such that there is a unique differentiable function $h : V \rightarrow U$ with $h(a_0) = b_0$ and $T_\theta[f](h(a), a) = 0$. The solutions $(h(a), a)$ are the only zeros of $T_\theta[f]$ in $U \times V$.
- 2) If there exists an $n > 1$, $n = 2p$ or $n = 2p + 1$, such that $\frac{\partial^m T_\theta[f]}{\partial b^m}(b_0, a_0) = 0$ for all $m < n$ and $\frac{\partial^n T_\theta[f]}{\partial b^n}(b_0, a_0) \neq 0$, then there are neighborhoods U of b_0 and V of a_0 , real numbers $t_j < 0$, $t_i \neq t_j$ and $n_j \in \mathbb{N}$ for $j = 1, \dots, p$ and p convergent power series $h_j : U \rightarrow V$ with

$$h_j(b) = a_0 + t_j(b - b_0)^2 + \sum_{i=1}^{\infty} a_{i,j}(b - b_0)^{2+\frac{i}{n_j}},$$

such that $T_\theta[f](b, a) = 0$.

- (a) Let $n = 2p$ and let x_1, \dots, x_p denote the p positive roots of the Hermite polynomial of order $2p$. Then $t_j = -\frac{1}{2a_0x_j^2}$ and the elements of

$$\bigcup_{j=1}^p \{(b, h_j(b)) \mid b \in U\}$$

are the only zeros of $T_\theta[f]$ in $U \times V$.

- (b) Let $n = 2p + 1$ and let y_1, \dots, y_p denote the p positive roots of the Hermite polynomial of order $2p + 1$. Then $t_j = -\frac{1}{2a_0y_j^2}$ and the elements of

$$\{(b_0, a) \mid a \in V\} \cup \bigcup_{j=1}^p \{(b, h_j(b)) \mid b \in U\}$$

are the only zeros of $T_\theta[f]$ in $U \times V$.

- 3) If $\frac{\partial^n T_\theta[f]}{\partial b^n}(b_0, a_0) = 0$ for all $n \in \mathbb{N}$, then $f \equiv 0$ almost everywhere.

In the following, we will present a complete proof of the above classification theorem. In section 2, we begin with a summary of results on convolutions with the Gaussian kernel and its relations to the Hermite polynomials. In section 3, we collect all necessary facts on the theory of Puiseux series which provides us with a tool to resolve singularities. Section 4 contains the proof of Theorem 1.1. Finally, in section 5, we characterize piecewise C^1 -functions $f : \mathbb{R} \rightarrow \mathbb{R}$, such that the zero set of $T_\theta[f]$ has a singularity of a given type.

2. CONVOLUTIONS WITH THE GAUSSIAN KERNEL

The following equation for the partial derivatives of $T_\theta[f]$ with respect to b follows easily from interchanging differentiation and integration:

$$(2) \quad \frac{\partial^n T_\theta[f]}{\partial b^n}(b, a) = \frac{(-1)^n}{a^{n+1}} \int_{-\infty}^{\infty} \theta^{(n)}\left(\frac{x-b}{a}\right) f(x) dx.$$

Moreover, the derivatives of the Gaussian function satisfy the following recursion formula:

$$(3) \quad \theta^{(n)}(x) = - \left[(n-1)\theta^{(n-2)}(x) + x\theta^{(n-1)}(x) \right].$$

This relation yields expressions for the partial derivatives of $T_\theta[f]$ with respect to a .

Lemma 2.1. For each $n \in \mathbb{N}$, there are constants $c_{i,n} \in \mathbb{R}$ for $i = 1, \dots, n$, such that

$$(4) \quad \frac{\partial^n T_\theta[f]}{\partial a^n}(b, a) = \frac{1}{a^{n+1}} \sum_{i=1}^n c_{i,n} \int_{-\infty}^{\infty} \theta^{(2i)}\left(\frac{x-b}{a}\right) f(x) dx.$$

Moreover, $c_{n,n} = 1$ for each $n > 0$.

Proof. Equation (4) is proved by induction over n , using the recursion formula (3). \square

Corollary 2.2. Let $(b_0, a_0) \in \mathbb{R} \times \mathbb{R}_+$ and $n \in \mathbb{N}$ such that

$$\frac{\partial^m T_\theta[f]}{\partial b^m}(b_0, a_0) = 0 \text{ for all } m < n \text{ and } \frac{\partial^n T_\theta[f]}{\partial b^n}(b_0, a_0) \neq 0.$$

Then

$$\frac{\partial^{i+k} T_\theta[f]}{\partial a^i \partial b^k}(b_0, a_0) = \begin{cases} (a_0)^i \frac{\partial^n T_\theta[f]}{\partial b^n}(b_0, a_0) & \text{for all } i, k \in \mathbb{N}, \text{ where } 2i + k = n, \\ 0 & \text{for all } i, k \in \mathbb{N}, \text{ where } 2i + k < n. \end{cases}$$

The proof is a consequence of Eq.(2) and Lemma 2.1.

The Gaussian function and its derivatives are closely related to Hermite polynomials. These polynomials will play an important role in the following parts of this paper. Therefore, we summarize a few of their properties. For further reading we refer to [5] and [6].

We start with one possible definition. The *Hermite-Polynomial* $H_n(x)$ of order n is defined by

$$H_n(x) = (-1)^n e^{\frac{1}{2}x^2} \cdot \frac{d^n}{dx^n} \left(e^{-\frac{1}{2}x^2} \right).$$

Our definition differs slightly from the definition given in [5] and [6]. It will turn out that our definition is more convenient for the further application to the Gaussian function. Nevertheless, the main properties of Hermite polynomials remain true for our definition.

Polynomial expansion:

$$H_n(x) = x^n - \binom{n}{2} x^{n-2} + 1 \cdot 3 \binom{n}{4} x^{n-4} - 1 \cdot 3 \cdot 5 \binom{n}{6} x^{n-6} + \dots$$

Generating function:

$$(5) \quad e^{tx - \frac{1}{2}t^2} = \sum_{n=0}^{\infty} H_n(x) \frac{t^n}{n!}.$$

Orthogonality property:

$$\int_{-\infty}^{\infty} H_n(x) H_m(x) e^{-\frac{1}{2}x^2} dx = \delta_{n,m} n! \sqrt{2\pi}.$$

If we define $r(x) = e^{-\frac{1}{2}x^2}$, then the Hermite polynomials form a complete orthogonal system with respect to the scalar product $\langle f, g \rangle_r = \int f(x)g(x)r(x)dx$ for

$f, g \in L^2_\theta(\mathbb{R})$. In this notation, we have

$$(6) \quad \frac{\partial^n T_\theta[f]}{\partial b^n}(b, a) = \frac{1}{a^n} \langle H_n, f(a \cdot + b) \rangle_r.$$

In the following, we are interested in the zeros of $T_\theta[f]$. In order to characterize the possible shapes of the zero set, we consider the Taylor expansion of $T_\theta[f]$ in a point (b_0, a_0) with $T_\theta[f](b_0, a_0) = 0$:

$$T_\theta[f](b, a) = \sum_{i, k=0}^\infty \frac{1}{k!i!} (b - b_0)^k (a - a_0)^i \frac{\partial^{k+i} T_\theta[f]}{\partial a^i \partial b^k}(b_0, a_0).$$

In the next section, we will show how certain conditions on the coefficients of the Taylor expansion allow us to resolve a critical point of the zero set.

3. RESOLUTION OF SINGULARITIES WITH PUISEUX SERIES

Usually, there is no general method to classify the set of zeros of a two-dimensional function $w : \mathbb{R}^2 \rightarrow \mathbb{R}, w \in C^\infty(\mathbb{R}^2)$, in a neighborhood of a singular point, even if w is represented by its infinite Taylor expansion. If we know more about the partial derivatives of w in a singular point, it is sometimes possible to reduce the Taylor expansion to a form which allows a resolution of the singularity. One of the well known methods to resolve singularities is the Newton Process, see e.g. [4], which we describe next.

Let $w : \mathbb{R}^2 \rightarrow \mathbb{R}$ be in $C^\infty(\mathbb{R}^2)$ given by the convergent power series

$$w(x, y) = \sum \beta_{k,i} x^k y^i$$

with $\beta_{k,i} \in \mathbb{R}$ and $w(0, 0) = 0$ near $(0, 0)$. Under certain conditions on the coefficients $\beta_{k,i}$, we can find parametrizations of all solutions of $w(x, y) = 0$. Here we will briefly explain the main steps of this process.

Without loss of generality let $m > 0, \beta_{0,m} \neq 0$ and $\beta_{i,m} = 0 \forall i < m$. Furthermore, let $k_0 > 0, i_0 > 0$ with $\beta_{k_0, i_0} \neq 0$. If there are positive rational numbers μ_0 and ν with $\mu_0 m = \nu$ such that

$$(7) \quad w(x, y) = \sum_{k+\mu_0 i \geq \nu} \beta_{k,i} x^k y^i,$$

we can split w into a sum of a polynomial \tilde{w} and a remainder h :

$$\begin{aligned} \tilde{w}(x, y) &= \sum_{k+\mu_0 i = \nu} \beta_{k,i} x^k y^i, \\ h(x, y) &= \sum_{k+\mu_0 i > \nu} \beta_{k,i} x^k y^i. \end{aligned}$$

With the substitution $y = tx^{\mu_0}$ for $t \in \mathbb{R}$ we obtain

$$\tilde{w}(x, y) = \tilde{w}(x, tx^{\mu_0}) = x^\nu \sum_{k+\mu_0 i = \nu} \beta_{k,i} t^i = x^\nu g(t),$$

where g is a polynomial in t of order m . If g has a real root $t_0 \neq 0$, $y_0 = t_0 x^{\mu_0}$ is a solution of $\tilde{w}(x, y_0) = 0$ and a first approximation to the zeros of f . Since $\mu_0 \in \mathbb{Q}_{>0}$ there are $p_0, q_0 \in \mathbb{N}$ with $\mu_0 = \frac{p_0}{q_0}$ and $(p_0, q_0) = 1$. The substitution $x_1 = x^{\frac{1}{q_0}}$ and $y = x_1^{p_0} (t_0 + y_1)$ leads to a new power series w_1 in x_1 and y_1 with

$$w(x_1, x_1^{p_0} (t_0 + y_1)) = x_1^{\nu q_0} w_1(x_1, y_1).$$

The iterative substitution process $y_i = x_i^{\mu_i}(t_i + y_{i+1})$ with $\mu_i = \frac{p_i}{q_i} \in \mathbb{Q}$ ($i = 0, 1, \dots$) gives the solution

$$y = t_0x^{\mu_0} + t_1x^{\mu_0+\frac{\mu_1}{q_0}} + t_2x^{\mu_0+\frac{\mu_1}{q_0}+\frac{\mu_2}{q_0q_1}} + \dots$$

In [4] it is proved that there is an index i_0 , such that $\mu_i \in \mathbb{N}$ for all $i \geq i_0$. Hence y is presented as a power series in $x^{\frac{1}{n}}$ where $n = q_0q_1 \dots q_{i_0}$. The power series

$$y(x) = \sum_{i=n\mu_0} \alpha_i \left(x^{\frac{1}{n}}\right)^i$$

is called *Puiseux expansion* for the curve with equation $f(x, y) = 0$. The convergence of the Puiseux expansion follows from the Weierstrass preparation theorem (see e.g. [4]). It is important to mention that we get the complete zero set of f if we determine the Puiseux expansions for all real roots of the polynomial g .

4. PROOF OF THEOREM 1.1

In this section we present the proof of the Main Theorem 1.1 stated in the introduction.

Ad 1) This is a consequence of the implicit function theorem.

Ad 3) By Equation (6), the condition $\frac{\partial^n T_\theta[f]}{\partial b^n}(b_0, a_0) = 0$ for all $n \in \mathbb{N}$ can be rewritten as $\langle H_n, f(a \cdot + b) \rangle_r = 0$ for all $n \in \mathbb{N}$. Since $\{H_n\}$ is complete, it follows that $f(a_0x + b_0)$ is orthogonal to all Hermite polynomials, which implies that f is equal to zero except on a set of measure zero.

Ad 2) We will start with some necessary lemmata.

Lemma 4.1. *Let $k \in \mathbb{N}$, and let*

$$P_k(x, y) = \frac{1}{k!}y^k + \frac{1}{(k-1)!2!}x^2y^{k-1} + \dots + \frac{1}{(2k-2)!}x^{2k-2}y + \frac{1}{(2k)!}x^{2k}$$

and

$$Q_k(x, y) = \frac{1}{k!}xy^k + \frac{1}{(k-1)!3!}x^3y^{k-1} + \dots + \frac{1}{(2k-1)!}x^{2k-1}y + \frac{1}{(2k+1)!}x^{2k+1}.$$

Then

$$y^{2k}P_k\left(1, -\frac{1}{2y^2}\right) = \frac{1}{(2k)!}H_{2k}(y)$$

and

$$y^{2k+1}Q_k\left(1, -\frac{1}{2y^2}\right) = \frac{1}{(2k+1)!}H_{2k+1}(y).$$

Proof. Using Equation (5) we set

$$f_t(y) := e^{ty - \frac{1}{2}t^2} = \sum_{n=0}^{\infty} H_n(y) \frac{t^n}{n!}.$$

Now we split f_t into its odd and even part f_t^e and f_t^o , respectively, and get

$$f_t^e(y) = \frac{1}{2} [f_t(y) + f_t(-y)] \quad \text{and} \quad f_t^o(y) = \frac{1}{2} [f_t(y) - f_t(-y)].$$

Furthermore,

$$(8) \quad \begin{aligned} f_t^e(y) &= \frac{1}{2} [e^{ty} + e^{-ty}] e^{-\frac{1}{2}t^2} = \sum_{n=0}^{\infty} H_{2n}(y) \frac{t^{2n}}{(2n)!} \quad \text{and} \\ f_t^o(y) &= \frac{1}{2} [e^{ty} - e^{-ty}] e^{-\frac{1}{2}t^2} = \sum_{n=0}^{\infty} H_{2n+1}(y) \frac{t^{2n+1}}{(2n+1)!}. \end{aligned}$$

The product of the power series of $\cosh x$ and e^y is absolute convergent. Therefore, we can arrange the summation in the following way:

$$\begin{aligned} &(\cosh x) e^y \\ &= \left(1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \frac{x^6}{6!} + \dots\right) \left(1 + y + \frac{y^2}{2!} + \frac{y^3}{3!} + \frac{y^4}{4!} + \frac{y^5}{5!} + \frac{y^6}{6!} + \dots\right) \\ &= (y+1) + \left(\frac{1}{2!}y^2 + \frac{1}{2!}x^2y + \frac{1}{4!}x^4\right) + \\ &\quad \left(\frac{1}{3!}y^3 + \frac{1}{2!2!}x^2y^2 + \frac{1}{4!}x^4y + \frac{1}{6!}x^6\right) + \dots \\ &= \sum_{k=0}^{\infty} P_k(x, y) \end{aligned}$$

For $s \in \mathbb{R}$ it follows:

$$(9) \quad \sum_{k=0}^{\infty} P_k(x, sx^2) = \sum_{k=0}^{\infty} x^{2k} P_k(1, s) = (\cosh x) e^{sx^2} = \frac{1}{2} [e^x + e^{-x}] e^{sx^2}.$$

By substituting $s = -\frac{1}{2y^2}$ and $x = ty$, the above equation together with Equation (8) yields:

$$\sum_{k=0}^{\infty} (ty)^{2k} P_k(1, -\frac{1}{2y^2}) = \sum_{k=0}^{\infty} H_{2k}(y) \frac{t^{2k}}{(2k)!}.$$

Now we get the formula

$$y^{2k} P_k(1, -\frac{1}{2y^2}) = \frac{1}{(2k)!} H_{2k}(y)$$

which proves the assertion.

In order to prove the second equation in Lemma 4.1 we consider the product of the power series of $\sinh x$ and e^y and obtain

$$\begin{aligned} &(\sinh x) e^y \\ &= \left(x + \frac{x^3}{3!} + \frac{x^5}{5!} + \frac{x^7}{7!} + \dots\right) \left(1 + y + \frac{y^2}{2!} + \frac{y^3}{3!} + \frac{y^4}{4!} + \frac{y^5}{5!} + \frac{y^6}{6!} + \dots\right) \\ &= \sum_{k=0}^{\infty} Q_k(x, y) \end{aligned}$$

By analogy with Equation (9) we obtain for $s \in \mathbb{R}$:

$$\sum_{k=0}^{\infty} Q_k(x, sx^2) = \sum_{k=0}^{\infty} x^{2k+1} Q_k(1, s) = (\sinh x) e^{sx^2} = \frac{1}{2} [e^x - e^{-x}] e^{sx^2}.$$

The substitutions $s = -\frac{1}{2y^2}$ and $x = ty$ lead to

$$\sum_{k=0}^{\infty} (ty)^{2k+1} Q_k(1, -\frac{1}{2y^2}) = \sum_{k=0}^{\infty} H_{2k+1}(y) \frac{t^{2k+1}}{(2k+1)!}$$

and thus

$$y^{2k+1} Q_k(1, -\frac{1}{2y^2}) = \frac{1}{(2k+1)!} H_{2k+1}(y).$$

□

Using this lemma we can determine the roots of the following polynomials in one variable: For $c \in \mathbb{R}_+$ let

$$\begin{aligned} g_{1,p}(t) &= \frac{1}{p!} (ct)^p + \frac{1}{(p-1)!2!} (ct)^{p-1} + \\ (10) \quad &\dots + \frac{1}{2!(2p-4)!} (ct)^2 + \frac{1}{(2p-2)!} ct + \frac{1}{(2p)!} \end{aligned}$$

and

$$\begin{aligned} g_{2,p}(t) &= \frac{1}{p!} (ct)^p + \frac{1}{(p-1)!3!} (ct)^{p-1} + \\ (11) \quad &\dots + \frac{1}{2!(2p-3)!} (ct)^2 + \frac{1}{(2p-1)!} ct + \frac{1}{(2p+1)!}. \end{aligned}$$

The roots of $g_{1,p}$ and $g_{2,p}$ are related to roots of the Hermite polynomials of order $2p$ and $2p+1$, because $g_{1,p}(\frac{x}{c}) = P_p(1, x)$ and $g_{2,p}(\frac{x}{c}) = Q_p(1, x)$. The roots of orthogonal polynomials are very well studied, and we can apply the following theorem about numbers of roots of Hermite polynomials (see e.g. [5]).

Theorem 4.2. *The Hermite polynomial of order n has exactly n distinct real roots.*

Theorem 4.2 and Lemma 4.1 give the following corollary:

Corollary 4.3. *The polynomials $g_{1,p}$ and $g_{2,p}$ from Equation (10) and (11), respectively, have exactly p negative roots. More precisely, the roots of $g_{1,p}$ are given by $-\frac{1}{2cx_i^2}, i = 1, \dots, p$, where x_i are the p positive roots of H_{2p} . The roots of $g_{2,p}$ are given by $-\frac{1}{2cy_i^2}, i = 1, \dots, p$, where y_i are the p positive roots of H_{2p+1} .*

In order to prove the main part 2 of Theorem 1.1 we first consider the more general function $F_l \in C^\infty(\mathbb{R}^2)$ for $l \in \mathbb{N}$ with

$$(12) \quad F_l(x, y) = \sum_{\{k,i \mid k+2i=l\}} \frac{1}{k!i!} c^i x^k y^i + \sum_{\{k,i \mid k+2i>l\}} \alpha_{i,k} x^k y^i,$$

where $c \in \mathbb{R}_+$ and $\alpha_{i,k} \in \mathbb{R}$. The resolution of the singularity $(0, 0)$ into its Puiseux expansions is given by the following theorem:

Theorem 4.4. *For $p \in \mathbb{N}$, let x_1, \dots, x_p be the p distinct positive roots of the Hermite polynomial of order $2p$ and y_1, \dots, y_p the p distinct positive roots of the Hermite polynomial of order $2p+1$. Furthermore, let U be an open subset of \mathbb{R}^2 with $0 \in U$. Then there exist open neighborhoods $V, W \subset \mathbb{R}$ of 0, $V \times W \subset U$, and p convergent power series $h_j : V \rightarrow W$ and $n_j \in \mathbb{N}$ for $j = 1, \dots, p$ with*

$$h_j(x) = t_j x^2 + \sum_{i=1}^{\infty} a_{i,j} x^{2+\frac{i}{n_j}}$$

and $t_j < 0$ for all $j = 1, \dots, p$, $t_i \neq t_j$, such that $F_l(x, h_j(x)) = 0$.

We distinguish two cases.

1. Let $l = 2p$ and $t_j = -\frac{1}{2cx_j^2}$. Then $\{(x, h_j(x)) \mid x \in V\}$ is exactly the set of solutions of $F_l(x, y) = 0$ in $V \times W$.
2. Let $l = 2p + 1$ and $t_j = -\frac{1}{2cy_j^2}$. There exist $\rho \in \mathbb{Q}$, $\rho > 0$, and a convergent power series $F_1(x, y)$, such that $F_l(x, y) = x^\rho F_1(x, y)$ and $\{(x, h_j(x)) \mid x \in V\}$ is exactly the set of solutions of $F_1(x, y) = 0$ in $V \times W$.

Proof. In the following we will resolve the singular point $(0, 0)$ of F_l by applying the Newton Process. We consider the cases l odd and l even separately.

1. Let $l = 2p$ for $p \in \mathbb{N}$. For $\mu_0 = 2$ and $\nu = 2p$, we can split the function F_l into the sum of a polynomial \tilde{F}_l and a remainder \tilde{H}_l (cf. Equation (7)). The approximation \tilde{F}_l is given by

$$\begin{aligned} \tilde{F}_l(x, y) &= \frac{1}{p!}(cy)^p + \frac{1}{(p-1)!2!}x^2(cy)^{p-1} + \dots \\ &\quad + \frac{1}{2!(2p-4)!}x^{2p-4}(cy)^2 + \frac{1}{(2p-2)!}x^{2p-2}cy + \frac{1}{(2p)!}x^{2p} \end{aligned}$$

and after substituting $y = tx^2$ we get $\tilde{F}_l(x, y) = x^{2p}g(t)$ with

$$\begin{aligned} g(t) &= \frac{1}{p!}(ct)^p + \frac{1}{(p-1)!2!}(ct)^{p-1} + \dots \\ &\quad + \frac{1}{2!(2p-4)!}(ct)^2 + \frac{1}{(2p-2)!}ct + \frac{1}{(2p)!} \end{aligned}$$

Since g is the polynomial $g_{1,p}$ from Equation (10), we know from Corollary 4.3 that g has p negative roots t_1, \dots, t_p with $t_j = -\frac{1}{2cx_j^2}$, where x_i are the positive roots of H_{2p} . Therefore, the zero set of F_l is parametrized by the Puiseux expansions

$$y_j(x) = \sum_{i=2n_j} a_{i,j} \left(x^{\frac{1}{n_j}}\right)^i$$

with $n_j \in \mathbb{N}$ and $a_{2n_j,j} = t_j$; this means $y_j(x) = t_j x^2 + O\left(x^{2+\frac{1}{n_j}}\right)$.

2. Let $l = 2p + 1$ for $p \in \mathbb{N}$. If we set $\mu_0 = 2$ and $\nu = 2p + 1$, then $\frac{l}{2} = \frac{2p+1}{2} \notin \mathbb{N}$, and therefore we cannot start the Newton process.

a) Assume that $\alpha_{0,i} = 0$ for all $i \in \mathbb{N}$. Then

$$F_l(x, y) = xF_l^*(x, y) \quad \text{with} \quad F_l^*(x, y) = \sum_{k+2i \geq 2p} \gamma_{k+1,i} x^k y^i,$$

where $\gamma_{k,i} = \frac{1}{k!i!} \alpha_{k,i}$, and the condition $\frac{\mu_0}{\nu} \in \mathbb{N}$ is valid for $\mu_0 = 2$ and $\nu = 2p$. The first approximation \tilde{F}_l^* is presented as

$$\begin{aligned} \tilde{F}_l^*(x, y) &= \frac{1}{p!}(cy)^p + \frac{1}{(p-1)!3!}x^2(cy)^{p-1} + \dots \\ &\quad + \frac{1}{2!(2p-3)!}x^{2p-4}(cy)^2 + \frac{1}{(2p-1)!}x^{2p-2}cy + \frac{1}{(2p+1)!}x^{2p}. \end{aligned}$$

The solutions of $F_l^*(x, y) = 0$ are determined in the same way as in 1), except that the t_j are the roots of the polynomial

$$g(t) = \frac{1}{p!}(ct)^p + \frac{1}{(p-1)!3!}(ct)^{p-1} + \dots + \frac{1}{2!(2p-3)!}(ct)^2 + \frac{1}{(2p-1)!}ct + \frac{1}{(2p+1)!}.$$

Since g is the polynomial $g_{2,p}$ from Equation (11), we know from Corollary 4.3 that g has p negative roots t_1, \dots, t_p with $t_j = -\frac{1}{2cy_i^2}$, where x_i are the positive roots of H_{2p+1} .

b) There exists $m \in \mathbb{N}$, such that $\alpha_{0,m} \neq 0$ and $\alpha_{0,i} = 0$ for all $i < m$. For $\gamma_{k,i} = \frac{1}{k!i!}\alpha_{k,i}$ we get

$$F_l(x, y) = \sum_{k+\mu_0 i \geq \nu} \gamma_{k,i} x^k y^i \quad \text{with} \quad \mu_0 = \frac{1}{m-p} \quad \text{and} \quad \nu = \frac{m}{m-p}.$$

The substitutions $y = x^{\frac{1}{m-p}} y_1$ and $x_1 = x^{\frac{1}{m-p}}$ yield

$$\begin{aligned} F_l(x, y) &= F_l(x, x^{\frac{1}{m-p}} y_1) = F_l(x_1^{m-p}, x_1 y_1) \\ &= \gamma_{1,p} x_1^{m-p} x_1^p y_1^p + \dots + \gamma_{2p-1,1} x_1^{(2p-1)(m-p)} x_1 y_1 \\ &\quad + \gamma_{2p+1,0} x_1^{(2p+1)(m-p)} + \gamma_{0,m} x_1^m y_1^m + \dots \\ &= x_1^m \left[\gamma_{1,p} y_1^p + \gamma_{3,p-1} x_1^{2(m-p)-1} y_1^{p-1} + \gamma_{5,p-2} x_1^{2[2(m-p)-1]} y_1^{p-2} + \dots \right. \\ &\quad \left. \dots + \gamma_{2p-1,1} x_1^{(p-1)[2(p-m)-1]} y_1 + \gamma_{2p+1,0} x_1^{p[2(p-m)-1]} + \gamma_{0,m} y_1^m + \dots \right] \\ &= x_1^m F_l^*(x_1, y_1). \end{aligned}$$

The power series F_l^* satisfies the assumptions of the Newton process ($\mu_1 = 2(m-p) - 1$, $\nu_1 = p[2(m-p) - 1]$), and for the first approximation \tilde{F}_l^* we get:

$$\begin{aligned} \tilde{F}_l^*(x_1, y_1) &= \gamma_{1,p}(cy_1)^p + \gamma_{3,p-1} x_1^{\mu_1} (cy_1)^{p-1} + \dots \\ &\quad + \gamma_{2p-1,1} x_1^{(p-1)\mu_1} cy_1 + \gamma_{2p+1,0} x_1^{p\mu_1}. \end{aligned}$$

If we substitute $y_1 = tx_1^{\mu_1}$, we obtain $\tilde{F}_l^*(x_1, y_1) = x_1^{p\mu_1} g(t)$, where g is the polynomial $g_{2,p}$ from Equation (11). Therefore, the zero set of F_l^* is parametrized by the Puiseux expansions

$$y_{1,j}(x) = \sum_{i=n_j \mu_1} a_{i,j} \left(x_1^{\frac{1}{n_j}} \right)^i,$$

where $n_j \in \mathbb{N}$ and $a_{2n_j,j} = t_j$. In terms of x and y the solutions of $F_l^*(x, y) = 0$ are given by

$$\begin{aligned} y_j(x) &= \sum_{i=n_j \mu_1} a_{i,j} x^{\frac{n_j+i}{n_j(m-p)}} \\ &= t_j x^2 + O\left(x^{2+\frac{1}{n_j(m-p)}}\right). \end{aligned}$$

□

We are now able to prove part 2 of the Main Theorem. Let (b_0, a_0) be a singular zero of $T_\theta[f]$ such that

$$\frac{\partial^m T_\theta[f]}{\partial b^m}(b_0, a_0) = 0 \quad \text{for all } m < n \quad \text{and} \quad \frac{\partial^n T_\theta[f]}{\partial b^n}(b_0, a_0) \neq 0.$$

Using the information about the partial derivatives of $T_\theta[f]$ from Corollary 2.2 we reduce the Taylor expansion of $T_\theta[f]$ in (b_0, a_0) to

$$\begin{aligned} T_\theta[f](b, a) &= \sum_{k+2i=n}^{\infty} \frac{1}{k!i!} (a_0)^i (b - b_0)^k (a - a_0)^i \frac{\partial^n T_\theta[f]}{\partial b^n}(b_0, a_0) + \\ &\quad \sum_{k+2i>n}^{\infty} \frac{1}{k!i!} (b - b_0)^k (a - a_0)^i \frac{\partial^{k+i} T_\theta[f]}{\partial a^i \partial b^k}(b_0, a_0). \end{aligned}$$

The Taylor expansion has exactly the form F_l for $l = n$. Therefore, part 2 of Theorem 1.1 follows from Theorem 4.4.

5. FURTHER RELATIONS TO HERMITE POLYNOMIALS

First, we give an example of a class of functions $f : \mathbb{R} \rightarrow \mathbb{R}$, such that the zero set of $T_\theta[f]$ has a singularity of type (a) of (b) in Theorem 1.1.

Example 5.1. Let $(b_0, a_0) \in \mathbb{R} \times \mathbb{R}_+$, $n \in \mathbb{N}$ and

$$f(a_0x + b_0) = \sum_{i=n}^{\infty} c_i H_i(x),$$

a convergent Hermite series with $c_i \in \mathbb{R}$, $c_n \neq 0$. From Equation (6) we get

$$\frac{\partial^m T_\theta[f]}{\partial b^m}(b_0, a_0) = \frac{1}{a_0^m} \int \left[c_n H_n(x) + \sum_{i=n+1}^{\infty} c_i H_i(x) \right] H_m(x) e^{-\frac{1}{2}x^2} dx = 0$$

for all $m < n$ and

$$\frac{\partial^n T_\theta[f]}{\partial b^n}(b_0, a_0) = \frac{1}{a_0^n} \int \left[c_n H_n(x) + \sum_{i=n+1}^{\infty} c_i H_i(x) \right] H_n(x) e^{-\frac{1}{2}x^2} dx = \frac{1}{a_0^n} c_n n! \sqrt{2\pi}.$$

Thus, for each $n \in \mathbb{N}$ there exist functions $f \in L^2_\theta(\mathbb{R})$ with the desired type of singularity.

Conversely, we can ask whether a given bifurcation point of type (a) or (b) in Theorem 1.1 determines the underlying function f . To answer this question we use the following theorem (see e.g. [8]):

Theorem 5.2. *Let $f \in L^2_\theta(\mathbb{R})$ be continuous with a piecewise continuous derivative $f' \in L^2_\theta(\mathbb{R})$. Then $f(x) = \sum_{n=0}^{\infty} c_n H_n(x)$ for all $x \in \mathbb{R}$, where $c_n = \frac{1}{n! \sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) H_n(x) e^{-\frac{1}{2}x^2} dx$. Moreover, the sequence $f_n(x) = \sum_{k=0}^n c_k H_k(x)$ converges uniformly to f on each interval $[x_1, x_2] \subset \mathbb{R}$.*

From this theorem we derive the following lemma:

Lemma 5.3. *Let $f \in L^2_\theta(\mathbb{R})$ be continuous and $f' \in L^2_\theta(\mathbb{R})$ be piecewise continuous. Then $\frac{\partial^m T_\theta[f]}{\partial b^m}(b, a) = 0$ for $m < n$, $\frac{\partial^n T_\theta[f]}{\partial b^n}(b, a) \neq 0$ iff $f(ax + b) = \sum_{i=n}^{\infty} c_i^{a,b} H_i(x)$, where $c_i^{a,b} \in \mathbb{R}$, $c_n^{a,b} \neq 0$.*

ACKNOWLEDGMENT

The author would like to thank Fritz von Haeseler, Heinz-Otto Peitgen and Raymond O. Wells, Jr. for many discussions on various parts of this paper. Part of this work was done while the author was working at the Center for Complex Systems and Visualization, University of Bremen. During that time she was a member of the Forschergruppe *Dynamische Systeme*, supported by the DFG.

REFERENCES

- [1] Anh, V.; Shi, J.Y.; Tsui, H.T. *Scaling Theorems for Zero Crossings of Bandlimited Signals*, IEEE Trans. Patt. Anal. Mach. Intell. Vol. 18, No. 3, 1996, pp. 309-320.
- [2] Babaud, J. et al. *Uniqueness of the Gaussian Kernel for Scale-Space Filtering*, IEEE Trans. Patt. Anal. Mach. Intell. Vol.PAMI-8, No.1, 1986, pp. 26-33.
- [3] Berkner, K. *Eine Charakterisierung der Nullstellen-Mengen von Gauß-Faltungen und eine daraus resultierende lokale Beschreibung von Kanten in Bildern*, Dissertation, Universität Bremen, 1996.
- [4] Brieskorn, E.; Knörrer, H. *Plane Algebraic Curves*, Birkhäuser Verlag, Basel, 1986. MR **88a**:14001
- [5] Chihara, T. S. *An Introduction to Orthogonal Polynomials*, Gordon and Breach, Science Publishers, New York, 1978. MR **58**:1979
- [6] Freud, G. *Orthogonale Polynome* Birkhäuser Verlag, Basel, 1969. MR **58**:1982
- [7] Johansen, P. et al. *Representing Signals by Their Topoints in Scale Space*, Proc. 8th Int. Conf. Pattern Recogn., Vol. 1, 1986, pp. 215-217.
- [8] Nikiforov, A.F.; Uvarov, V.B. *Special Functions of Mathematical Physics*, Birkhäuser Verlag, Basel, 1988. MR **89h**:33001
- [9] Wu, L.; Xie, Z. *Scaling Theorems for Zero Crossings*, IEEE Trans. Patt. Anal. Mach. Intell. Vol. 12, No.1, 1990, pp. 46-54.
- [10] Yuille, A.L.; Poggio, T.A. *Scaling Theorems for Zero Crossings*, IEEE Trans. Patt. Anal. Mach. Intell. Vol. 8, No.1, 1986, pp. 15-25.

DEPARTMENT OF MATHEMATICS, RICE UNIVERSITY, HOUSTON, TEXAS 77251-1892
E-mail address: `berkner@cml.rice.edu`