

**THE VOLUME OF A CLOSED HYPERBOLIC 3-MANIFOLD
IS BOUNDED BY π TIMES THE LENGTH OF ANY
PRESENTATION OF ITS FUNDAMENTAL GROUP**

DARYL COOPER

(Communicated by Ronald A. Fintushel)

Theorem 0.1. *Suppose M is a closed hyperbolic 3-manifold. Given a presentation of $\pi_1 M$ let L be the sum of the word-lengths of the relations and n the number of relations of length at least 3. Then $\text{volume}(M) < \pi(L - 2n)$.*

Proof. A **pleated disc** is a map covered by a map of a disc into \mathbb{H}^3 such that there is a triangulation of the disc with vertices only on the boundary of the disc and with the property that the image of each 2-simplex is a geodesic 2-simplex in \mathbb{H}^3 . A presentation of M gives a set of generators and relations. For simplicity, we will assume every relation has word length at least 3. This may be realized geometrically by a map $f : S \rightarrow M$ of a 2-complex S which induces an isomorphism of $\pi_1 S$ onto $\pi_1 M$. The map f may be homotoped so that edges map to geodesics and f restricted to each 2-cell is a pleated disc. The area of a pleated disc is at most π times the number of 2-simplices. The boundary of a 2-cell, D , in S represents a relation, and the number of 2-simplices in D is the number of edges in ∂D minus 2. The number of edges in ∂D is the word length of the relation represented by ∂D . Thus the total surface area of $f(S)$ is at most $\pi(L - 2n)$.

Let X be the closure of a component of $M - f(S)$; then X lifts to \mathbb{H}^3 . For otherwise, there is a loop γ in X which is not contractible in M . Since S is mapped into $M - \gamma$, the isomorphism $f_* : \pi_1 S \rightarrow \pi_1 M$ factors through $\pi_1(M - \gamma)$. Thus the composite

$$\pi_1 M \cong \pi_1 S \rightarrow \pi_1(M - \gamma) \rightarrow \pi_1 M$$

is the identity, where the second map is induced by inclusion. Now M is aspherical, hence $\pi_2(M - \gamma) = 0$ because otherwise, by the sphere theorem, γ would be contained inside a ball and thus contractible in M . Hence $M - \gamma$ is a $K(\pi, 1)$ and thus the first homomorphism is induced by a continuous map $M \rightarrow (M - \gamma)$. Thus the composite

$$M \rightarrow (M - \gamma) \rightarrow M$$

is a π_1 -isomorphism, hence a homotopy equivalence. Consideration of the induced map on H_3 gives a contradiction:

$$H_3(M) \rightarrow H_3(M - \gamma) \rightarrow H_3(M)$$

since the composite is an isomorphism and M is closed.

Received by the editors August 3, 1998.

1991 *Mathematics Subject Classification.* Primary 30F40, 57M50.

The author's research was supported in part by the NSF.

The isoperimetric inequality (for example [1], p. 283) for \mathbb{H}^3 states that, for a given volume, the smallest ratio of surface area divided by volume is attained by a sphere. Computation shows that this ratio is always greater than 2. This asymptotic ratio is attained by a horosphere. Thus the surface area of the polyhedron X in \mathbb{H}^3 is at least 2 times its volume. Now S may be subdivided so that each 2-cell appears in exactly two such polyhedra, thus the total surface area of S is greater than 1 times the volume of M . Putting this together with the first part gives the result. \square

In his thesis, Matt White [2] has obtained (a much deeper result) an explicit bound on the **diameter** of M in terms of the sum of the lengths of the relations. He also extends these results to the finite volume case.

REFERENCES

- [1] I. Chavel. *Riemannian Geometry: A Modern Introduction*. Cambridge University Press (1993). MR **95j**:53001
- [2] M. White. UCSB Thesis, to appear.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT SANTA BARBARA, SANTA BARBARA, CALIFORNIA 93106-0001
E-mail address: `cooper@math.ucsb.edu`