

## UNIVERSAL $\mathbb{Z}$ -LATTICES OF MINIMAL RANK

BYEONG-KWEON OH

(Communicated by David E. Rohrlich)

**ABSTRACT.** Let  $U_{\mathbb{Z}}(n)$  be the minimal rank of  $n$ -universal  $\mathbb{Z}$ -lattices, by which we mean positive definite  $\mathbb{Z}$ -lattices which represent all positive  $\mathbb{Z}$ -lattices of rank  $n$ . It is a well known fact that  $U_{\mathbb{Z}}(n) = n + 3$  for  $1 \leq n \leq 5$ . In this paper, we determine  $U_{\mathbb{Z}}(n)$  and find all  $n$ -universal lattices of rank  $U_{\mathbb{Z}}(n)$  for  $6 \leq n \leq 8$ .

### 1. INTRODUCTION

A positive definite  $\mathbb{Z}$ -lattice (or simply a lattice) is said to be *n-universal* if it represents all positive definite  $\mathbb{Z}$ -lattices of rank  $n$ . It is well known that the ranks of  $n$ -universal lattices should be greater than or equal to  $n + 3$ . In fact, for each  $n$ ,  $1 \leq n \leq 5$ , the lattice  $I_{n+3}$  is  $n$ -universal because  $I_{n+3}$  has class number 1 and is universal over the  $p$ -adic integer ring  $\mathbb{Z}_p$  for all  $p$ , where  $I_n$  is the lattice  $\mathbb{Z}^n$  equipped with the standard inner product (see [10], [12] and [15]). For  $n \geq 6$ , however, no diagonal lattice can be  $n$ -universal. Moreover, there does not exist a lattice of rank  $n + 3$  which has class number 1 and represents all integral lattices of rank  $n$  over  $\mathbb{Z}_p$  for all  $p$  (see [18], [20]). To be more precise, we define

$$U_{\mathbb{Z}}(n) = \min \{ \text{rank}(L) \mid L \text{ is } n\text{-universal} \}.$$

Let  $L_1, L_2, \dots, L_k$  be all unimodular lattices of rank  $n + 3$  up to isometry. Then the lattice  $L_1 \perp L_2 \perp \dots \perp L_k$  is  $n$ -universal and therefore  $U_{\mathbb{Z}}(n)$  exist for all  $n$ . As was mentioned above,  $U_{\mathbb{Z}}(n) = n + 3$  for  $1 \leq n \leq 5$ . In this paper, we investigate the minimal rank  $U_{\mathbb{Z}}(n)$  of  $n$ -universal  $\mathbb{Z}$ -lattices for  $6 \leq n \leq 10$ . We prove that  $U_{\mathbb{Z}}(n) = 13, 15, 16, 28, 30$  for  $n = 6, 7, 8, 9, 10$  respectively, and find all 6, 7, 8-universal  $\mathbb{Z}$ -lattices of rank 13, 15, 16, respectively. For the complete list of 1, 2-universal  $\mathbb{Z}$ -lattices of minimal rank, see [6], [7], [16] and [21].

In [1], Bannai proved that most unimodular lattices (even or odd) have trivial automorphism groups if the rank is sufficiently large, and that such lattices are indecomposable. If a lattice  $L$  is  $n$ -universal, then  $L$  must represent all indecomposable unimodular lattices of rank  $n$  as direct summands. So from this we may guess that  $U_{\mathbb{Z}}(n)$  grows very quickly.

*Remark.* Note that if we define  $U_{\mathbb{Q}}(n)$  to be the minimal rank of  $n$ -universal positive definite quadratic space over  $\mathbb{Q}$ , then  $U_{\mathbb{Q}}(n) = n + 3$  for all  $n$ .

---

Received by the editors April 27, 1998.

1991 *Mathematics Subject Classification.* Primary 11E12, 11H06.

*Key words and phrases.*  $n$ -universal lattice,  $U_{\mathbb{Z}}(n)$ , root lattice, additively indecomposable.

The author was partially supported by GARC and BSRI-98-1414.

We adopt terminologies and notations from [2], [3] and [14]. By  $l \rightarrow L$  we mean that the lattice  $L$  represents the lattice  $l$ . For a sublattice  $l$  of  $L \perp M$  of the form  $l = \mathbb{Z}(x_1 + y_1) + \mathbb{Z}(x_2 + y_2) + \cdots + \mathbb{Z}(x_n + y_n)$  for  $x_i \in L$  and  $y_i \in M$ , we define sublattices  $l(L) = \mathbb{Z}x_1 + \mathbb{Z}x_2 + \cdots + \mathbb{Z}x_n$  and  $l(M) = \mathbb{Z}y_1 + \mathbb{Z}y_2 + \cdots + \mathbb{Z}y_n$ . A lattice  $l$  is said to be *additively indecomposable* if either  $l(L) = 0$  or  $l(M) = 0$  whenever  $l \rightarrow L \perp M$ .

2. DETERMINATION OF  $U_{\mathbb{Z}}(6)$

We assume that  $L$  is a 6-universal  $\mathbb{Z}$ -lattice. Since  $L$  must represent the lattice  $I_6$ , it decomposes into  $I_6 \perp L'$ . Furthermore, since the root lattice  $E_6$  is additively indecomposable, it should be represented by  $L'$ . Therefore  $U_{\mathbb{Z}}(n) \geq 12$ . Suppose  $\text{rank } L = 12$ ; then  $L = I_6 \perp E_6$ . But this cannot represent the root lattice  $A_6$ . On the other hand, the lattice  $E_6 \perp I_{10}$  is 6-universal because  $E_6$  is the unique additively indecomposable lattice of rank 6 and  $E_6^{(2)}$  is represented by  $I_8$ , where  $E_6^{(2)}$  is the lattice obtained from scaling  $E_6$  by 2 (see [8] and [11]). Therefore  $13 \leq U_{\mathbb{Z}}(n) \leq 16$ . If  $L$  is 6-universal and  $\text{rank } L = 13$ , then  $L$  must be equal to  $E_6 \perp I_7$  or  $E_7 \perp I_6$  because the only lattice of rank 7 which represents both  $A_6$  and  $E_6$  is  $E_7$ . In this section, we prove that  $E_6 \perp I_7$  and  $E_7 \perp I_6$  are indeed 6-universal lattices of rank 13.

**Lemma 2.1.** *If a lattice  $L$  of rank  $n + 3$  has a square free determinant and its quadratic norm  $Q(L)$  is not contained in  $2\mathbb{Z}$ , then every lattice  $l$  of rank  $n$  is represented by a lattice in the genus of  $L$ .*

*Proof.* The local lattice  $L_p$  is  $n$ -universal over  $\mathbb{Z}_p$  by [13]. So the lemma follows directly from [14, 102:5]. (See also [4].) □

Now, we prove the following technical lemma, which is useful in the sequel.

**Lemma 2.2.** *Let  $l$  be a lattice of rank  $n$  which is represented by  $I_m$ ,  $m \geq 7$ .*

(1) *If  $5 \leq n \leq m - 2$ , then  $l$  is represented by  $D_{n-i} \perp I_{m-n+i}$  for some  $i = 1, 2, \dots, n - 1$ , where  $D_k$  is the root lattice of type  $D$  for  $k \geq 4$ ,  $D_3 = A_3$ ,  $D_2 = A_1 \perp A_1$ , and  $D_1 = \langle 4 \rangle$ .*

(2) *If  $n = m - 1$ , then  $l$  is represented by  $D_{n+1-i} \perp I_i$  for some  $i = 0, 1, 2, \dots, n$ .*

*Furthermore, if  $l$  is represented by  $D_{n+1-i} \perp I_i$  only for  $i = 0$  or  $1$  and  $n \equiv i \pmod{2}$ , then  $dl \equiv n - i + 1 \pmod{4}$ .*

*Proof.* We only prove (1). The proof of (2) is quite similar to that of (1). It suffices to show this when  $m$  is equal to  $n + 2$ . We may assume that  $l = \bigoplus_{i=1}^n \mathbb{Z}(\sum_{k=1}^m a_{ik} e_k)$  is a sublattice of  $I_m$ , where the  $e_i$ 's are the standard orthonormal basis of  $I_m$ . By suitable base change, we may also assume that  $a_{ij} = 0$  for all  $i, j$  satisfying  $i \geq 2$  and  $j \geq m + 2 - i$ , and that the  $a_{k(m+1-i)}$ 's are even for  $1 \leq k \leq i - 1$  if  $a_{i(m+1-i)}$  is odd for some  $i \geq 2$ . For a subset  $J = \{j_1, j_2, \dots, j_r\} \subset \{1, 2, \dots, m\}$ , we define the lattice  $M_J = \bigoplus_{i=1}^n \mathbb{Z}(\sum_{t=1}^r a_{ijt} e_{j_t})$ . We let  $J = \{m\}$  if  $a_{1m}$  is even. Then  $M_J = \mathbb{Z}(a_{1m} e_m) \rightarrow D_1$ . Assume that  $a_{i(m+1-i)}$  is even for some  $i$ ,  $2 \leq i \leq n - 1$ . Let  $i$  be the smallest such. Let  $J$  be the set containing  $(m + 1 - i)$  and all  $(m - k + 1)$ 's,  $1 \leq k \leq i - 1$ , for which the  $a_{k(m+1-i)}$ 's are odd. Then  $M_J \rightarrow D_{|J|}$  and hence  $l \rightarrow D_{|J|} \perp I_{m-|J|}$ . Therefore we may assume that the  $a_{i(m+1-i)}$ 's are odd for all  $i$ ,  $1 \leq i \leq n - 1$ .

Now assume that  $a_{nj}$  is even for some  $j$ ,  $1 \leq j \leq 3$ . If not all  $a_{kj}$ 's are odd for  $1 \leq k \leq n - 1$ , then  $M_J \rightarrow D_{|J|}$  as above. Hence if one of  $a_{nj}$  is even for  $j = 1, 2, 3$ ,

then we may assume that the  $a_{kj}$ 's are all odd for  $k = 1, 2, \dots, n - 1$ . If two of the  $a_{nj}$ 's are even, then  $l \rightarrow D_2 \perp I_{m-2}$ . Therefore, without loss of generality, we may assume that  $a_{n3}$  is odd and  $a_{k3}$  are all even for  $k = 1, 2, \dots, n - 1$ . For a fixed  $s$ ,  $s = 1$  or  $2$ , if the number of  $a_{ks}$ 's which are odd is less than  $n - 1$  for  $k = 1, 2, \dots, n$ , then  $l \rightarrow D_{|J|} \perp I_{m-|J|}$ , where  $J$  is the set containing  $s$  and the  $(m - k + 1)$ 's for which the  $a_{ks}$ 's are odd. In the remaining case, it is easy to see that  $l \rightarrow D_k \perp I_{m-k}$ , where  $k$  is 2 or 3 or 4.  $\square$

**Theorem 2.3.** *The lattice  $E_7 \perp I_6$  is 6-universal. In particular,  $U_{\mathbb{Z}}(n) = 13$ .*

*Proof.* First observe that  $gen(I_8 \perp A_1) = \{I_8 \perp A_1, E_7 \perp I_2\}$ . Hence it suffices to show that every sublattice  $l$  of  $A_1 \perp I_8$  of rank 6 is represented by  $E_7 \perp I_6$  by Lemma 2.1. By Lemma 2.2 (1),  $l(I_8) \rightarrow D_{5-i} \perp I_{3+i}$ , for some  $i = 0, 1, \dots, 4$ . If  $i \neq 4$ , then we have

$$l \rightarrow A_1 \perp I_8 \rightarrow A_1 \perp D_{5-i} \perp I_{i+3} \rightarrow E_7 \perp I_6.$$

If  $i = 4$ , then  $l' = l(I_8) \rightarrow D_1 \perp I_7$ . We apply Lemma 2.2 (2) to  $l'(I_7)$ . By similar reasoning as above, we need only consider the case when  $l'(I_7) \rightarrow D_7$ . This indeed implies  $l \rightarrow A_1 \perp D_1 \perp D_7$  and  $d(l(D_7)) \equiv 7 \pmod{8}$ . For all prime  $p$  (including  $\infty$ ), since  $l(D_7)_p$  is represented by  $(E_7 \perp A_1)_p$  and the class number of  $E_7 \perp A_1$  is 1 [19], we have  $l(D_7) \rightarrow E_7 \perp A_1$ , which proves the theorem.  $\square$

In order to prove that  $E_6 \perp I_7$  is the other 6-universal lattice of rank 13, we need the following lemma.

**Lemma 2.4.** *If a  $\mathbb{Z}$ -lattice  $l$  of rank 6 is not represented by a sum of squares, then  $l \rightarrow E_6 \perp I_5$ .*

*Proof.* We may assume that  $l \rightarrow E_7 \perp I_2$ . By [8], we may also assume that  $d(l(E_7))$  is an odd determinant. Since the class number of  $E_6 \perp A_2$  is 1, it can easily be checked that  $l(E_7) \rightarrow E_6 \perp A_2$ , and hence  $l \rightarrow E_6 \perp I_5$  if  $d(l(E_7)) \not\equiv 1 \pmod{3}$ . So we assume that  $d(l(E_7)) \equiv 1 \pmod{6}$ . By considering local conditions for representation, we can conclude that  $l(E_7) \rightarrow gen(E_6 \perp I_2)$  and consequently  $l \rightarrow E_6 \perp I_5$  from the fact that  $gen(E_6 \perp I_2) = \{E_6 \perp I_2, \langle 3 \rangle \perp I_7\}$ .  $\square$

*Remark.* Ko conjectured [11] that if  $l$  is of rank 6 and represented by a sum of squares, then  $l \rightarrow I_9$ , and if  $l$  is of rank 6 and not represented by a sum of squares, then  $l \rightarrow E_6 \perp I_3$  and  $l(E_6) = E_6$ . But both conjectures are false because  $l = A_2 \perp A_2 \perp A_1 10[1\frac{1}{2}]$  is represented by  $I_{10}$  but not by  $I_9$  for the former conjecture (see [8], [9] for further results) and  $l = D_5 124[1\frac{1}{4}]$ , which is not represented by a sum of squares, is represented by  $E_6 \perp I_3$  but does not satisfy  $l(E_6) = E_6$ .

**Theorem 2.5.** *The lattice  $E_6 \perp I_7$  is 6-universal.*

*Proof.* Let  $l$  be a  $\mathbb{Z}$ -lattice of rank 6. By the above lemma, we may assume that  $l$  is represented by a sum of squares, and hence by [8] we may assume that  $l \rightarrow I_{10}$ . This implies that  $l \rightarrow D_{5-i} \perp I_{5+i}$  for some  $i = 0, 1, \dots, 4$  by Lemma 2.2 (1). If  $i \neq 3, 4$ , then  $l \rightarrow D_{5-i} \perp I_{5+i} \rightarrow E_6 \perp I_7$ . The desired conclusion for the case when  $i = 3, 4$  can be deduced by applying Lemma 2.2 again if necessary.  $\square$

3. DETERMINATION OF  $U_{\mathbb{Z}}(n)$  FOR  $7 \leq n \leq 10$

**Theorem 3.1.** *The lattice  $E_8 \perp I_8$  is a unique 8-universal  $\mathbb{Z}$ -lattice of rank 16, and  $U_{\mathbb{Z}}(8) = 16$ .*

*Proof.* Note that the lattice  $E_8 \perp I_8$  is the unique candidate of 8-universal  $\mathbb{Z}$ -lattice of rank 16, for  $E_8$  is the unique additively indecomposable  $\mathbb{Z}$ -lattice of rank 8. Let  $l$  be a  $\mathbb{Z}$ -lattice of rank 8. Since  $l \rightarrow \text{gen}(E_8 \perp I_3) = \{E_8 \perp I_3, I_{11}\}$ , we may assume that  $l \rightarrow I_{11}$ . By Lemma 2.2, we may further assume that  $l \rightarrow A_1 \perp A_1 \perp D_9$  and  $d(l(D_9)) \equiv 1 \pmod{8}$ . Clearly,  $l(D_9)$  is contained in one of the sublattices of  $I_9$  of rank 9 with determinant 9. The following are all such sublattices of  $I_9$ :

$$\begin{aligned} &\langle 9 \rangle \perp I_8, A_1 18[1\frac{1}{2}] \perp I_7, A_2 \perp \langle 3 \rangle \perp I_6, A_3 36[1\frac{1}{4}] \perp I_5, A_4 45[2\frac{1}{5}] \perp I_4, \\ &A_5 6[3\frac{1}{2}] \perp I_3, A_6 63[3\frac{1}{7}] \perp I_2, A_7 72[3\frac{1}{8}] \perp I_1, \text{ and } A_8 9[3\frac{1}{3}]. \end{aligned}$$

One can easily check that if  $l(D_9)$  is represented by one of these lattices except the first one, then  $l \rightarrow E_8 \perp I_8$ . So assume that  $l(D_9) \rightarrow \langle 9 \rangle \perp I_8$ . Then  $l(D_9)$  is represented by  $\mathbb{Z}(e_1 - e_2) + \mathbb{Z}(e_2 - e_3) + \dots + \mathbb{Z}(e_7 - e_8) + \mathbb{Z}(e_8 - 3e_9) + \mathbb{Z}(e_8 + 3e_9)$  and hence is represented by  $A_8 \perp I_5$ . Therefore  $l$  is represented by  $E_8 \perp I_8$ .  $\square$

*Remark.* In [5], Conway and Schneeberger proved the so-called 15-Theorem, i.e., every integral  $\mathbb{Z}$ -lattice which represents 1, 2, 3, 5, 6, 7, 10, 14, 15 is 1-universal. An analogy for 8-universal  $\mathbb{Z}$ -lattices can be deduced from Theorem 3.1: Every  $\mathbb{Z}$ -lattice which represents both  $I_8$  and  $E_8$  is 8-universal.

**Corollary 3.2.** *The lattice  $E_8 \perp I_7$  is 7-universal and  $U_{\mathbb{Z}}(7) = 15$ .*

*Proof.* The 7-universality of  $E_8 \perp I_7$  follows from the above theorem. Consider the only possible candidate for a 7-universal  $\mathbb{Z}$ -lattice of rank 14; namely,  $E_7 \perp I_7$ . But this cannot represent  $A_6 77[2\frac{1}{7}]$ , and the result follows.  $\square$

**Theorem 3.3.** *There are exactly three 7-universal  $\mathbb{Z}$ -lattices of rank 15. They are  $E_8 \perp I_7$ ,  $E_7 \perp I_8$ , and  $E_7 6[1\frac{1}{2}] \perp I_7$ .*

*Proof.* Suppose that  $L$  is a 7-universal  $\mathbb{Z}$ -lattice of rank 15. Then  $L = I_7 \perp L'$  and  $\text{rank}(L') = 8$ . Clearly,  $E_7 \rightarrow L'$ . If the lattice  $L'$  represents 1, then  $L = I_8 \perp E_7$ . So assume that  $L'$  does not represent 1. Since  $A_6 77[2\frac{1}{7}] \rightarrow L$ , either  $D_7 \rightarrow L'$  or  $A_6 77[2\frac{1}{7}] \rightarrow L'$ . In the first case,  $L'$  must be  $E_8$ , for  $E_8$  is the only lattice of rank 8 which represents  $E_7$  and  $D_7$  simultaneously. In the second case, since the minimum quadratic norm of the dual lattice  $E_7^\#$  of  $E_7$  is  $\frac{3}{2}$ , it can be easily deduced that  $L'$  must be  $E_7 6[1\frac{1}{2}]$ . Hence we have exactly three candidates  $E_8 \perp I_7$ ,  $E_7 \perp I_8$  and  $E_7 6[1\frac{1}{2}] \perp I_7$  for 7-universal  $\mathbb{Z}$ -lattices of minimal rank, 15.

It suffices to show the 7-universality for the latter two. First, we show that  $E_7 \perp I_8$  is 7-universal. Let  $l$  be any  $\mathbb{Z}$ -lattice of rank 7. Note that

$$l \rightarrow \text{gen}(E_8 \perp I_2) = \{E_8 \perp I_2, I_{10}\}.$$

If  $l \rightarrow I_{10}$ , it is easy to check that  $l \rightarrow E_7 \perp I_8$  by Lemma 2.2(1). So assume that  $l \rightarrow E_8 \perp I_2$ . Note that  $l(E_8)$  can be represented by one of the sublattices of  $E_8$  with determinant 4; the only such sublattices are  $E_7 \perp A_1$  and  $D_8$ . Therefore the 7-universality of  $E_7 \perp I_8$  follows immediately.

Now we prove that  $E_76[1\frac{1}{2}] \perp I_7$  is 7-universal. Note that for every  $\mathbb{Z}$ -lattice  $l$  of rank 7

$$l \rightarrow \text{gen}(E_76[1\frac{1}{2}] \perp I_2) = \{E_76[1\frac{1}{2}] \perp I_2, A_2 \perp I_8\}.$$

So we assume that  $l \rightarrow A_2 \perp I_8$ . Then  $l(I_8)$  is contained in one such sublattice of  $I_8$  of rank 8 with determinant 9. It is easy to check that  $l \rightarrow E_76[1\frac{1}{2}] \perp I_7$  if  $l(I_8)$  is contained in one such sublattice except  $A_772[3\frac{1}{8}]$ . Therefore, we may restrict ourselves to the case when

$$l(I_8) \rightarrow A_772[3\frac{1}{8}] = \{ \sum_{i=1}^8 a_i e_i \mid \sum_{i=1}^8 a_i \equiv 0 \pmod{3} \}.$$

Furthermore, we may assume that  $d(l(I_8)) \equiv 2 \pmod{3}$ , for we may assume that  $l(I_8)$  is not contained in any sublattice of  $I_8$  of rank 8 with determinant 9 other than  $A_772[3\frac{1}{8}]$ . By Lemma 2.2, we obtain  $l \rightarrow D_{8-i} \perp I_i \perp A_2$  for  $i = 0, 1, \dots, 7$ . If  $i \neq 0, 1$ , then this implies  $l \rightarrow E_76[1\frac{1}{2}] \perp I_7$ , as desired. If  $i = 0$ , then

$$l(I_8) \rightarrow A_772[2\frac{1}{4}] = \{ \sum_{i=1}^8 a_i e_i \mid \sum_{i=1}^8 a_i \equiv 0 \pmod{6} \} \rightarrow E_76[1\frac{1}{2}] \perp I_3$$

and hence  $l \rightarrow E_76[1\frac{1}{2}] \perp I_7$ . If  $i = 1$ , then we may assume that  $d(l(I_8)) \equiv 11 \pmod{12}$  by Lemma 2.2(2). Therefore

$$l(I_8) \rightarrow \text{gen}(A_2 \perp I_7) = \{A_2 \perp I_7, E_76[1\frac{1}{2}] \perp I_1\}.$$

Consequently,  $l \rightarrow E_76[1\frac{1}{2}] \perp I_7$  as desired.  $\square$

**Theorem 3.4.** *The lattice  $E_8 \perp I_9 \perp D_{10}A_1[11]$  is a 9-universal  $\mathbb{Z}$ -lattice and  $U_{\mathbb{Z}}(9) = 28$ .*

*Proof.* Suppose that  $L$  is a 9-universal  $\mathbb{Z}$ -lattice. Then  $L$  must decompose into  $E_8 \perp I_9 \perp L'$ . There exist exactly two additively indecomposable  $\mathbb{Z}$ -lattices of rank 9, namely,  $A_863[4\frac{1}{9}]$  and  $A_4A_415[33\frac{1}{5}]$  (see [17]). Since  $L'$  must represent these lattices, the rank of  $L'$  is greater than 9. Suppose that the rank of  $L'$  is 10. Then  $A_9 \rightarrow L'$ , since  $1 \notin Q(L')$ . Furthermore,  $L'$  has a vector of norm 3, since  $A_863[4\frac{1}{9}] \rightarrow L'$ . The possible candidates for  $L'$  satisfying these properties are the following:

$$A_9210[1\frac{1}{10}], A_935[2\frac{1}{5}], A_990[3\frac{1}{10}], A_915[4\frac{1}{5}], A_9A_1[5\frac{1}{2}], \text{ and } A_9 \perp \langle 3 \rangle.$$

Among these lattices, only  $A_915[4\frac{1}{5}]$  and  $A_9A_1[5\frac{1}{2}]$  can represent  $A_863[4\frac{1}{9}]$  and  $A_4A_415[33\frac{1}{5}]$  simultaneously. But neither  $E_8 \perp I_9 \perp A_915[4\frac{1}{5}]$  nor  $E_8 \perp I_9 \perp A_9A_1[5\frac{1}{2}]$  can represent  $A_8117[2\frac{1}{9}]$ . Therefore the rank of  $L'$  is greater than 10. On the other hand, it can be easily checked by using Lemma 2.2 that every  $\mathbb{Z}$ -lattice  $l$  of rank 9, which is represented by  $A_1 \perp I_{11}$ , is represented by  $E_8 \perp D_{10}A_1[11] \perp I_9$ . Hence from

$$\text{gen}(A_1 \perp I_{11}) = \{A_1 \perp I_{11}, E_7 \perp I_5, D_{10}A_1[11] \perp I_1, E_8 \perp I_3 \perp A_1\}$$

we may conclude that  $E_8 \perp D_{10}A_1[11] \perp I_9$  is 9-universal, and the result follows.  $\square$

**Theorem 3.5.** *The lattice  $E_8 \perp I_{10} \perp D_{12}[1]$  is 10-universal and  $U_{\mathbb{Z}}(10) = 30$ .*

*Proof.* Suppose that  $L$  is a 10-universal  $\mathbb{Z}$ -lattice. The lattice  $L$  must decompose into  $E_8 \perp I_{10} \perp L'$ . The lattices  $D_9 12[1\frac{1}{4}]$ ,  $A_9 A_1[51]$  are additively indecomposable  $\mathbb{Z}$ -lattices of rank 10 (see [17]), so  $L'$  must represent these lattices. Suppose that the rank of  $L'$  is 11; then  $A_{10} \rightarrow L'$ , since  $1 \notin Q(L')$ . But there does not exist a lattice of rank 11 which represents the lattices  $A_{10}$  and  $D_9 12[1\frac{1}{4}]$  simultaneously. Hence the rank of  $L'$  is greater than 11. On the other hand, since

$$\text{gen}(I_{13}) = \{I_{13}, E_8 \perp I_5, D_{12}[1] \perp I_1\},$$

it can be easily checked by applying Lemma 2.2 that  $E_8 \perp I_{10} \perp D_{12}[1]$  is 10-universal. Therefore the result follows.  $\square$

*Remark.* It seems to be a very difficult problem to find the exact value of  $U_{\mathbb{Z}}(n)$  for large  $n$ . For example, one can easily obtain  $U_{\mathbb{Z}}(24) \geq 6673$  from a simple counting of all indecomposable unimodular  $\mathbb{Z}$ -lattices of rank less than or equal to 24 (see [2]).

#### ACKNOWLEDGEMENTS

This article is a part of the author's doctoral thesis. I wish to thank my doctoral supervisor M-H. Kim for continuous encouragement.

#### REFERENCES

- [1] E. Bannai, *Positive definite unimodular lattices with trivial automorphism groups*, Ohio State Univ., Thesis, 1988.
- [2] J.H. Conway, N.J.A. Sloane, *Sphere packings, lattices and groups*, Springer-Verlag, 1988. MR **89a**:11067
- [3] ———, *Low dimensional lattices. I. Quadratic forms of small determinant*, Proc. Royal Soc. Lond. A. **418** (1988), 17-41. MR **90a**:11071
- [4] ———, *Low dimensional lattices. V. Integral coordinates for integral lattices*, Proc. Royal Soc. Lond. A **426** (1989), 211-232. MR **90m**:11100
- [5] J.H. Conway, W. Schneeberger, *A 15-theorem for universal quadratic forms*, to appear.
- [6] B.M. Kim, M-H. Kim, S. Raghavan, *2-universal positive definite integral quinary diagonal quadratic forms*, Ramanujan J. **1** (1997), 333-337. CMP 98:10
- [7] B.M. Kim, M-H. Kim, B-K. Oh, *2-universal positive definite integral quinary quadratic forms*, Preprint.
- [8] M-H. Kim, B-K. Oh, *Representations of positive definite senary integral-quadratic forms by a sum of squares*, J. Number Theory **63** (1997), 89-100. MR **98a**:11045
- [9] ———, *A lower bound for the number of squares whose sum represents integral quadratic forms*, J. Korean Math. Soc. **33** (1996), 651-655. MR **97j**:11017
- [10] C. Ko, *On the representation of a sum of squares of linear forms*, Quart. J. Math. Oxford **8** (1937), 81-98.
- [11] ———, *On the decomposition of quadratic forms in six variables*, Acta Arith. **3** (1939), 64-78.
- [12] J.L. Lagrange, *Oeuvres* **3** (1869), 189-201.
- [13] O.T. O'Meara, *The integral representations of quadratic forms over local fields*, Amer. J. Math. **80** (1958), 843-878. MR **20**:4526
- [14] ———, *Introduction to quadratic forms*, Springer-Verlag, 1973.
- [15] L.J. Mordell, *A new Waring's problem with squares of linear forms*, Quart. J. Math. Oxford **1** (1930), 276-288.
- [16] S. Ramanujan, *On the expression of a number in the form  $ax^2 + by^2 + cz^2 + dw^2$* , Proc Cambridge Phil. Soc. **19** (1917), 11-21.
- [17] W. Plesken, *Additively indecomposable positive integral quadratic forms*, J. Number Theory **47** (1994), 273-283. MR **95c**:11045
- [18] G.L. Watson, *The class number of a positive quadratic form*, Proc. London Math. Soc. (3) **13** (1963), 549-576. MR **27**:107

- [19] ———, *One-class genera of positive quadratic forms in at least five variables*, *Acta Arith.* **26** (1975), 309-327. MR **52**:274
- [20] ———, *One-class genera of positive quadratic forms in nine and ten variables*, *Math.* **25** (1978), 57-67. MR **58**:10738
- [21] M.F. Willerding, *Determination of all classes of (positive) quaternary quadratic forms which represent all positive integers*, *Bull. Amer. Math. Soc.* **54** (1948), 334-337. MR **9**:571e

DEPARTMENT OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SEOUL, 151-742, KOREA  
E-mail address: oandhan@math.snu.ac.kr