

ON ABSORBING EXTENSIONS

KLAUS THOMSEN

(Communicated by David R. Larson)

ABSTRACT. Building on the work of Kasparov we show that there always exists a trivial absorbing extension of A by $B \otimes \mathcal{K}$, provided only that A and B are separable. If A is unital there is a unital trivial extension which is unitaly absorbing.

1. INTRODUCTION

Absorbing trivial extensions play an important role in the theory of extensions of C^* -algebras; cf. 15.12 in [Bl]. Recently the interest in such extensions has been renewed because of the way KK -theory comes into the classification program. In this connection, as well as in the proper theory of C^* -extensions, it is slightly disturbing that the existence of an absorbing trivial extension has only been established in the case where at least one of the C^* -algebras involved is nuclear; cf. Theorem 5 of [K]. The purpose of the present note is to show that such extensions always exist when both C^* -algebras are separable. The argument for this is a modification of Kasparov's approach from [K]. The absorbing trivial extensions were constructed, in [K] as well as before Kasparov's work, by taking the infinite direct sum of the same copy of a faithful unital representation of the separable C^* -algebra A (for the moment assumed to be unital) which plays the role of the quotient in the extensions. The resulting representation $\pi : A \rightarrow \mathcal{B}(\mathcal{H})$ was then composed with the natural imbedding $\mathcal{B}(\mathcal{H}) \subseteq \mathcal{M}(B \otimes \mathcal{K})$, where $B \otimes \mathcal{K}$ is the stable C^* -algebra which features as the ideal in the extensions. So in practice this means that the absorbing extension was constructed by taking a weak* dense sequence of states of A , repeating all states in the sequence infinitely often, and then adding the corresponding GNS-representations. This procedure has nothing to do with the C^* -algebra B , and it is a highly non-trivial task to show that it often results in an absorbing extension when prolonged to a map $A \rightarrow \mathcal{M}(B \otimes \mathcal{K})$; cf. [K]. The observation we offer here is that if one instead takes a sequence $s_n : A \rightarrow B \otimes \mathcal{K}$ of completely positive contractions which is dense for the topology of pointwise norm-convergence among all completely positive contractions (such a sequence exists when both A and B are separable), repeats each s_n infinitely often and add up the unital representations

$$\pi_n : A \rightarrow \mathcal{M}(B \otimes \mathcal{K}), \quad n \in \mathbb{N},$$

Received by the editors August 11, 1999.

2000 *Mathematics Subject Classification*. Primary 46L80, 19K35.

©2000 American Mathematical Society

coming from the Kasparov-Stinespring decompositions

$$s_n(\cdot) = W_n^* \pi_n(\cdot) W_n ,$$

the resulting representation $A \rightarrow \mathcal{M}(B \otimes \mathcal{K})$ will be a unitaly absorbing trivial extension. The general trivial absorbing extensions can then be obtained (for a not necessarily unital C^* -algebra A) by taking a unitaly absorbing representation $\pi : A^+ \rightarrow \mathcal{M}(B \otimes \mathcal{K})$ and restricting it to A .

In order to illustrate how the absorbing $*$ -homomorphisms constructed here can be used to extend known results we prove a general version of the Paschke-Valette-Skandalis duality which realizes the group $KK(A, B)$ as the K_1 -group of a C^* -algebra D_π built out of A and B by using an absorbing $*$ -homomorphism $\pi : A \rightarrow \mathcal{M}(B)$; cf. [P], [V], [S], [H].

2. ABSORBING $*$ -HOMOMORPHISMS

Given Hilbert B -modules E and F , we let $\mathcal{L}_B(E, F)$ denote the Banach space of adjointable operators from E to F . The ideal of ‘compact’ operators from E to F is denoted by $\mathcal{K}_B(E, F)$. When $E = F$ we write $\mathcal{L}_B(E)$ and $\mathcal{K}_B(E)$ instead of $\mathcal{L}_B(E, E)$ and $\mathcal{K}_B(E, E)$, respectively. In the special case where $E = B$ there are well-known identifications $\mathcal{L}_B(B) = \mathcal{M}(B) =$ the multiplier algebra of B and $\mathcal{K}_B(B) = B$, which we shall use freely.

Theorem 2.1. *Let A and B be separable C^* -algebras with A unital and B stable. Let $\pi : A \rightarrow \mathcal{M}(B)$ be a unital $*$ -homomorphism. Then the following conditions are equivalent:*

- 1) *For any completely positive contraction $\varphi : A \rightarrow B$ there is a sequence $\{W_n\} \subseteq \mathcal{M}(B)$ such that*
 - 1a) $\lim_{n \rightarrow \infty} \|\varphi(a) - W_n^* \pi(a) W_n\| = 0$ for all $a \in A$,
 - 1b) $\lim_{n \rightarrow \infty} \|W_n^* b\| = 0$ for all $b \in B$.
- 2) *For any completely positive unital map $\varphi : A \rightarrow \mathcal{M}(B)$ there is a sequence $\{V_n\}$ of isometries in $\mathcal{M}(B)$ such that*
 - 2a) $V_n^* \pi(a) V_n - \varphi(a) \in B$, $n \in \mathbb{N}$, $a \in A$,
 - 2b) $\lim_{n \rightarrow \infty} \|V_n^* \pi(a) V_n - \varphi(a)\| = 0$, $a \in A$.
- 3) *For any unital $*$ -homomorphism $\varphi : A \rightarrow \mathcal{M}(B)$ there is a sequence $\{U_n\}$ of unitaries $U_n \in \mathcal{L}_B(B \oplus B, B)$ such that*
 - 3a) $U_n \begin{pmatrix} \pi(a) & 0 \\ 0 & \varphi(a) \end{pmatrix} U_n^* - \pi(a) \in B$, $n \in \mathbb{N}$, $a \in A$,
 - 3b) $\lim_{n \rightarrow \infty} \|U_n \begin{pmatrix} \pi(a) & 0 \\ 0 & \varphi(a) \end{pmatrix} U_n^* - \pi(a)\| = 0$, $a \in A$.
- 4) *For any unital $*$ -homomorphism $\varphi : A \rightarrow \mathcal{M}(B)$ there is a sequence $\{U_n\}$ of unitaries $U_n \in \mathcal{L}_B(B \oplus B, B)$ such that*

$$\lim_{n \rightarrow \infty} \|U_n \begin{pmatrix} \pi(a) & 0 \\ 0 & \varphi(a) \end{pmatrix} U_n^* - \pi(a)\| = 0, a \in A .$$

Proof. 1) \Rightarrow 2): Let $F \subseteq A$ be a finite set containing 1 and $\epsilon > 0$. Let $\varphi : A \rightarrow \mathcal{M}(B)$ be a completely positive unital map. It suffices to find an element $V \in \mathcal{M}(B)$ such that

$$(2.1) \quad V^* \pi(a) V - \varphi(a) \in B$$

for all $a \in A$ and

$$(2.2) \quad \|V^* \pi(x)V - \varphi(x)\| < 3\epsilon$$

for all $x \in F$. If namely ϵ is small enough this will imply that $W = V[V^*V]^{-\frac{1}{2}}$ is an isometry close to V such that $V - W \in B$, and we can then work with W instead of V . We repeat Kasparov's arguments: Let X be a compact subset of A containing F and with dense span in A . By Lemma 10 of [K] there is a sequence $\psi_k : A \rightarrow B$, $k \in \mathbb{N}$, of completely positive contractions such that $\psi(a) = \sum_{k=1}^{\infty} \psi_k(a)$ converges in the strict topology, $\varphi(a) - \psi(a) \in B$ for all $a \in A$, and $\|\varphi(x) - \psi(x)\| < \epsilon$ for all $x \in X$. Let $\{b_i\}$ be a countable approximate unit for B . It follows from 1) that we can find a sequence $\{m_i\} \subseteq B$ such that

- 1) $\|\psi_i(x) - m_i^* \pi(x) m_i\| \leq \epsilon 2^{-i}$, $x \in X$, $i \in \mathbb{N}$,
- 2) $\|m_i^* \pi(x) m_j\| \leq \epsilon 2^{-i-j}$, $x \in X$, $i, j \in \mathbb{N}$, $i \neq j$,
- 3) $\sum_{i=1}^{\infty} \|m_i^* b_k\| < \infty$ for all $k \in \mathbb{N}$.

The argument from the proof of Theorem 5 in [K] shows that $\sum_{i=1}^{\infty} m_i$ converges in the strict topology to an element $V \in \mathcal{M}(B)$ satisfying (2.1) and (2.2).

2) \Rightarrow 3): This follows from the arguments of Arveson given on pp. 338-339 of [A] by substituting Hilbert spaces with Hilbert B -modules. We leave this to the reader.

3) \Rightarrow 4) is trivial.

4) \Rightarrow 1): Let $\varphi : A \rightarrow B$ be a completely positive contraction. Let $F \subseteq A$ and $G \subseteq B$ be finite sets and $\epsilon > 0$. Since A and B are separable it suffices to find an element $L \in \mathcal{M}(B)$ such that $\|\varphi(a) - L^* \pi(a)L\| < \epsilon$, $a \in F$, and $\|Lb\| < \epsilon$ for all $b \in B$. By Kasparov's Stinespring theorem (Theorem 3 of [K]), there is a unital $*$ -homomorphism $\chi : A \rightarrow \mathcal{M}(B)$ and an element $W \in \mathcal{M}(B)$ such that $\varphi(\cdot) = W^* \chi(\cdot) W$. Let S_i , $i = 1, 2, 3, \dots$, be a sequence of isometries in $\mathcal{M}(B)$ such that $S_i^* S_j = 0$, $i \neq j$, and $\sum_{i=1}^{\infty} S_i S_i^* = 1$ in the strict topology, and set $\chi^\infty(a) = \sum_{i=1}^{\infty} S_i \chi(a) S_i^*$. It follows from 4) that there is a sequence $\{U_n\}$ of unitaries in $\mathcal{L}_B(B \oplus B, B)$ such that

$$\lim_{n \rightarrow \infty} \|U_n \begin{pmatrix} \pi(a) & 0 \\ 0 & \chi^\infty(a) \end{pmatrix} U_n^* - \pi(a)\| = 0, \quad a \in A.$$

Define $T_i : B \rightarrow B \oplus B$ by $T_i b = (0, S_i b)$. Then $\chi(a) = T_i^* \begin{pmatrix} \pi(a) & 0 \\ 0 & \chi^\infty(a) \end{pmatrix} T_i$ and

$$\varphi(a) = W^* T_i^* \begin{pmatrix} \pi(a) & 0 \\ 0 & \chi^\infty(a) \end{pmatrix} T_i W \text{ for all } a \text{ and } i. \text{ Choose } n \text{ so large that}$$

$$\left\| \begin{pmatrix} \pi(a) & 0 \\ 0 & \chi^\infty(a) \end{pmatrix} - U_n^* \pi(a) U_n \right\| < \frac{\epsilon}{1 + \|W\|^2}, \quad a \in F.$$

Then

$$\|\varphi(a) - W^* T_i^* U_n^* \pi(a) U_n T_i W\| < \epsilon, \quad a \in F,$$

for all i . Since $\lim_{i \rightarrow \infty} \|T_i^* x\| = 0$ for all $x \in B \oplus B$, we can choose i so large that $\|W^* T_i^* U_n^* b\| < \epsilon$ for all $b \in G$. Set $L = U_n T_i W$. \square

Definition 2.2. Let A and B be separable C^* -algebras with A unital and B stable. A unital $*$ -homomorphism $\pi : A \rightarrow \mathcal{M}(B)$ which satisfies the four equivalent conditions in Theorem 2.1 is called *unittally absorbing* (for (A, B)).

The following lemma is surely known, but it is so crucial for us here that we include a proof.

Lemma 2.3. *Let A and B be separable C^* -algebras. There is then a countable set X of completely positive contractions $A \rightarrow B$ such that for any completely positive contraction $\mu : A \rightarrow B$, any finite set $F \subseteq A$ and any $\epsilon > 0$ there is an element $l \in X$ such that*

$$\|\mu(f) - l(f)\| \leq \epsilon, \quad f \in F.$$

Proof. Let $\{a_1, a_2, a_3, \dots\}$ be a dense sequence in the unit ball of A and set $F_n = \text{span}\{a_1, a_2, \dots, a_n\}$. Let ω be a faithful state of A and let (π_ω, H_ω) be the GNS-representation coming from ω . We can then consider A as a subspace of H_ω . The orthogonal projection $P_n : H_\omega \rightarrow F_n$ gives us then by restriction a continuous idempotent map $P_n : A \rightarrow F_n$. Let $1 < m_1 < m_2 < m_3 < \dots$ be a sequence of numbers such that $\|P_n\| \leq m_n$ for all n . We can then define a metric d on the space $\mathcal{B}(A, B)$ of continuous linear maps $L : A \rightarrow B$ by

$$d(L_1, L_2) = \sum_{i=1}^{\infty} \frac{2^{-i}}{m_i} \|L_1(a_i) - L_2(a_i)\|.$$

Choose a linear basis $\{x_1, x_2, \dots, x_{n_0}\}$ for F_n . For each n_0 -tuple $\underline{b} = (b_1, b_2, \dots, b_{n_0}) \in B^{n_0}$ there is a linear map $L_{\underline{b}} : F_n \rightarrow B$ such that $L_{\underline{b}}(x_i) = b_i$, $i = 1, 2, \dots, n_0$. By using that B^{n_0} is separable this construction gives us a countable set \mathcal{M} of linear maps $F_n \rightarrow B$ which is dense in the strong topology of $\mathcal{B}(F_n, B)$. Now let $0 < \epsilon < 1$ and let a finite set $D \subseteq F_n$ be given. Let $\mu \in \mathcal{B}(F_n, B)$ be a contraction. There is a finite subset G of F_n such that every $x \in F_n$ with $\|x\| \leq 1 - \epsilon$ is a convex combination of elements from G . Choose $l \in \mathcal{M}$ such that

$$(2.3) \quad \|\mu(z) - l(z)\| < \epsilon, \quad z \in D \cup G.$$

Then $\|\mu(x) - l(x)\| \leq \epsilon$ for all $x \in F_n$ with $\|x\| \leq 1 - \epsilon$, and hence $\|l\| \leq \frac{1+\epsilon}{1-\epsilon}$. Let q be a positive rational number in $]\frac{1-2\epsilon}{1+\epsilon}, \frac{1-\epsilon}{1+\epsilon}[$. Then $ql \in \mathbb{Q}_+ \mathcal{M}$ is a contraction and we find that

$$\begin{aligned} \|\mu(z) - ql(z)\| &\leq \|\mu(z) - l(z)\| + \|l(z) - ql(z)\| \\ &\leq \epsilon + |1 - q| \|l\| \sup\{\|z\| : z \in D\} \\ &< \frac{2\epsilon + 2\epsilon^2}{1 - \epsilon^2} \sup\{\|z\| : z \in D\} + \epsilon \end{aligned}$$

for all $z \in D$. It follows that we can find a countable set $\mathcal{Y}_n \subseteq \mathbb{Q}_+ \mathcal{M}$ of linear contractions which is strongly dense among all contractions in $\mathcal{B}(F_n, B)$. Set

$$\mathcal{Y} = \bigcup_{n=1}^{\infty} \{l \circ P_n : l \in \mathcal{Y}_n\}.$$

Let $\mu : A \rightarrow B$ be a linear contraction and let $\epsilon > 0$. Choose n so large that $2 \sum_{i \geq n+1} 2^{-i} < \frac{\epsilon}{2}$. From what we have just proved there is an element $l \in \mathcal{Y}_n$ such that

$$\|\mu(a_i) - l(a_i)\| < \frac{\epsilon}{2}, \quad i = 1, 2, \dots, n.$$

Then $l \circ P_n \in \mathcal{Y}$ and

$$\begin{aligned} d(\mu, l \circ P_n) &\leq \sum_{i=1}^n \frac{2^{-i} \epsilon}{m_i} \frac{1}{2} + \sum_{i \geq n+1} \frac{2^{-i}}{m_i} (1 + \|P_n\|) \\ &\leq \frac{\epsilon}{2} + \sum_{i \geq n+1} \frac{2^{-i}}{m_i} (1 + m_i) \leq \epsilon. \end{aligned}$$

It follows that \mathcal{Y} is a countable set in $\mathcal{B}(A, B)$ with the property that for any linear contraction $\mu : A \rightarrow B$ and any $\epsilon > 0$ there is an element $l \in \mathcal{Y}$ such that $d(\mu, l) < \epsilon$. For each $l \in \mathcal{Y}$ choose a completely positive contraction $l' : A \rightarrow B$ such that

$$d(l, l') \leq 2 \inf\{d(l, L) : L \in \mathcal{B}(A, B) \text{ is a completely positive contraction}\}.$$

Then $\mathcal{Y}' = \{l' : l \in \mathcal{Y}\}$ is a countable set of completely positive contractions in $\mathcal{B}(A, B)$ with the property that for any completely positive linear contraction $\mu : A \rightarrow B$ and any $\epsilon > 0$ there is an element $l \in \mathcal{Y}'$ such that $d(\mu, l) < \epsilon$. \square

Theorem 2.4. *Let A and B be separable C^* -algebras. Assume that B is stable and A unital. Then there exists a unitaly absorbing $*$ -homomorphism $\pi : A \rightarrow \mathcal{M}(B)$ for (A, B) .*

Proof. By Lemma 2.3 there is a dense sequence $\{s_n\}$ in the set of completely positive contractions from A to B . We may assume that each s_n is repeated infinitely often in this sequence. By Kasparov’s Stinespring Theorem (Theorem 3 of [K]), there are elements $V_n \in \mathcal{M}(B)$ and unital $*$ -homomorphisms $\pi_n : A \rightarrow \mathcal{M}(B)$ such that

$$s_n(\cdot) = V_n^* \pi_n(\cdot) V_n$$

for all n . Note that $\|V_n\|^2 = \|V_n^* V_n\| = \|s_n(1)\| \leq 1$ for all n . Define a unital $*$ -homomorphism $\pi_\infty : A \rightarrow \mathcal{L}_B(l_2(B))$ by

$$\pi_\infty(a)(b_1, b_2, b_3, \dots) = (\pi_1(a)b_1, \pi_2(a)b_2, \pi_3(a)b_3, \dots).$$

Define $L_n \in \mathcal{L}_B(B, l_2(B))$ by

$$L_n b = (0, 0, \dots, 0, V_n b, 0, 0, \dots),$$

where the non-trivial entry occurs at the n ’th coordinate. Since we repeated the s_n ’s infinitely often there is, for each n , a sequence $k_1 < k_2 < k_3 < \dots$ in \mathbb{N} such that

$$(2.4) \quad s_n(a) = L_{k_i}^* \pi_\infty(a) L_{k_i}$$

for all $a \in A$, $i \in \mathbb{N}$, and

$$(2.5) \quad \lim_{i \rightarrow \infty} \|L_{k_i}^* \psi\| = 0, \quad \psi \in l_2(B).$$

By Lemma 1.3.2 of [KJT] there is an isomorphism $S : l_2(B) \rightarrow B$ of Hilbert B -modules. Set $T_n = S L_n \in \mathcal{M}(B)$ and $\pi(\cdot) = S \pi_\infty(\cdot) S^*$. We assert that π satisfies condition 1) of Theorem 2.1, and to prove it we let $\varphi : A \rightarrow B$ be a completely positive contraction. In order to construct a sequence $\{W_n\}$ in $\mathcal{M}(B)$ such that 1a) and 1b) of Theorem 2.1 hold it suffices, because A and B are separable, to pick $\epsilon > 0$ and finite subsets $F_1 \subseteq A$ and $F_2 \subseteq B$ and find an element $W \in \mathcal{M}(B)$ such that $\|\varphi(a) - W^* \pi(a) W\| < \epsilon$, $a \in F_1$, and $\|W^* b\| < \epsilon$, $b \in F_2$. Choose first $n \in \mathbb{N}$ such that $\|\varphi(a) - s_n(a)\| < \epsilon$, $a \in F_1$. If we then choose $k_1 < k_2 < k_3 < \dots$

such that (2.4) and (2.5) hold we have that $T_{k_i}^* \pi(a) T_{k_i} = s_n(a)$ for all $a \in F_1$ and $\|T_{k_i}^* b\| < \epsilon$ for all $b \in F_2$, provided only that i is large enough. We can then set $W = T_{k_i}$ for such an i . □

We now turn to the case of a not necessarily unital C^* -algebra A and the general notion of absorbing $*$ -homomorphisms. Given a C^* -algebra A we denote in the following by A^+ the C^* -algebra obtained by adding a unit to A . Let B be another C^* -algebra. Any linear completely positive contraction $\varphi : A \rightarrow \mathcal{M}(B)$ admits a unique linear extension $\varphi^+ : A^+ \rightarrow \mathcal{M}(B)$ such that $\varphi^+(1) = 1$. φ^+ is automatically a completely positive contraction (cf. e.g. Lemma 3.2.8 of [KJT]), and is automatically a $*$ -homomorphism when φ is. The following theorem is therefore an immediate consequence of Theorem 2.1.

Theorem 2.5. *Let A and B be separable C^* -algebras with B stable. Let $\pi : A \rightarrow \mathcal{M}(B)$ be a $*$ -homomorphism. Then the following conditions are equivalent:*

- 1) $\pi^+ : A^+ \rightarrow \mathcal{M}(B)$ is unittally absorbing for (A^+, B) .
- 2) For any completely positive contraction $\varphi : A \rightarrow \mathcal{M}(B)$ there is a sequence $\{V_n\}$ of isometries in $\mathcal{M}(B)$ such that
 - 2a) $V_n^* \pi(a) V_n - \varphi(a) \in B$, $n \in \mathbb{N}$, $a \in A$,
 - 2b) $\lim_{n \rightarrow \infty} \|V_n^* \pi(a) V_n - \varphi(a)\| = 0$, $a \in A$.
- 3) For any $*$ -homomorphism $\varphi : A \rightarrow \mathcal{M}(B)$ there is a sequence $\{U_n\}$ of unitaries $U_n \in \mathcal{L}_B(B \oplus B, B)$ such that
 - 3a) $U_n \begin{pmatrix} \pi(a) & 0 \\ 0 & \varphi(a) \end{pmatrix} U_n^* - \pi(a) \in B$, $n \in \mathbb{N}$, $a \in A$,
 - 3b) $\lim_{n \rightarrow \infty} \|U_n \begin{pmatrix} \pi(a) & 0 \\ 0 & \varphi(a) \end{pmatrix} U_n^* - \pi(a)\| = 0$, $a \in A$.
- 4) For any $*$ -homomorphism $\varphi : A \rightarrow \mathcal{M}(B)$ there is a sequence $\{U_n\}$ of unitaries $U_n \in \mathcal{L}_B(B \oplus B, B)$ such that

$$\lim_{n \rightarrow \infty} \|U_n \begin{pmatrix} \pi(a) & 0 \\ 0 & \varphi(a) \end{pmatrix} U_n^* - \pi(a)\| = 0, a \in A.$$

Definition 2.6. Let A and B be separable C^* -algebras with B stable. A $*$ -homomorphism $\pi : A \rightarrow \mathcal{M}(B)$ is *absorbing* (for (A, B)) when it satisfies the four equivalent conditions of Theorem 2.5.

Theorem 2.7. *Let A and B be separable C^* -algebras with B stable. There exists an absorbing $*$ -homomorphism $\pi : A \rightarrow \mathcal{M}(B)$ for (A, B) .*

Proof. Combine Theorem 2.5 and Theorem 2.4. □

An absorbing $*$ -homomorphism is clearly unique in the following sense: Given two absorbing $*$ -homomorphisms $\pi_1, \pi_2 : A \rightarrow \mathcal{M}(B)$ there is a sequence $\{U_n\} \subseteq \mathcal{M}(B)$ of unitaries such that $U_n \pi_1(a) U_n^* - \pi_2(a) \in B$, $a \in A$, $n \in \mathbb{N}$, and $\lim_{n \rightarrow \infty} U_n \pi_1(a) U_n^* - \pi_2(a) = 0$, $a \in A$.

3. DUALITY IN KK -THEORY

Throughout this section A and B will be separable C^* -algebras and B will be stable. A $*$ -homomorphism $\pi : A \rightarrow \mathcal{M}(B)$ is of *infinite multiplicity* when π is unitarily equivalent to π^∞ , where $\pi^\infty : A \rightarrow \mathcal{M}(B)$ is the $*$ -homomorphism given by $\pi^\infty(a) = \sum_{i=1}^\infty S_i \pi(a) S_i^*$, for some sequence S_i , $i \in \mathbb{N}$, of isometries in $\mathcal{M}(B)$ such that $S_i^* S_j = 0$, $i \neq j$, and $\sum_{i=1}^\infty S_i S_i^* = 1$ in the strict topology.

Lemma 3.1. *Let $\pi : A \rightarrow \mathcal{M}(B)$ be a $*$ -homomorphism of infinite multiplicity and set*

$$E = \{m \in \mathcal{M}(B) : m\pi(a) = \pi(a)m \ \forall a \in A\} .$$

Then $K_*(E) = \{0\}$.

Proof. Since π has infinite multiplicity,

$$E \simeq \{m \in \mathcal{L}_B(l_2(B)) : m\mu(a) = \mu(a)m \ \forall a \in A\}$$

where $\mu : A \rightarrow \mathcal{L}_B(l_2(B))$ is given by

$$\mu(a)(b_1, b_2, b_3, \dots) = (\pi(a)b_1, \pi(a)b_2, \pi(a)b_3, \dots).$$

The usual proof that $K_*(\mathcal{L}_B(l_2(B))) = 0$ works to show that $K_*(E) = 0$; cf. e.g. Proposition 12.2.1 of [Bl]. □

Given an absorbing $*$ -homomorphism $\pi : A \rightarrow \mathcal{M}(B)$ we set

$$C_\pi = \{x \in \mathcal{M}(B) : x\pi(a) - \pi(a)x \in B, \ a \in A\}$$

and

$$A_\pi = \{x \in C_\pi : x\pi(A) \subseteq B\} .$$

Then A_π is a closed two-sided ideal in C_π and we set $D_\pi = C_\pi/A_\pi$. The quotient map $C_\pi \rightarrow D_\pi$ will be denoted by q . If $\tau : A \rightarrow \mathcal{M}(B)$ is another absorbing $*$ -homomorphism there is a unitary $w \in \mathcal{M}(B)$ such that $\text{Ad } w \circ \pi(a) - \tau(a) \in B$ for all $a \in A$ and then $x \mapsto wxw^*$ defines a $*$ -isomorphism of C_π onto C_τ which takes A_π onto A_τ . In particular, $D_\pi \simeq D_\tau$.

Let u be a unitary in $M_n(D_\pi)$. Choose $v \in M_n(C_\pi)$ such that $\text{id}_{M_n} \otimes q(v) = u$. Define $\pi^n : A \rightarrow \mathcal{L}_B(B^n)$ by $\pi^n(a)(b_1, b_2, \dots, b_n) = (\pi(a)b_1, \pi(a)b_2, \dots, \pi(a)b_n)$. Let $B^n \oplus B^n$ be graded by $(x, y) \mapsto (x, -y)$. Then

$$(B^n \oplus B^n, \begin{pmatrix} \pi^n & \\ & \pi^n \end{pmatrix}, (v^* \ v))$$

is a Kasparov $A - B$ -module. We leave it to the reader to check that the class of this module in $KK(A, B)$ only depends on the class of u in $K_1(D_\pi)$, and that the construction gives rise to a group homomorphism $\Theta : K_1(D_\pi) \rightarrow KK(A, B)$.

Theorem 3.2. *Assume that $\pi : A \rightarrow \mathcal{M}(B)$ is an absorbing $*$ -homomorphism. Then $\Theta : K_1(D_\pi) \rightarrow KK(A, B)$ is an isomorphism.*

Proof. When τ is another absorbing $*$ -homomorphism there is a commuting diagram

$$(3.1) \quad \begin{array}{ccc} K_1(D_\pi) & \xrightarrow{\Theta} & KK(A, B) \\ \downarrow & \nearrow & \\ K_1(D_\tau) & & \end{array}$$

where $K_1(D_\pi) \rightarrow K_1(D_\tau)$ is induced by the isomorphism $D_\pi \rightarrow D_\tau$ described above, and $K_1(D_\tau) \rightarrow KK(A, B)$ is the map obtained by using τ instead of π in the definition of Θ . Indeed if one considers a specific unitary in $M_n(D_\pi)$, the Kasparov $A - B$ -module which results by going down and up in the diagram differs from the one which arises by going across by an isomorphism and a compact perturbation. Thus if we prove that $\Theta : K_1(A_\pi) \rightarrow KK(A, B)$ is an isomorphism for one absorbing

*-homomorphism π it will follow that it is an isomorphism for any other. Hence by working with π^∞ instead of π we may assume that π is of infinite multiplicity.

Θ is injective: Let $u \in M_n(D_\pi)$ be a unitary and choose $v \in M_n(C_\pi)$ such that $\text{id}_{M_n} \otimes q(v) = u$. Assume that $[B^n \oplus B^n, (\pi^n \ \pi^n), (v^* \ v)] = 0$ in $KK(A, B)$. This means that there are degenerate Kasparov $A - B$ -modules \mathcal{D}_1 and \mathcal{D}_2 such that $(B^n \oplus B^n, (\pi^n \ \pi^n), (v^* \ v)) \oplus \mathcal{D}_1$ is operator homotopic to $(B^n \oplus B^n, (\pi^n \ \pi^n), (1 \ 1)) \oplus \mathcal{D}_2$. Since \mathcal{D}_1 and \mathcal{D}_2 are degenerate we can define a new degenerate Kasparov $A - B$ -module \mathcal{D} by $\mathcal{D} = \mathcal{D}_1 \oplus \mathcal{D}_2 \oplus \mathcal{D}_1 \oplus \mathcal{D}_2 \oplus \mathcal{D}_1 \oplus \mathcal{D}_2 \oplus \dots$. Then $\mathcal{D}_1 \oplus \mathcal{D}$ and $\mathcal{D}_2 \oplus \mathcal{D}$ are both isomorphic to \mathcal{D} and hence $(B^n \oplus B^n, (\pi^n \ \pi^n), (v^* \ v)) \oplus \mathcal{D}$ is operator homotopic to $(B^n \oplus B^n, (\pi^n \ \pi^n), (1 \ 1)) \oplus \mathcal{D}$. By combining Kasparov's stabilization theorem (Theorem 2.12 of [KJT]) with Lemma 1.3.2 of [KJT] we may assume that $a = w$ and $b = w^*$ for some unitary $w \in \mathcal{M}(B)$. Finally, by applying the unitary of the Hilbert B -module $B \oplus B$ given by $(x, y) \mapsto (x, wy)$, we see that we can assume that $w = 1$. So all in all we have that

$$(B^n \oplus B^n, (\pi^n \ \pi^n), (v^* \ v)) \oplus (B \oplus B, \begin{pmatrix} \lambda_+ & \\ & \lambda_- \end{pmatrix}, (1 \ 1))$$

is operator homotopic to

$$(B^n \oplus B^n, (\pi^n \ \pi^n), (1 \ 1)) \oplus (B \oplus B, \begin{pmatrix} \lambda_+ & \\ & \lambda_- \end{pmatrix}, (1 \ 1)).$$

Note that $\lambda_+ = \lambda_-$ since $(B \oplus B, \begin{pmatrix} \lambda_+ & \\ & \lambda_- \end{pmatrix}, (1 \ 1))$ is degenerate. Finally, by adding on an infinite number of copies of $(B \oplus B, \begin{pmatrix} \lambda_+ & \\ & \lambda_- \end{pmatrix}, (1 \ 1))$ we find that there is a *-homomorphism of infinite multiplicity $\lambda : A \rightarrow \mathcal{M}(B)$ such that

$$(B^n \oplus B^n, (\pi^n \ \pi^n), (v^* \ v)) \oplus (B \oplus B, (\lambda \ \lambda), (1 \ 1))$$

is operator homotopic to

$$(B^n \oplus B^n, (\pi^n \ \pi^n), (1 \ 1)) \oplus (B \oplus B, (\lambda \ \lambda), (1 \ 1)).$$

Furthermore, by adding on $(B \oplus B, (\pi \ \pi), (1 \ 1))$ we may assume that there is a unitary $w \in \mathcal{M}(B)$ such that $w\lambda(a)w^* - \pi(a) \in B$, $a \in A$. The operator homotopy consists of an isomorphism of Kasparov $A - B$ modules and a norm-continuous path of operators. The isomorphism gives us a unitary $S \in M_{n+1}(\mathcal{M}(B))$ such that $S \begin{pmatrix} \pi^n(a) & \\ & \lambda(a) \end{pmatrix} = \begin{pmatrix} \pi^n(a) & \\ & \lambda(a) \end{pmatrix} S$ for all $a \in A$, and in addition we have a norm-continuous path F_t , $t \in [0, 1]$, in $M_{n+1}(\mathcal{M}(B))$ such that $F_0 = S$, $F_1 = \begin{pmatrix} v & \\ & 1 \end{pmatrix}$, and $(F_t F_t^* - 1_{n+1}) \begin{pmatrix} \pi^n(a) & \\ & \lambda(a) \end{pmatrix}$, $(F_t^* F_t - 1_{n+1}) \begin{pmatrix} \pi^n(a) & \\ & \lambda(a) \end{pmatrix}$, $F_t \begin{pmatrix} \pi^n(a) & \\ & \lambda(a) \end{pmatrix} - \begin{pmatrix} \pi^n(a) & \\ & \lambda(a) \end{pmatrix} F_t$ are in $M_{n+1}(B)$ for all t and a . Here and in the following we let 1_k denote the unit of $M_k(\mathcal{M}(B))$. Note that $\nu = \begin{pmatrix} \pi^n & \\ & \lambda \end{pmatrix}$ is of infinity multiplicity, as a *-homomorphism $A \rightarrow \mathcal{M}(M_{n+1}(B))$, since π and λ both are of infinite multiplicity. By Lemma 3.1 we can therefore find an $m \in \mathbb{N}$ and a norm-continuous path of unitaries in $\{x \in M_{m(n+1)}(\mathcal{M}(B)) : x\nu^m(a) = \nu^m(a)x, a \in A\}$ connecting $\begin{pmatrix} S & \\ & 1_{(m-1)(n+1)} \end{pmatrix}$ to $1_{m(n+1)}$. In combination with F this gives us a norm-continuous path H_t , $t \in [0, 1]$, in $M_{m(n+1)}(\mathcal{M}(B))$ such that $H_0 = 1_{m(n+1)}$, $H_1 = \begin{pmatrix} v & \\ & 1_{m(n+1)-n} \end{pmatrix}$, $(H_t H_t^* - 1_{m(n+1)})\nu^m(a)$, $(H_t^* H_t - 1_{m(n+1)})\nu^m(a)$, $H_t \nu^m(a) - \nu^m(a)H_t$ are in $M_{m(n+1)}(B)$ for all t and a . Set

$$W = \text{diag}(\underbrace{1_n, w, 1_n, w, \dots, 1_n, w}_{m \text{ times}}) \in M_{m(n+1)}(\mathcal{M}(B))$$

and $G_t = WH_tW^*$. Then G_t is a norm-continuous path in $M_{m(n+1)}(\mathcal{M}(B))$ such that $G_0 = 1_{m(n+1)}$, $G_1 = \begin{pmatrix} v & \\ & 1_{m(n+1)-n} \end{pmatrix}$ and $(G_tG_t^* - 1_{m(n+1)})\pi^{m(n+1)}(a)$, $(G_t^*G_t - 1_{m(n+1)})\pi^{m(n+1)}(a)$, $G_t\pi^{m(n+1)}(a) - \pi^{m(n+1)}(a)G_t$ are in $M_{m(n+1)}(B)$ for all t and a . Thus $(\text{id}_{M_{m(n+1)}} \otimes q)(G_t)$ is a path of unitaries in $M_{m(n+1)}(D\pi)$ connecting $\begin{pmatrix} u & \\ & 1_{m(n+1)-n} \end{pmatrix}$ to $1_{m(n+1)}$.

Θ is surjective: Let (E, ψ, F) be a Kasparov $A - B$ -module. The constructions on pages 125-126 of [KJT] show that $[E, \psi, F] \in KK(A, B)$ is also represented by a Kasparov $A - B$ -module of the form $(B \oplus B, \begin{pmatrix} \varphi_+ & \\ & \varphi_- \end{pmatrix}, \begin{pmatrix} v^* & v \end{pmatrix})$ for some $*$ -homomorphisms $\varphi_{\pm} : A \rightarrow \mathcal{M}(B)$ and some unitary $v \in \mathcal{M}(B)$. Using the trick from p. 354 of [H] we may assume that $\varphi_- = \varphi_+ = \varphi$. By adding on $(B \oplus B, \begin{pmatrix} \pi & \\ & \pi \end{pmatrix}, \begin{pmatrix} 1 & \\ & 1 \end{pmatrix})$ and using that π is absorbing we may assume that there is a unitary $u \in \mathcal{M}(B)$ such that $u\varphi(a)u^* - \pi(a) \in B$ for all $a \in A$. Then $(B \oplus B, \begin{pmatrix} \varphi & \\ & \varphi \end{pmatrix}, \begin{pmatrix} v^* & v \end{pmatrix})$ is isomorphic to

$$\left(B \oplus B, \begin{pmatrix} \text{Ad } u \circ \varphi & \\ & \text{Ad } u \circ \varphi \end{pmatrix}, \begin{pmatrix} uv^*u^* & uvu^* \end{pmatrix} \right)$$

which in turn is a compact perturbation of $(B \oplus B, \begin{pmatrix} \pi & \\ & \pi \end{pmatrix}, \begin{pmatrix} uv^*u^* & uvu^* \end{pmatrix})$. Then uvu^* is a unitary C_{π} such that $\Theta([q(uvu^*)]) = [E, \psi, F]$ in $KK(A, B)$. \square

Of course there is also an isomorphism

$$K_0(D\pi) \simeq \text{Ext}^{-1}(A, B)$$

which can be proved in basically the same way.

REFERENCES

- [A] W. Arveson, *Notes on extensions*, Duke Math. J. **44** (1977), 329-355. MR **55**:11056
- [Bl] B. Blackadar, *K-theory for Operator Algebras*, MSRI publications, Springer Verlag, New York, 1986. MR **88g**:46082
- [H] N. Higson, *C*-Algebra Extension Theory and Duality*, J. Func. Analysis **129** (1995), 349-363. MR **96c**:46072
- [K] G. Kasparov, *Hilbert C*-modules: theorems of Stinespring and Voiculescu*, J. Oper. Th. **4** (1980), 133-150. MR **82b**:46074
- [KJT] K. Knudsen-Jensen and K. Thomsen, *Elements of KK-theory*, Birkhäuser, Boston, 1991. MR **94b**:19008
- [P] W. Paschke, *K-theory for commutants in the Calkin algebra*, Pacific J. Math. **95** (1981), 427-437. MR **82k**:46101
- [S] G. Skandalis, *Une Notion de Nucléarité en K-Théorie (d'après J. Cuntz)*, K-theory **1** (1988), 549-573. MR **90b**:46131
- [V] A. Valette, *A remark on the Kasparov Groups $\text{Ext}^i(A, B)$* , Pacific J. Math. **109** (1983), 247-255. MR **85f**:46128

INSTITUT FOR MATEMATISKE FAG, NY MUNKEGADE, 8000 AARHUS C, DENMARK
E-mail address: matkt@imf.au.dk