

## ON THE HOMOLOGY OF SPLIT EXTENSIONS WITH $p$ -ELEMENTARY KERNEL

MARCIN CHALUPNIK

(Communicated by Ralph Cohen)

ABSTRACT. We study a Hochschild–Serre spectral sequence associated to a split group extension with kernel  $(\mathbf{Z}/p)^n$ . It is shown that a large part of  $E_2^{0*}$  must survive to infinity. We also sketch the general procedure of computing this surviving group.

### 1. INTRODUCTION

It is often useful to decompose a spectral sequence into eigenspaces of an automorphism of the sequence. In the case of a Hochschild–Serre spectral sequence associated to a split extension with abelian kernel the Liebermann trick (see [Sa], p. 262) provides an important example of this. One takes the automorphism induced by multiplication by a scalar in the kernel of the extension. For example, it is easy using this method to show that in a split extension with abelian kernel

$$0 \rightarrow A \rightarrow H \rightarrow G \rightarrow 1$$

we have  $H_i(H, \mathbf{Q}) = \bigoplus_{0 \leq j \leq i} (H_{i-j}(G, \Lambda^j(A \otimes \mathbf{Q})))$ . If one considers the homology with  $\mathbf{F}_p$ -coefficients, the situation becomes more involved. The first problem is that scalars have only finite multiplicative order and the second is that the homology of an abelian group also contains a part generated by elements of degree 2 (for  $p$  odd). For these reasons a Hochschild–Serre spectral sequence can have many nontrivial differentials and is generally hard to understand.

In the present paper we use the Liebermann trick together with an analysis of a scalar extension to show triviality of some differentials when one takes  $\mathbf{F}_p$ -coefficients. We apply this technique in Section 2 to show that a part of the 0-th column survives which is close to  $\Lambda^*(A)_G$ . In Section 3 we discuss some examples, in particular we show that  $\Lambda^*(A)_G$  does not always embed into  $H_*(H, \mathbf{F}_p)$ .

It is a pleasure to thank Stanislaw Betley for many interesting discussions and strong encouragement.

### 2. THE THEOREM

Let

$$(1) \quad 0 \rightarrow A \rightarrow H \rightarrow G \rightarrow 1$$

---

Received by the editors August 25, 1999 and, in revised form, March 8, 2000.

2000 *Mathematics Subject Classification*. Primary 20J06.

The author was partially supported by the Polish scientific grant (KBN) 2 P03A 01113.

be a split extension with  $A = (\mathbf{Z}/p)^n$  (we specialize to this case only to simplify notation, for general (abelian)  $A$  our Theorem 1 remains true if we replace  $A$  by  $A \otimes_{\mathbf{Z}} \mathbf{F}_p$ ). We will consider a homological Hochschild–Serre spectral sequence with  $\mathbf{F}_p$ -coefficients corresponding to this extension. Denoting by  $D^j$  a  $j$ -th divided power we have the natural identification of the  $E^2$ -term:

$$(2) \quad E_{ij}^2 = \bigoplus_{k+2l=j} H_i(G, \Lambda^k(A) \otimes D^l(A)) \quad \text{for } p \text{ odd,}$$

$$(3) \quad E_{ij}^2 = H_i(G, D^j(A)) \quad \text{for } p = 2$$

(unless otherwise stated  $\otimes$  means  $\otimes_{\mathbf{F}_p}$ ).

Let us define  $H_i(G, \Lambda^*(A))_{reg}$  to be

$$H_i(G, \Lambda^*(A))/\ker(H_i(G, \Lambda^*(A)) \rightarrow H_i(G, A^{\otimes*}))$$

where the arrow is induced by the natural embedding  $\Delta : \Lambda^*(A) \hookrightarrow A^{\otimes*}$ . Similarly, we put

$$H_i(G, D^*(A))_{reg} = H_i(G, D^*(A))/\ker(H_i(G, D^*(A)) \rightarrow H_i(G, A^{\otimes*}))$$

for  $\Gamma : D^*(A) \hookrightarrow A^{\otimes*}$ .

In our spectral sequence only some pieces of the groups we are interested in survive, hence we should carefully differ between  $E^2$  and the higher  $E^r$ . Thus, in order to make the formulation of Theorem 1 clear and to avoid a confusion in its proof we introduce some notation. For  $i \leq 1$  we have natural epimorphisms  $\alpha_{ij}^r : E_{ij}^2 \rightarrow E_{ij}^r$ . Then we put  $B_{ij} = \bigcup_{r \geq 2} \ker(\alpha_{ij}^r) \cap H_i(G, \Lambda^j(A))$  for  $p$  odd and  $B_{ij} = \bigcup_{r \geq 2} \ker(\alpha_{ij}^r) \cap H_i(G, D^j(A))$  for  $p = 2$ .

**Theorem 1.** *In the sequence (2) we have  $B_{ij} \subset \ker(\Delta_*)$  for  $i \leq 1$ . In other words, the spaces  $H_i(G, \Lambda^j(A))_{reg}$  for  $i \leq 1$  survive to infinity. The same holds for the sequence (3) when we replace  $\Lambda^j(A)$  by  $D^j(A)$ .*

*Proof.* We begin with some remarks concerning functoriality of semidirect products. It is well known that the class of split extensions of a fixed group  $G$  by abelian kernels is in bijection with the class of  $\mathbf{Z}[G]$ -modules via construction of semidirect product. Moreover, this assignment yields an isomorphism of the category of  $\mathbf{Z}[G]$ -modules and the category of split extensions of  $G$  with abelian kernels where morphisms are morphisms of extensions being identity on  $G$ . The practical consequence is that any  $G$ -homomorphism between kernels of two extensions induces a morphism of spectral sequences.

The idea of the proof (for  $p$  odd) is as follows. We look at the automorphism of the spectral sequence (2) induced by the  $G$ -automorphism of  $A$  defined by the formula  $x \mapsto cx$  for a given scalar  $c \in \mathbf{F}_p^*$  (we will frequently use the structure of  $\mathbf{F}_p$ -linear space on  $A$ ). Then it is easy to see that the space

$$H_*(G, \Lambda^k(A) \otimes D^l(A))$$

belongs to the eigenspace of the induced automorphism for the eigenvalue  $c^{k+l}$  and that the whole spectral sequence is a direct sum of eigensequences for eigenvalues  $1, c, c^2, \dots, c^{p-1}$ . At this point it is clear for example that there are no differentials coming to  $H_*(G, \Lambda^k(A))$  for  $k < p$  because all  $E_{ij}^*$  for  $j < k$  belong to eigenspaces of  $c^s$  for  $s < k$  and differentials must preserve the decomposition. Thus  $H_0(G, \Lambda^k(A))$  and  $H_1(G, \Lambda^k(A))$  for  $k < p$  survives. Unfortunately, this argument fails for  $k \geq p$

since  $c^p = c$  for any  $c \in \mathbf{F}_p^*$ . We partially overcome this difficulty comparing the sequence (2) with a sequence associated to a split extension with kernel equipped with  $G$ -automorphism of higher order.

Therefore let us consider a split extension

$$0 \rightarrow A \otimes L \rightarrow H(L) \rightarrow G \rightarrow 1$$

where  $L$  is a one-dimensional space over a field  $\mathbf{F}_q$  with  $q = p^d$  elements regarded as a trivial  $G$ -module. We shall describe  $'E^2$ —the second page of a Hochschild–Serre spectral sequence (with coefficients in  $\mathbf{F}_q$ ) associated to it. We focus here on the case when  $p$  is odd. We should take into account the  $\mathbf{F}_q$ -structure appearing in this new sequence. More precisely, we describe  $'E_{**}^2$  as evaluations on  $L$  of functors from the category of finite  $\mathbf{F}_q$ -spaces to itself assigning to a  $\mathbf{F}_q$ -space  $V$  the entries in the spectral sequence associated to the extension

$$0 \rightarrow A \otimes V \rightarrow H(V) \rightarrow G \rightarrow 1.$$

According to a functorial description of the homology of an abelian group (see e.g. [Qu2], p. 210) and the natural isomorphism  $V \otimes \mathbf{F}_q = \bigoplus_{t=0}^{d-1} V^{(t)}$  (here  $V^{(t)}$  means the space  $V$  with  $\mathbf{F}_q$ -structure twisted by  $t$ -th Frobenius) we get

$$\begin{aligned} (4) \quad 'E_{ij}^2 &= \bigoplus_{k+2l=j} H_i(G, \Lambda^k(A \otimes L \otimes \mathbf{F}_q) \otimes_{\mathbf{F}_q} D^l(A \otimes L \otimes \mathbf{F}_q)) \\ &= \bigoplus_{k+2l=j} H_i\left(G, \Lambda^k\left(\bigoplus_{t=0}^{d-1} A \otimes L^{(t)}\right) \otimes_{\mathbf{F}_q} D^l\left(\bigoplus_{t=0}^{d-1} A \otimes L^{(t)}\right)\right) \\ &= \bigoplus_{k+2l=j} \bigoplus_{\sum k_t=k} \bigoplus_{\sum l_t=l} H_i\left(G, \Lambda^{k_0}(A \otimes L^{(0)}) \otimes_{\mathbf{F}_q} \dots \otimes_{\mathbf{F}_q} \Lambda^{k_{d-1}}(A \otimes L^{(d-1)}) \right. \\ &\quad \left. \otimes_{\mathbf{F}_q} D^{l_0}(A \otimes L^{(0)}) \otimes_{\mathbf{F}_q} \dots \otimes_{\mathbf{F}_q} D^{l_{d-1}}(A \otimes L^{(d-1)})\right) \end{aligned}$$

(we alert the reader that in these formulas exterior and divided powers are taken over  $\mathbf{F}_q$ ). Now take a scalar  $c \in \mathbf{F}_q^*$  of multiplicative order  $p^d - 1$ . We define the  $G$ -automorphism of  $A \otimes L$  by the formula  $a \otimes x \mapsto a \otimes cx$ . In order to understand the induced automorphism of the spectral sequence (4) observe that multiplication by  $c$  on  $L$  induces on  $A \otimes L^{(t)}$  multiplication by  $c^{p^t}$ . Thus the space

$$H_*\left(G, \bigotimes_{t=0}^{d-1} \Lambda^{k_t}(A \otimes L^{(t)}) \otimes_{\mathbf{F}_q} \bigotimes_{t=0}^{d-1} D^{l_t}(A \otimes L^{(t)})\right)$$

(big tensor products are over  $\mathbf{F}_q$ ) belongs to the eigenspace of the eigenvalue  $c^{\sum_{t=0}^{d-1} (k_t + l_t)p^t}$ . The crucial fact is that here exponents in eigenvalues are taken modulo  $p^d - 1$  hence more differentials must be trivial than in sequence (2). So our next task will be to compare both spectral sequences. Before doing this however, we introduce some notation. For a sequence of nonnegative integers  $\mathbf{k} = (k_0, \dots, k_{d-1})$  we define  $r(\mathbf{k})$  to be the number  $r(\mathbf{k}) = \sum_t k_t$  and  $|\mathbf{k}|$  to be  $\sum_t k_t p^t \pmod{p^d - 1}$ . The following elementary arithmetic lemma holds

**Lemma 1.** *Let  $0 \leq j < d(p - 1)$ . Then there exists  $\mathbf{k}$  and a number  $a$  such that  $r(\mathbf{k}) = j$ ,  $|\mathbf{k}| = a$  and  $|\mathbf{k}'| \neq a$  for any  $\mathbf{k}'$  such that  $r(\mathbf{k}') \leq j$ .*

*Proof.* Let  $j = f(p - 1) + g$  where  $g < p - 1$ . We put

$$k_i = \begin{cases} 0 & \text{for } i < d - f - 1, \\ g & \text{for } i = d - f - 1, \\ p - 1 & \text{for } d - f - 1 < i \leq d - 1. \end{cases}$$

We are going to show that for any  $\mathbf{k}'$  satisfying  $|\mathbf{k}'| = |\mathbf{k}|$  we have  $r(\mathbf{k}') > r(\mathbf{k})$ . Let us take such  $\mathbf{k}'$ . If there exists  $k'_{i_0} \geq p$ , we may replace  $\mathbf{k}'$  by  $\mathbf{k}''$  having the same  $|\cdot|$  but smaller  $r$  defining

$$k''_i = \begin{cases} k'_i - p & \text{for } i = i_0, \\ k'_{i_0+1} + 1 & \text{for } i = i_0 + 1, \\ k'_i & \text{for } i \neq i_0, i_0 + 1 \end{cases}$$

(we use here the convention  $k_d = k_0$ ). Thus, iterating this procedure, we may assume that all  $k'_i$  are smaller than  $p$ . But in this case the only possibility for  $|\mathbf{k}'| = |\mathbf{k}|$  is  $\mathbf{k}' = \mathbf{k}$ .

Let  $\Lambda^{\mathbf{k}}(A \otimes L)$  denote  $\Lambda^{k_0}(A \otimes L) \otimes \dots \otimes \Lambda^{k_{d-1}}(A \otimes L^{(d-1)})$ . If for  $j < d(p - 1)$  we take  $\mathbf{k}$  as in Lemma 1, then there cannot be any differentials coming to  $H_*(G, \Lambda^{\mathbf{k}}(A \otimes L))$  (and its subspaces in the higher  $'E^r$ ). Thus the spaces  $H_i(G, \Lambda^{\mathbf{k}}(A \otimes L))$  for  $i \leq 1$  survive in sequence (4).

We now want to construct a morphism from the spectral sequence (2) to (4). First we should replace (2) by (5)—its counterpart with  $\mathbf{F}_q$ -coefficients. In this new sequence we have

$$(5) \quad {}''E^2_{i,j} = \bigoplus_{k+2l=j} H_i(G, \Lambda^k(A \otimes \mathbf{F}_q) \otimes_{\mathbf{F}_q} D^l(A \otimes \mathbf{F}_q)).$$

There is a morphism  $\Phi$  from (2) to (5) which is, by the Kunneth formula, on  $E^2$  just induced by scalar extension in all tensors appearing as the coefficients of the homology of  $G$ . Now to obtain a morphism from (5) to (4) it suffices to choose a  $G$ -homomorphism from  $A$  to  $A \otimes L$  which is possible since  $L$  is a trivial  $G$ -module. In order to make formulas explicit let us identify  $L$  with  $\mathbf{F}_q$ . Then we take the homomorphism from  $A$  to  $A \otimes \mathbf{F}_q$  sending  $a$  to  $a \otimes 1$  and we will consider the morphism of spectral sequences  $\Psi$  induced by this  $G$ -homomorphism. Under the isomorphism  $(A \otimes \mathbf{F}_q) \otimes_{\mathbf{F}_q} = \bigoplus_{t=0}^{d-1} (A \otimes \mathbf{F}_q^{(t)})$  the morphism  $\Psi$  from (5) to (4) is induced on  $E^2$  by the morphism of coefficients  $\psi : A \otimes \mathbf{F}_q \rightarrow \bigoplus_{t=0}^{d-1} (A \otimes \mathbf{F}_q^{(t)})$  sending  $a \otimes x$  to  $\bigoplus_{t=0}^{d-1} (a \otimes x^{p^t})$ . We focus on the groups  $H_*(G, \Lambda^j(A \otimes \mathbf{F}_q))$ . We would like to describe the map  $\pi_{\mathbf{k}} \circ \psi_* : H_*(G, \Lambda^j(A \otimes \mathbf{F}_q)) \rightarrow H_*(G, \Lambda^{\mathbf{k}}(A \otimes \mathbf{F}_q))$  where for given a sequence  $\mathbf{k}$  with  $r(\mathbf{k}) = j$  the map  $\pi_{\mathbf{k}}$  is the projection from  $'E^2_{*j}$  onto  $H_*(G, \Lambda^{\mathbf{k}}(A \otimes \mathbf{F}_q))$ . According to the above formulas,  $\pi_{\mathbf{k}} \circ \psi_*$  may be factorized as  $f_* \circ com_*$  where

$$com_* : H_*(G, \Lambda^j(A \otimes \mathbf{F}_q)) \rightarrow H_*(G, \Lambda^{k_0}(A \otimes \mathbf{F}_q) \otimes \dots \otimes \Lambda^{k_{d-1}}(A \otimes \mathbf{F}_q))$$

is induced by the iterated comultiplication map in the  $\mathbf{F}_q$ -Hopf algebra  $\Lambda_{\mathbf{F}_q}^*(A \otimes \mathbf{F}_q)$  while

$$\begin{aligned} f_* : H_*(G, \Lambda^{k_0}(A \otimes \mathbf{F}_q) \otimes \dots \otimes \Lambda^{k_{d-1}}(A \otimes \mathbf{F}_q)) \\ \rightarrow H_*(G, \Lambda^{k_0}(A \otimes \mathbf{F}_q^{(0)}) \otimes \dots \otimes \Lambda^{k_{d-1}}(A \otimes \mathbf{F}_q^{(d-1)})) \end{aligned}$$

is determined by the  $G$ -isomorphism  $f_{k_0} \otimes \dots \otimes f_{k_{d-1}}$  defined on a factor  $\Lambda^{k_t}$  by the formula  $f_{k_t}(a \otimes x) = a \otimes x^{p^t}$ .

Now we are in a position to prove the theorem. Given  $x \in H_i(G, \Lambda^j(A))$  ( $i \leq 1$ ) belonging to  $\ker(\alpha_{ij}^r)$ , choose  $\mathbf{k}$  as in Lemma 1 and consider the commutative diagram

$$\begin{array}{ccccc}
 E_{ij}^2 & \xrightarrow{\psi_* \circ \Phi^2} & {}'E_{ij}^2 & \xrightarrow{\pi_{|\mathbf{k}|}^2} & {}_{|\mathbf{k}|}'E_{ij}^r \\
 \alpha_{ij}^r \downarrow & & {}'\alpha_{ij}^r \downarrow & & {}'\alpha_{ij}^r \downarrow \\
 E_{ij}^r & \xrightarrow{\Psi^r \circ \Phi^r} & {}'E_{ij}^r & \xrightarrow{\pi_{|\mathbf{k}|}^r} & {}_{|\mathbf{k}|}'E_{ij}^r
 \end{array}$$

where  ${}_{|\mathbf{k}|}'E$  denotes the subsequence corresponding to the eigenvalue  $|\mathbf{k}|$  and  $\pi_{|\mathbf{k}|}$  is a natural projection (it is a morphism of spectral sequences in contrast to  $\pi_{\mathbf{k}}$ ). Now suppose that

$${}'\alpha_{ij}^r \circ \pi_{|\mathbf{k}|}^2 \circ \psi_* \circ \Phi^2(x) = 0.$$

Since  $\pi_{|\mathbf{k}|}^2 \circ \psi_* \circ \Phi^2(x) = \pi_{\mathbf{k}} \circ \psi_* \circ \Phi^2(x) \in H_i(G, \Lambda^{\mathbf{k}}(A \otimes \mathbf{F}_q))$ , which by the paragraph after Lemma 1 survives to infinity, we thus obtain

$$\pi_{\mathbf{k}} \circ \psi_* \circ \Phi^2(x) = 0.$$

Now by identifications we have worked out earlier we get

$$0 = \pi_{\mathbf{k}} \circ \psi_* \circ \Phi^2(x) = f_* \circ \text{com}_* \circ \Phi^2(x).$$

But since  $f_*$  is an isomorphism, we have

$$\text{com}_* \circ \Phi^2(x) = 0.$$

At last, by the Kunneth formula,  $\ker(\text{com}_* \circ \Phi^2) = \ker(\text{com}'_*)$  where  $\text{com}'_*$  is iterated comultiplication in  $\mathbf{F}_p$ -Hopf algebra  $\Lambda_{\mathbf{F}_p}^*(A)$ . Thus we get that  $\text{com}'_*(x) = 0$ . Since  $\Delta$  is also iterated comultiplication (corresponding to the partition  $(1, \dots, 1)$ ), then  $\Delta$  factors through  $\text{com}'_*$  and we obtain  $\ker(\text{com}'_*) \subset \ker(\Delta_*)$  concluding the proof. We note that in fact  $\ker(\text{com}'_*) = \ker(\Delta_*)$ , and we have introduced  $\Delta$  only in order to simplify the statement of the theorem, since  $\text{com}$  depends on  $j$  in a more complicated way.

For the case  $p = 2$  we proceed analogously. The only difference is the different description of the homology of an abelian group which does not affect our arguments. □

### 3. REMARKS AND EXAMPLES

This paper was motivated by the following example. We consider a split extension of finite  $\mathbf{F}_p$ -algebras

$$(6) \quad \mathbf{F}_p \rightarrow \mathbf{F}_p[x]/x^2 \rightarrow \mathbf{F}_p.$$

This extension induces a split group extension

$$(7) \quad 0 \rightarrow M(J) \rightarrow GL(R) \rightarrow GL(S) \rightarrow 1$$

where  $GL$  is the colimit of general linear groups,  $M$  is the colimit of additive groups of matrices and  $GL(\mathbf{F}_p)$  acts on  $M(\mathbf{F}_p)$  by conjugation (of course, a group extension exists already at the level of  $GL_n$  and  $M_n$ ). It was shown by Goodwillie that for any split extension of rings  $J \rightarrow R \rightarrow S$ , where  $J$  is a free  $S$ -bimodule regarded as an ideal with trivial multiplication, that  $\Lambda^*(M(J) \otimes \mathbf{Q})_{GL(S)} = H_*(F, \mathbf{Q})$  where  $F$  is the homotopy fiber of the induced map  $BGL^+(R) \rightarrow BGL^+(S)$  ([Go], p. 395). This result awakened my interest to the space  $\Lambda^*(A)_G$ . For example, if Goodwillie's theorem was also true with coefficients in  $\mathbf{F}_p$ , then thanks to  $\tilde{H}_*(GL(\mathbf{F}_p), \mathbf{F}_p) = 0$  ([Qu1]) we would obtain

$$H_*(GL(\mathbf{F}_p[x]/x^2), \mathbf{F}_p) = \Lambda^*(M(\mathbf{F}_p), \mathbf{F}_p)_{GL(\mathbf{F}_p)}.$$

It has been known since the early eighties (see e.g. [EF]) that this equality cannot hold because  $H_*(GL(\mathbf{F}_p[x]/x^2), \mathbf{F}_p)$  is too big, but initially I conjectured that  $\Lambda^*(M(\mathbf{F}_p), \mathbf{F}_p)_{GL(\mathbf{F}_p)}$  embeds into  $H_*(GL(\mathbf{F}_p[x]/x^2), \mathbf{F}_p)$  through the edge homomorphism in the sequence (2) associated to the extension (7). This hope was destroyed by results of [HM]. Hesselholt and Madsen have computed  $K_*(\mathbf{F}_p[x]/x^2)$ , but since  $BGL^+(\mathbf{F}_p[x]/x^2)_p^\wedge$  is a generalized Eilenberg–Mac Lane spectrum, it also determines  $H_*(GL(\mathbf{F}_p[x]/x^2), \mathbf{F}_p)$ . In particular, their formulas give

$$H_2(GL(\mathbf{F}_2[x]/x^2), \mathbf{F}_2) = \mathbf{F}_2,$$

but it is easy to see that  $H_0(GL(\mathbf{F}_2), D^2(M(\mathbf{F}_2))) = \mathbf{F}_2^2$ . This shows that the spectral sequence (3) corresponding to the extension  $\mathbf{F}_2 \rightarrow \mathbf{F}_2[x]/x^2 \rightarrow \mathbf{F}_2$  must have a nontrivial differential arriving at  $H_0(GL(\mathbf{F}_2), D^2(M(\mathbf{F}_2)))$ . A similar example may be also constructed for  $p = 3$ . Thus the restriction to  $(H_i)_{reg}$  in Theorem 1 is necessary.

We look more closely at the groups in Theorem 1, and focus on  $H_0(G, \Lambda^*(A))_{reg}$  (for  $p$  odd) which, as we have seen, appears in another context but is also more computable than  $H_1(G, \Lambda^*(A))_{reg}$ . In general, the process of computing  $H_0(G, \Lambda^j(A))_{reg}$  divides into two steps. The first requires knowledge not only about  $H_0(G, A)$  but also about the whole representation  $G \rightarrow Aut(A)$  to determine  $H_0(G, A^{\otimes j})$ . The second is to describe the action of the group  $\Sigma_j$  on  $H_0(G, A^{\otimes j})$  induced by permutation of factors in  $A^{\otimes j}$ . If one completes this program, in order to obtain a formula for  $H_0(G, \Lambda^j(A))_{reg}$  it suffices to observe that it may be identified with the image of the endomorphism  $Alt_*$  of  $H_0(G, A^{\otimes j})$  given by the antisymmetrization formula

$$(8) \quad Alt_*(x) = \sum_{\sigma \in \Sigma_j} sgn(\sigma) \sigma.x$$

(an analogous fact is not true for  $p = 2$  because  $D^j(A)$  is not an image of  $A^{\otimes j}$ ).

To illustrate this algorithm we apply it to extension (7). By the First Fundamental Theorem of (co)Invariant Theory [dCP] we have

$$H_0(GL(\mathbf{F}_p), M(\mathbf{F}_p)^{\otimes j}) = \mathbf{F}_p[\Sigma_j]$$

and the action of the symmetric group on the group algebra is given by the formula  $\sigma.e_\tau = e_{\sigma\tau\sigma^{-1}}$ . Now we should describe the image of the antisymmetrization map (8). Let us take  $e_\tau \in \mathbf{F}_p[\Sigma_j]$  and consider two different cases: if the centralizer of  $\tau$  contains an odd permutation, and if it does not. In the first case we have  $Alt_*(e_\tau) = 0$  so we focus on the case when the centralizer consists of only even

permutations. Then choosing representatives for  $\Sigma_j/Centr(\tau)$  we may write

$$Alt_*(e_\tau) = |Centr(\tau)| \cdot \sum_{\tau' \in \Sigma_j/Centr(\tau)} sgn(\tau') \cdot e_{\tau'\tau}.$$

From this formula the following consequences may be immediately derived:  $Alt_*(e_\tau)$  depends only on the conjugacy class of  $\tau$ , it is nontrivial if  $Centr(\tau)$  contains no odd permutation and is of order prime to  $p$ , elements in different conjugacy classes have images linearly independent. Using elementary combinatorics of the symmetric group we may translate these conditions into the language of partitions of  $j$ . The result is

$$\dim(H_0(G, \Lambda^j(A))_{reg}) = \{\text{the number of partitions of } j \text{ into} \\ \text{different odd numbers prime to } p\}.$$

We point out that the last requirement is nothing but the condition for regularity of a conjugacy class in the sense of representation theory. This explains our notation for  $(H_*)_{reg}$ .

We finish by making one disappointing remark concerning the group  $H_1(GL(\mathbf{F}_p), \Lambda^*(M(\mathbf{F}_p)))_{reg}$ . Namely, in contrast to  $H_0(GL(\mathbf{F}_p), \Lambda^j(M(\mathbf{F}_p)))_{reg}$ , it quite easily follows from [Be1] and [Be2] that  $H_1(GL(\mathbf{F}_p), \Lambda^j(M(J))) = 0$  for  $p > 2$  and  $j < p$ .

#### REFERENCES

- [Be1] S. Betley, *Calculations in THH-Theory*, J. Algebra **180**, (1996), 445–458. MR **97a**:19002
- [Be2] S. Betley, *Stable K-Theory of Finite Fields*, K-Theory **17** (1999), 103–111. MR **2000d**:18006
- [dCP] C. de Concini, C. Procesi, *A characteristic free approach to invariant theory*, Advances in Math. **21** (1976), no. 3, 330–354. MR **54**:10305
- [EF] L. Evens, E. Friedlander, *On  $K_*(\mathbb{Z}/p^2)$  and related homology groups*, Trans. Amer. Math. Soc. **270** (1982), 1–46. MR **83j**:18013
- [Go] T. Goodwillie, *Relative algebraic K-theory and cyclic homology*, Ann. Math. **124** (1986), 347–402. MR **88b**:18008
- [HM] L. Hesselholt, I. Madsen, *On the K-theory of finite algebras over Witt vectors of perfect fields*, Topology **36** (1997), 29–101. MR **97i**:19002
- [Qu1] D. Quillen, *On the cohomology and K-theory of the general linear groups over a finite field*, Ann. Math. **96** (1972), 552–586. MR **47**:3565
- [Qu2] D. Quillen, *Characteristic classes of representations*, Lect. Notes in Math. **551**, (1976) (189–216). MR **58**:28215
- [Sa] C.-H. Sah, *Cohomology of split extensions*, J. Algebra **29** (1974), 255–302. MR **52**:14083

INSTYTUT MATEMATYKI, UNIVERSITY OF WARSAW, UL. BANACHA 2, 02-097 WARSAW, POLAND  
E-mail address: mchal@mimuw.edu.pl