

ON AN ADJOINT FUNCTOR TO THE THOM FUNCTOR

YULI B. RUDYAK

(Communicated by Ralph Cohen)

ABSTRACT. We construct a right adjoint functor to the Thom functor, i.e., to the functor which assigns the Thom space $T\xi$ to a vector bundle ξ .

INTRODUCTION

Let τ denote the functor which assigns the Thom space $T\xi$ to a vector bundle ξ , and similarly for maps. The goal of this paper is to construct the right adjoint functor λ to the functor τ .

To motivate this result, I remark that it is always nice to know whether a functor admits an adjoint one. However, here we have a more interesting motivation. Namely, it is useful to know when a space is the Thom space of a certain vector bundle (spherical fibration). For example, de-Thomification plays an important role in the theory of immersion of manifolds; see [BP], [C] and the survey [L]. In fact, Brown and Peterson [BP] de-Thomify a space, while Cohen [C] de-Thomifies a map. However, these de-Thomifications are very ad hoc. So, it is reasonable to want a de-Thomification machine, like the de-looping machine of May [M] or Boardman and Vogt [BV]. The following observation of Beck [B] plays the crucial role in the de-looping theory. The suspension functor S is the left adjoint to the loop functor Ω , and so there is a monad $M := \Omega S$. Clearly, every loop space is a space over M . Conversely, if a space X is a space over M , then, using the simplicial resolution of the M -space X , one can provide a de-looping of X “at the simplicial level”, and then certain additional arguments enable us to lift this “simplicial de-looping” to the geometric level; see [B], [M].

Here we have a dual situation. As usual, the functor $C := \tau\lambda$ is a comonad, and every Thom space is a space over C . Conversely, if X is a space over C then, dually to what we said above, one can take the cosimplicial resolution of X and provide a de-Thomification of X “at the cosimplicial level”. However, in order to do the next step, a lifting to the geometrical level, one must prove that the Thomification commutes with the functor Tot, and this problem looks quite complicated; cf. [Bo].

Summarizing, one can consider this paper as a first step in an attack on the de-Thomification problem.

Notice that the above arguments enable us to prove that a certain space is not a Thom space: it suffices to check that it is not a space over the comonad C . For

Received by the editors April 2, 1999 and, in revised form, March 26, 2000.
2000 *Mathematics Subject Classification*. Primary 55R25; Secondary 18A40.

example, we have (implicitly) used these arguments in [R1] in order to prove that the spectra k and kO are not Thom spectra.

THE CASE OF NON-ORIENTABLE BUNDLES

Let O_n be the group of orthogonal transformations of the Euclidean space \mathbb{R}^n , let BO_n denote its classifying space, and let γ denote the universal n -dimensional vector bundle over BO_n . Given a locally trivial bundle ξ with the fiber \mathbb{R}^n and structure group O_n , let $T\xi$ denote the Thom space of ξ , i.e., $T\xi := D(\xi)/S(\xi)$, where $D(\xi)$ is the total space of the unit disc bundle and $S(\xi)$ is the total space of the unit sphere subbundle of $D(\xi)$. We regard $T\xi$ as a pointed space with the base point given by $S(\xi)$.

Let \mathcal{K} be the category whose objects are maps $f : B \rightarrow BO_n$, where B is a connected space and f is a map such that

$$\pi_1(B) \xrightarrow{f_*} \pi_1(BO_n) = \mathbb{Z}/2$$

is an epimorphism, and whose morphisms are commutative diagrams

$$\begin{array}{ccc} B & \xrightarrow{\varphi} & C \\ f \downarrow & & \downarrow g \\ BO_n & \xlongequal{\quad} & BO_n, \end{array}$$

where f and g are objects of \mathcal{K} . Let \mathcal{S} be the category whose objects are pointed spaces X with $\pi_i(X) = 0$ for $i < n$ and $\pi_n(X) = \mathbb{Z}/2$, and whose morphisms are maps $f : X \rightarrow Y$ such that $f_* : \pi_n(X) \rightarrow \pi_n(Y)$ is an isomorphism.

Let $\tau : \mathcal{K} \rightarrow \mathcal{S}$ be the Thom functor which assigns the object $\tau f := T(f^*\gamma) \in \mathcal{S}$ to the object $f : X \rightarrow BO_n$ of \mathcal{K} .

Theorem 1. *The functor τ admits a right adjoint functor $\lambda : \mathcal{S} \rightarrow \mathcal{K}$.*

Proof. We construct λ as follows. Choose any $X \in \mathcal{S}$. Given an integer k , let $\Omega_k^n X$ be the component of $\Omega^n X$ corresponding to $k \in \pi_n(X) = \pi_0(\Omega^n X) = \mathbb{Z}/2$. The standard O_n -action on \mathbb{R}^n yields the obvious O_n -action on S^n , which, in turn, induces a (right) O_n -action on $\Omega^n X = (X, *)^{(S^n, *)}$, and it is clear that every component $\Omega_k^n X, k = 0, 1$, is O_n -invariant. Convert the right O_n -action on $\Omega_1^n X$ into a left O_n -action by setting $ga = ag^{-1}, g \in O_n, a \in \Omega_1^n X$. Consider the locally trivial bundle

$$p : EO_n \times_{O_n} \Omega_1^n X \rightarrow BO_n$$

which is associated with the universal principal O_n -bundle $\Gamma := \{EO_n \rightarrow BO_n\}$; cf. [PS]. We define λX to be the map p . The λ -action on morphisms is clear.

We prove that λ is right adjoint to τ , i.e., that $\mathcal{K}(f, \lambda X) = \mathcal{S}(T(f^*\gamma), X)$ for every $f : B \rightarrow BO_n$; cf. [R2]. Indeed, consider the principal O_n -bundle

$$f^*\Gamma = \{q : E \rightarrow B\},$$

and let ξ be the $\Omega_1^n X$ -bundle associated with $f^*\Gamma$, i.e.

$$\xi = \{E \times_{O_n} \Omega_1^n X \rightarrow B\}.$$

Then ξ is induced by f from the bundle $\lambda X = \{p : EO_n \times_{O_n} \Omega_1^n X \rightarrow BO_n\}$. So, $\mathcal{K}(f, \lambda X) = \text{Sec } \xi$, where $\text{Sec } \xi$ denotes the set of all sections of ξ .

For every $b \in B$ choose any O_n -equivariant map $i_b : O_n \rightarrow E$ with $qi_b(O_n) = b$. We have (the first equality can be found e.g. in [H])

$$\begin{aligned} \text{Sec } \xi &= \{O_n\text{-equivariant maps } E \rightarrow \Omega_1^n X\} \\ &= \{O_n\text{-equivariant maps } f : E \rightarrow (X, *)^{(S^n, *)} \\ &\quad \text{such that } f(x) \in \Omega_1^n X \text{ for every } a \in E\} \\ &= \{\text{maps } f : E \times_{O_n} (S^n, *) \rightarrow (X, *) \text{ such that the map} \\ &\quad (S^n, *) = O_n \times_{O_n} (S^n, *) \xrightarrow{i_b} E \times_{O_n} (S^n, *) \xrightarrow{f} (X, *) \\ &\quad \text{belongs to } \Omega_1^n X \text{ for every } b\} \\ &= \mathcal{S}(T(f^* \gamma), X). \end{aligned}$$

□

THE CASE OF ORIENTABLE BUNDLES

Let BSO_n be the classifying space for the connected component SO_n of O_n . Let \mathcal{K}' be the category whose objects are maps $f : B \rightarrow BSO_n$, where B is a connected space, and whose morphisms are commutative diagrams

$$\begin{array}{ccc} B & \xrightarrow{\varphi} & C \\ f \downarrow & & \downarrow g \\ BSO_n & \xlongequal{\quad} & BSO_n, \end{array}$$

where f and g are objects of \mathcal{K}' . Let \mathcal{S}' be the category whose objects are pairs (X, a_X) , where X is a pointed space with $\pi_i(X) = 0$ for $i < n$ and a_X is a generator (one of two) of $\pi_n(X) = \mathbb{Z}$, and whose morphisms are maps $\varphi : X \rightarrow Y$ with $\varphi_*(a_X) = a_Y$.

Let γ' be the universal oriented n -dimensional vector bundle over BSO_n . There is a unique element $a \in \pi_n(T\gamma') = \mathbb{Z}$ such that $\langle u, h(a) \rangle = 1$, where $u \in H^n(T\gamma') = \mathbb{Z}$ is the orientation of γ' , $h : \pi_n(T\gamma') \rightarrow H_n(T\gamma')$ is the Hurewicz homomorphism and $\langle -, - \rangle$ is the Kronecker pairing.

Given an object $f : X \rightarrow BSO_n$ of \mathcal{K}' , we have the canonical map $F : T(f^* \gamma') \rightarrow T\gamma'$, and $F_* : \mathbb{Z} = \pi_n(T(f^* \gamma')) \rightarrow \pi_n(T\gamma) = \mathbb{Z}$ is an isomorphism. Now define the Thom functor $\tau' : \mathcal{K}' \rightarrow \mathcal{S}'$ by setting $\tau' f = (T(f^* \gamma), (F_*)^{-1}(a))$.

Theorem 2. *The functor τ' admits a right adjoint functor $\lambda' : \mathcal{S}' \rightarrow \mathcal{K}'$.*

Proof. Given an object (X, a_X) of \mathcal{K}' , consider the isomorphism $\pi_n(X) \cong \pi_0(\Omega^n X)$, and let $\Omega_1^n X$ be the component of $\Omega^n X$ which corresponds to a_X . As in §1, we have the left SO_n -action on $\Omega^n X$, and clearly the component $\Omega_1^n X$ is invariant under the SO_n -action on $\Omega^n X$. We construct a fibre bundle $p : ESO_n \times_{SO_n} \Omega_1^n X \rightarrow BSO_n$, and we define $\lambda'(X, a_X) := p$. Now the proof can be completed similarly to that of Theorem 1. □

REFERENCES

[B] J. Beck, *On H-spaces and infinite loop spaces*, Category Theory, Homotopy Theory and Their Applications III. Lecture Notes in Mathematics **99**, Springer, Berlin Heidelberg New York, 1969., pp. 139-153. MR **40**:2079

- [BV] J.M. Boardman, R.M. Vogt, *Homotopy invariant algebraic structures on topological spaces*, Lecture Notes in Mathematics **347**, Springer, Berlin Heidelberg New York, 1973. MR **54**:8623a
- [Bo] A.K. Bousfield, *On the homology spectral sequence of a cosimplicial space*, Amer. J. Math. **109** (1987), 361–394. MR **88j**:55017
- [BP] E.H. Brown, F.P. Peterson, *A universal space for normal bundles of n -manifolds*, Comment. Math. Helv. **54** (1979), 405–430. MR **80k**:57055
- [C] R. Cohen, *The immersion conjecture for differentiable manifolds.*, Ann of Math **122** (1985), 237–328. MR **86m**:57030
- [H] D. Husemoller, *Fibre bundles.* – McGraw-Hill, New York, 1966. MR **37**:4821
- [L] J. Lannes, *La conjecture des immersions (d'après R.L. Cohen, E.H. Brown, F.P. Peterson et al.)*, Bourbaki Seminar **1981/1982**, Soc. Math. France, Paris, 1982, pp. 331–346. MR **84i**:57020
- [M] J.P. May, *The geometry of iterated loop spaces*, Lecture Notes in Mathematics **271**, Springer, Berlin Heidelberg New York, 1972. MR **54**:8623b
- [PS] R. Patterson, R. Stong, *Orientability of bundles*, Duke Math. J. **39** (197), 619–622. MR **47**:4248
- [R1] Yu. Rudyak, *The spectra k and kO are not Thom spectra*, Group Representations: Cohomology, Group Actions and Topology, Proc. Symp. Pure Math. **63**, Amer. Math. Soc, Providence, RI, 1998, pp. 475–483. MR **98m**:55007
- [R2] Yu. Rudyak, *On Thom Spectra, Orientability and Cobordism*, Springer, Berlin, 1998. MR **99f**:55001

MATHEMATISCHES INSTITUT, UNIVERSITÄT HEIDELBERG, IM NEUENHEIMER FELD 288, 69120 HEIDELBERG, GERMANY

E-mail address: `rudyak@mathi.uni-heidelberg.de`

Current address: Department of Mathematics, University of Florida, Gainesville, Florida 32611-8105

E-mail address: `rudyak@math.ufl.edu`