

POINTWISE DIMENSIONS AND RÉNYI DIMENSIONS

RYSZARD RUDNICKI

(Communicated by David Preiss)

ABSTRACT. We prove that the local lower and upper pointwise dimensions of a probability measure μ are bounded from below by the lower generalized dimension $\underline{D}_\mu(q)$ for $q > 1$ and from above by the upper generalized dimension $\bar{D}_\mu(q)$ for $q < 1$.

Let μ be a probability Borel measure on a metric space (M, ρ) and let X be the support of μ . For any $q \neq 1$ we define the lower generalized dimensions $\underline{D}_\mu(q)$ by

$$\underline{D}_\mu(q) = \underline{\lim}_{\varepsilon \rightarrow 0^+} \frac{\log \int_X \mu(B(x, \varepsilon))^{q-1} \mu(dx)}{(q-1) \log \varepsilon},$$

where $B(x, \varepsilon)$ is the open ball with centre $x \in X$ and radius $\varepsilon > 0$. Replacing in the above definition the lower limit by the upper one we define the upper generalized dimensions $\bar{D}_\mu(q)$. The generalized dimensions $\underline{D}_\mu(q)$ and $\bar{D}_\mu(q)$ introduced by Hentschel and Procaccia [2] are decreasing functions of q (see [1]). The generalized dimensions are frequently called Rényi dimensions. They are intensively studied in mathematical and physical literature concerning dynamical systems (see [4] and ref. therein). The lower and upper pointwise dimensions $\underline{d}_\mu(x)$ and $\bar{d}_\mu(x)$ are given by

$$\underline{d}_\mu(x) = \underline{\lim}_{\varepsilon \rightarrow 0^+} \frac{\log \mu(B(x, \varepsilon))}{\log \varepsilon}, \quad \bar{d}_\mu(x) = \overline{\lim}_{\varepsilon \rightarrow 0^+} \frac{\log \mu(B(x, \varepsilon))}{\log \varepsilon}.$$

The following theorem was formulated in another but equivalent way in [3] in the case when μ is a Borel measure on \mathbb{R}^d with bounded support. The proof given in [3] is based on the Vitali Covering Lemma and cannot be adapted to the general case.

Theorem. *If $q > 1$, then $\underline{D}_\mu(q) \leq \underline{d}_\mu(x)$ for μ -a.e. x . If $q < 1$, then $\bar{D}_\mu(q) \geq \bar{d}_\mu(x)$ for μ -a.e. x .*

Proof. To prove the first part of the Theorem we assume, contrary to our claim, that there exist $q > 1$ and $\lambda_0 > 0$ such that $\underline{D}_\mu(q) > \lambda_0$ and $\mu\{x : \underline{d}_\mu(x) < \lambda_0\} > 0$. Let λ, λ_1 be constants such that $\lambda_0 < \lambda < \lambda_1 < \underline{D}_\mu(q)$. Let $A = \{x : \underline{d}_\mu(x) < \lambda_0\}$

Received by the editors January 15, 2001.

2000 *Mathematics Subject Classification.* Primary 28A80; Secondary 28A78.

Key words and phrases. Pointwise dimension, generalized (Rényi) dimension, measure.

This research was partially supported by the State Committee for Scientific Research (Poland) Grant No. 2 P03A 010 161.

and $\mathcal{E}(x) = \{\varepsilon \in (0, 1) : \mu(B(x, \varepsilon)) > \varepsilon^{\lambda_0}\}$ for $x \in X$. Then from the definition of the set A it follows immediately that $\inf \mathcal{E}(x) = 0$ for $x \in A$. Let $\bar{\varepsilon} \in \mathcal{E}(x)$. Then

$$(1) \quad \mu(B(x, \varepsilon)) > \varepsilon^\lambda \quad \text{for } \varepsilon \in [\bar{\varepsilon}, \bar{\varepsilon}^{\lambda_0/\lambda}].$$

Set $A(\varepsilon) = \{x : \mu(B(x, \varepsilon)) > \varepsilon^\lambda\}$ and $\varepsilon_n = 2^{-(\lambda/\lambda_0)^n}$ for $n \in \mathbb{N}$. Observe that

$$(2) \quad A \subset \bigcap_{m=1}^{\infty} \bigcup_{n=m}^{\infty} A(\varepsilon_n).$$

Indeed, fix an $x \in A$. Then for every $m \in \mathbb{N}$ there exist an $\bar{\varepsilon} \in \mathcal{E}(x)$ and $n \geq m$ such that $\varepsilon_{n+1} \leq \bar{\varepsilon} \leq \varepsilon_n$. From (1) it follows that $x \in A(\varepsilon_n)$ and condition (2) holds. Since $\mu(A) > 0$, from (2) we obtain

$$(3) \quad \sum_{n=1}^{\infty} \mu(A(\varepsilon_n)) = \infty.$$

Since $\underline{D}_\mu(q) > \lambda_1$, there exists $r > 0$ such that

$$(4) \quad \int_X \mu(B(x, \varepsilon))^{q-1} \mu(dx) \leq \varepsilon^{\lambda_1(q-1)} \quad \text{for } \varepsilon \in (0, r).$$

From (1) and (4) we obtain $\mu(A(\varepsilon))\varepsilon^{\lambda(q-1)} \leq \varepsilon^{\lambda_1(q-1)}$ for $\varepsilon \in (0, r)$, which implies that $\mu(A(\varepsilon)) \leq \varepsilon^{(\lambda_1-\lambda)(q-1)}$. Since $\sum_{n=1}^{\infty} \varepsilon_n^{(\lambda_1-\lambda)(q-1)} < \infty$, the series (3) is convergent, a contradiction.

In the same manner we can prove the second part of the Theorem. In this case we assume, contrary to our claim, that there are some constants $\lambda_1, \lambda, \lambda_0$ and $q < 1$ such that $\overline{D}_\mu(q) < \lambda_1 < \lambda < \lambda_0$ and $\mu\{x : \bar{d}_\mu(x) > \lambda_0\} > 0$. Replacing in the definitions of the sets of $A, \mathcal{E}(x), A(\varepsilon)$ the inequalities by opposite ones we can check that $\mu(B(x, \varepsilon)) < \varepsilon^\lambda$ for $\varepsilon \in [\bar{\varepsilon}^{\lambda_0/\lambda}, \bar{\varepsilon}]$. Also, (2) holds if $\varepsilon_n = 2^{-(\lambda_0/\lambda)^n}$ for $n \in \mathbb{N}$. The rest of the proof is exactly the same as the proof of the first part. \square

Remark. In the statement of the Theorem inequalities cannot be replaced by equalities. Let $f = \sum_{n=1}^{\infty} 2^{-n+n^2} f_n$, where f_n is the characteristic function of the interval $[2^{-n}, 2^{-n} + 2^{-n^2}]$. If μ is the probability measure on the interval $[0, 1]$ with the density f , then $\underline{D}_\mu(q) = \overline{D}_\mu(q) = 0$ for all $q > 1$ but $\underline{d}_\mu(x) = \bar{d}_\mu(x) = 1$ for all $x \in \text{supp } \mu$. If $\mu = \sum_{n=1}^{\infty} a_n \delta_{x_n}$, where $a_n = cn^{-1} \log^{-2}(n+1)$ and $\sum_{n=1}^{\infty} a_n = 1$, $x_1 = 1$, $x_n = 1 - \sum_{k=1}^{n-1} a_k$ for $n \geq 2$ and δ_{x_n} is the Dirac measure at x_n , then $\overline{D}_\mu(q) \geq \underline{D}_\mu(q) \geq 1$ for all $q < 1$ but $\underline{d}_\mu(x) = \bar{d}_\mu(x) = 0$ for all $x \in \text{supp } \mu$.

REFERENCES

1. C.D. Cutler, *Some results on the behavior and estimation of the fractal dimensions of distribution on attractors*, J. Statist. Phys. **62** (1991), 651–708. MR **92b**:58137
2. H. Hentschel and I. Procaccia, *The infinite number of generalized dimensions of fractals and strange attractors*, Phys. D, **8** (1983), 435–444. MR **85a**:58064
3. S-N. Ngai, *A dimension result arising from the L^q -spectrum of a measure*, Proc. Amer. Math. Soc. **125** (1997), 2943–2951. MR **97m**:28007
4. Y.B. Pesin, *Dimension Theory in Dynamical Systems. Contemporary views and applications*, University of Chicago Press, Chicago, 1997. MR **99b**:58003

INSTITUTE OF MATHEMATICS, POLISH ACADEMY OF SCIENCES AND INSTITUTE OF MATHEMATICS,
SILESIAN UNIVERSITY, BANKOWA 14, 40-007 KATOWICE, POLAND
E-mail address: rudnicki@us.edu.pl