

L^1 CONVERGENCE OF THE RECONSTRUCTION FORMULA FOR THE POTENTIAL FUNCTION

YA-TING CHEN, Y. H. CHENG, C. K. LAW, AND J. TSAY

(Communicated by Carmen C. Chicone)

ABSTRACT. It is known that the potential function of the Sturm-Liouville problem can be reconstructed from the nodal data by a pointwise limit. We show that this convergence is in fact L^1 .

1. INTRODUCTION

Consider the Sturm-Liouville problem

$$-y'' + q(x)y = \lambda y$$

such that

$$\begin{cases} y(0) \cos \alpha + y'(0) \sin \alpha = 0, \\ y(1) \cos \beta + y'(1) \sin \beta = 0, \end{cases}$$

where $q \in L^1(0, 1)$, and $\alpha, \beta \in [0, \pi)$.

In 1988, J.R. McLaughlin [8] showed that knowledge of the nodal set of the eigenfunctions alone can determine the potential function of the Sturm-Liouville problem up to a constant. This is the so-called inverse nodal problem [1]. C.F. Yang [13] showed that this uniqueness result is valid for any $q \in L^1(0, 1)$. Independently C.L. Shen [10] had similar ideas for the density function ρ of the string equation $y'' + \lambda\rho y = 0$. See also [1, 11, 12]. The uniqueness result holds for the situation that ρ is only assumed to be of bounded variation [3].

The inverse nodal problem in two dimension is much more difficult. Hald and McLaughlin [2] proved the uniqueness result for the potential function of the Schrödinger operator defined on a rectangular domain in \mathbf{R}^2 . Some results for density function of string equation on a membrane can be found in [9, 7].

In this note we concern ourselves with the reconstruction of the potential function of the one-dimensional Sturm-Liouville problem. Law-Shen-Yang [4], improving a result of X.F. Yang [13], gave a reconstruction formula for the potential function q .

Let λ_n be the n -th eigenvalue, $s_n = \sqrt{\lambda_n}$ and $x_i^{(n)}$ be the i -th nodal point of the n -th eigenfunction y_n . In other words, $y_n(x_i^{(n)}) = 0$, $i = 1, 2, \dots, n-1$. Let $I_i^{(n)} = (x_i^{(n)}, x_{i+1}^{(n)})$, and $l_i^{(n)}$ be the nodal length where $l_i^{(n)} = |I_i^{(n)}| = x_{i+1}^{(n)} - x_i^{(n)}$.

Received by the editors November 21, 2000 and, in revised form, February 20, 2001.

2000 *Mathematics Subject Classification*. Primary 34A55; Secondary 34B24.

Key words and phrases. Inverse nodal problem, potential function, reconstruction formula.

Define $x_0^{(n)} = 0, x_n^{(n)} = 1$. We also define the function $j_n(x)$ to be the largest index j such that $0 \leq x_j^{(n)} \leq x$. Thus, $j = j_n(x)$ if and only if $x \in [x_j^{(n)}, x_{j+1}^{(n)})$.

Theorem 1.1 ([4]). *The potential function $q \in L^1(0, 1)$ satisfies*

$$q(x) = \lim_{n \rightarrow \infty} 2s_n^2 \left(\frac{s_n l_j^{(n)}}{\pi} - 1 \right)$$

for a.e. $x \in (0, 1)$, with $j = j_n(x)$.

Note that Theorem 1.1, with the asymptotic expression for s_n (see [5, Lemma 2] and [6, Lemma 2.2]) implies that $q(x) = \lim_{n \rightarrow \infty} F_n(x)$, where F_n is determined only by the nodal data and the constant $\int_0^1 q$:

(a) If $\alpha = \beta = 0$ or $\alpha, \beta > 0$, then

$$F_n(x) = 2n^2 \pi^2 \left\{ n l_j^{(n)} - 1 + \frac{l_j^{(n)}}{n \pi^2} \left(\frac{1}{2} \int_0^1 q - \operatorname{scot} \alpha + \operatorname{scot} \beta \right) \right\}.$$

(b) If $\alpha = 0 < \beta$ or $\beta = 0 < \alpha$, then

$$F_n(x) = 2(n - \frac{1}{2})^2 \pi^2 \left\{ (n - \frac{1}{2}) l_j^{(n)} - 1 + \frac{l_j^{(n)}}{n \pi^2} \left(\frac{1}{2} \int_0^1 q - \operatorname{scot} \alpha + \operatorname{scot} \beta \right) \right\}.$$

Here, $\operatorname{scot} \gamma = 0$ if $\gamma = 0$; $\operatorname{scot} \gamma = \cot \gamma$ otherwise.

Since q is in L^1 , one may ask whether the convergence is in fact L^1 . The answer is affirmative.

Theorem 1.2. *F_n converges to q in L^1 .*

2. PROOF

Lemma 2.1 ([5]). *Let $q \in L^1$. Then*

$$(2.1) \quad l_i^{(n)} = \frac{\pi}{s_n} + \frac{1}{2s_n^2} \int_{x_i^{(n)}}^{x_{i+1}^{(n)}} (1 + \alpha_0 \cos(2s_n y)) q(y) dy + o\left(\frac{1}{s_n^3}\right),$$

where $\alpha_0 = 1$ if $\alpha > 0$, $\alpha_0 = -1$ if $\alpha = 0$.

In the above lemma, the order estimate is independent of i . As a result,

$$\frac{s_n l_i^{(n)}}{\pi} = 1 + O\left(\frac{1}{s_n}\right) = 1 + O\left(\frac{1}{n}\right).$$

Lemma 2.2. *Suppose the sequence $f_k \in C[0, 1]$ converges to f in L^1 . For any $\epsilon > 0$, then with $j = j_n(x)$,*

$$\left\| \frac{s_n}{\pi} \int_{x_j^{(n)}}^{x_{j+1}^{(n)}} (f_k(y) - f(y)) dy \right\|_1 < \epsilon$$

for all sufficiently large n and k .

Proof. By Lemma 2.1 and the observation that the integral $\int_{x_j^{(n)}}^{x_{j+1}^{(n)}} (f_k(y) - f(y)) dy$ is constant on any nodal interval $I_j^{(n)}$,

$$\begin{aligned} & \int_0^1 \left| \frac{s_n}{\pi} \int_{x_j^{(n)}}^{x_{j+1}^{(n)}} (f_k(y) - f(y)) dy \right| dx \\ &= \sum_{i=0}^{n-1} \frac{s_n l_i^{(n)}}{\pi} \left| \int_{x_i^{(n)}}^{x_{i+1}^{(n)}} (f_k(y) - f(y)) dy \right| \\ &\leq \sum_{i=0}^{n-1} \left(1 + O\left(\frac{1}{n}\right)\right) \int_{x_i^{(n)}}^{x_{i+1}^{(n)}} |f_k(y) - f(y)| dy \\ &= \left(1 + O\left(\frac{1}{n}\right)\right) \int_0^1 |f_k(y) - f(y)| dy, \end{aligned}$$

and hence converges to 0 as $k \rightarrow \infty$. □

Lemma 2.3. *Suppose $q \in L^1(0, 1)$. Then as $n \rightarrow \infty$, with $j = j_n(x)$,*

$$\left\| \frac{s_n}{\pi} \int_{x_j^{(n)}}^{x_{j+1}^{(n)}} q(y) dy - q(x) \right\|_1 \rightarrow 0.$$

Proof. We first show that the result holds if q is continuous on $[0, 1]$. Let $M = \max_{x \in (0, 1)} |q(x)|$ and $x_0 \in [x, y]$. By intermediate value theorem, there exists $\xi \in (x, y)$ such that

$$\left| \frac{1}{y-x} \int_x^y q - q(x_0) \right| = |q(\xi) - q(x_0)|.$$

Due to the uniform continuity of q there is a $\delta > 0$ such that the difference is small whenever $|y-x| < \delta$. Hence, given $\epsilon > 0$, when n is large enough such that $l_j^{(n)} < \delta$ and $|\frac{s_n l_j^{(n)}}{\pi} - 1| < \min\{\frac{\epsilon}{M}, 1\}$, with $j = j_n(x)$, we have

$$\begin{aligned} & \left| \frac{s_n}{\pi} \int_{x_j^{(n)}}^{x_{j+1}^{(n)}} q(y) dy - q(x) \right| \\ &\leq \left| \frac{s_n(x_{j+1}^{(n)} - x_j^{(n)})}{\pi} \left[\frac{1}{x_{j+1}^{(n)} - x_j^{(n)}} \int_{x_j^{(n)}}^{x_{j+1}^{(n)}} q - q(x) \right] \right| + |q(x)| \left| \frac{s_n l_j^{(n)}}{\pi} - 1 \right| \\ &\leq 2\epsilon. \end{aligned}$$

Therefore, if $q \in C[0, 1]$, then $\left\| \frac{s_n}{\pi} \int_{x_j^{(n)}}^{x_{j+1}^{(n)}} q(y) dy - q(x) \right\|_1$ can be arbitrarily small for all large n .

Since $C[0, 1]$ is dense in $L^1(0, 1)$, for any $q \in L^1(0, 1)$ there exists a sequence $q_k \in C[0, 1]$ that converges to q in $L^1(0, 1)$. Now

$$\begin{aligned} & \int_0^1 \left| \frac{s_n}{\pi} \int_{x_j^{(n)}}^{x_{j+1}^{(n)}} q(y) dy - q(x) \right| dx \\ & \leq \int_0^1 \left| \frac{s_n}{\pi} \int_{x_j^{(n)}}^{x_{j+1}^{(n)}} (q(y) - q_k(y)) dy \right| dx \\ & \quad + \int_0^1 \left| \frac{s_n}{\pi} \int_{x_j^{(n)}}^{x_{j+1}^{(n)}} q_k(y) dy - q_k(x) \right| dx \\ & \quad + \int_0^1 |q_k(x) - q(x)| dx . \end{aligned}$$

For any $\epsilon > 0$, fix k large enough such that the first and last terms are smaller than ϵ . Then for all n large enough, the second term is smaller than 2ϵ by above. Hence, as $n \rightarrow \infty$,

$$\left\| \frac{s_n}{\pi} \int_{x_j^{(n)}}^{x_{j+1}^{(n)}} q(y) dy - q(x) \right\|_1 \rightarrow 0 .$$

□

Proof of Theorem 1.2. Since

$$\left| F_n(x) - 2s_n^2 \left(\frac{s_n l_j^{(n)}}{\pi} - 1 \right) \right| = O\left(\frac{1}{n}\right),$$

by an asymptotic estimate of s_n , it suffices to show that as $n \rightarrow \infty$,

$$\left\| 2s_n^2 \left(\frac{s_n l_j^{(n)}}{\pi} - 1 \right) - q \right\|_1 \rightarrow 0 .$$

By (2.1), we have

$$2s_n^2 \left(\frac{s_n l_j^{(n)}}{\pi} - 1 \right) = \frac{s_n}{\pi} \int_{x_j^{(n)}}^{x_{j+1}^{(n)}} q + \frac{\alpha_0 s_n}{\pi} \int_{x_j^{(n)}}^{x_{j+1}^{(n)}} \cos(2s_n y) q(y) dy + o(1) .$$

Hence we only need to prove that as $n \rightarrow \infty$,

$$\left\| \frac{s_n}{\pi} \int_{x_j^{(n)}}^{x_{j+1}^{(n)}} q - q \right\|_1 \rightarrow 0 ,$$

and

$$(2.2) \quad \left\| \frac{s_n}{\pi} \int_{x_j^{(n)}}^{x_{j+1}^{(n)}} \cos(2s_n y) q(y) dy \right\|_1 \rightarrow 0 .$$

The first limit holds because of Lemma 2.3. By the proof of [4, Theorem 3.2], the sequence of functions

$$h_n(x) = \frac{s_n}{\pi} \int_{x_j^{(n)}}^{x_{j+1}^{(n)}} \cos(2s_n y) q(y) dy$$

converges to 0 for a.e. $x \in (0, 1)$. Furthermore,

$$|h_n(x)| \leq \frac{s_n}{\pi} \int_{x_j^{(n)}}^{x_{j+1}^{(n)}} |q(y)| dy = g_n(x)$$

and

$$\begin{aligned} \int_0^1 g_n(x) dx &= \sum_{i=0}^{n-1} \frac{s_n l_i^{(n)}}{\pi} \int_{x_i^{(n)}}^{x_{i+1}^{(n)}} |q(y)| dy \\ &= \left(1 + O\left(\frac{1}{n}\right)\right) \|q\|_1. \end{aligned}$$

Thus we may apply the Lebesgue dominated convergence theorem to show that (2.2) is valid. The proof is complete. \square

ACKNOWLEDGEMENT

We thank Ngai-Ming Mok of the University of Hong Kong for suggesting the problem. The authors are partially supported by National Science Council, Taiwan, Republic of China.

REFERENCES

- [1] O.H. Hald and J.R. McLaughlin, *Solutions of inverse nodal problems*, Inverse Problems, **5** (1989), 307–347. MR **90c**:34015
- [2] O.H. Hald and J.R. McLaughlin, *Inverse problems: finding the potential from nodal lines*, AMS Memoir, **119**, No. 572 (1996). MR **97d**:35240
- [3] O.H. Hald and J.R. McLaughlin, *Inverse problems: recovery of BV coefficients from nodes*, Inverse Problems, **14** (1998), 245–273. MR **99b**:34028
- [4] C.K. Law, C.L. Shen and C.F. Yang, *The inverse nodal problem on the smoothness of the potential function*, Inverse Problems, **15** (1999), 253–263. MR **2000a**:34020
- [5] C.K. Law, C.L. Shen and C.F. Yang, *Errata: The inverse nodal problem on the smoothness of the potential function*, Inverse Problems, **17** (2001), no. 2, 361–363.
- [6] C.K. Law and J. Tsay, *On the well-posedness of the inverse nodal problem*, Inverse Problems, **17** (2001), no. 5, 1493–1512.
- [7] C.J. Lee and J.R. McLaughlin, *Finding the density for a membrane from nodal lines*, Inverse Problems in Wave Propagation, eds. G. Chavent et al., Springer, New York (1997), 325–345. MR **98h**:35236
- [8] J.R. McLaughlin, *Inverse spectral theory using nodal points as data – a uniqueness result*, J. Diff. Eqns. **73** (1988), 354–362. MR **89f**:34035
- [9] J.R. McLaughlin, *Solving inverse problems with spectral data*, Surveys on Solution Methods for Inverse Problems, eds. D. Colton et al., Springer, New York (2000), 169–194. MR **2001f**:35431
- [10] C.L. Shen, *On the nodal sets of the eigenfunctions of the string equations*, SIAM J. Math. Anal., **19** (1988), 1419–1424. MR **89j**:34035
- [11] C.L. Shen, *On the nodal sets of the eigenfunctions of certain homogeneous and nonhomogeneous membranes*, SIAM J. Math. Anal., **24** (1993) 1277–1282. MR **94i**:35138

- [12] C.L. Shen and T.M. Tsai, *On uniform approximation of the density function of a string equation using eigenvalues and nodal points and some related inverse problems*, *Inverse Problems*, **11** (1995), 1113–1123. MR **96g**:34021
- [13] X.F. Yang, *A solution of the inverse nodal problem*, *Inverse Problems*, **13** (1997), 203–213. MR **98c**:34017

DEPARTMENT OF APPLIED MATHEMATICS, NATIONAL SUN YAT-SEN UNIVERSITY, KAOHSIUNG,
TAIWAN 80424, REPUBLIC OF CHINA
E-mail address: `chenyt@math.nsysu.edu.tw`

DEPARTMENT OF APPLIED MATHEMATICS, NATIONAL SUN YAT-SEN UNIVERSITY, KAOHSIUNG,
TAIWAN 80424, REPUBLIC OF CHINA
E-mail address: `jengyh@math.nsysu.edu.tw`

DEPARTMENT OF APPLIED MATHEMATICS, NATIONAL SUN YAT-SEN UNIVERSITY, KAOHSIUNG,
TAIWAN 80424, REPUBLIC OF CHINA
E-mail address: `law@math.nsysu.edu.tw`

DEPARTMENT OF APPLIED MATHEMATICS, NATIONAL SUN YAT-SEN UNIVERSITY, KAOHSIUNG,
TAIWAN 80424, REPUBLIC OF CHINA
E-mail address: `tsay@math.nsysu.edu.tw`