

FORCING A BOOLEAN ALGEBRA WITH PREDESIGNED AUTOMORPHISM GROUP

TAPANI HYTTINEN AND SAHARON SHELAH

(Communicated by Carl G. Jockusch, Jr.)

ABSTRACT. For suitable groups G we will show that one can add a Boolean algebra B by forcing in such a way that $\text{Aut}(B)$ is almost isomorphic to G . In particular, we will give a positive answer to the following question due to J. Roitman: Is \aleph_ω a possible number of automorphisms of a rich Boolean algebra?

In [Ro], J. Roitman asked, is \aleph_ω a possible number of automorphisms of a rich Boolean algebra? A Boolean algebra is rich if the number of automorphisms is greater than the size of the algebra and possible means consistent with ZFC (since GCH implies trivially that the answer is no). We will answer this question positively. For this we will give a method for adding a Boolean algebra by forcing in such a way that we have a lot of control on the automorphism group of the Boolean algebra. Notice that by [Mo], Theorem 4.3, one cannot hope of giving a positive answer to Roitman's question from the assumption that $2^\omega > \aleph_\omega$.

Partially, our methods are similar to those in [Ro].

We say that a po-set P is κ -Knaster if for all $p_i \in P$, $i < \kappa$, there is a set $Z \subseteq \kappa$ of power κ such that for all $i, j \in Z$, p_i and p_j are compatible.

1. Theorem. *Assume that $\kappa = \kappa^{<\kappa}$ is an infinite cardinal. Let (I, \leq) be a partial order, G_t , $t \in I$, be groups of power $\leq \kappa$ and $\pi_{s,t} : G_t \rightarrow G_s$, $s < t \in I$, be homomorphisms such that*

- (a) I is κ^+ -directed, every pair of elements of I has the least upper bound and for all $t \in I$, $|\{s \in I \mid s \leq t\}| \leq \kappa$,
- (b) for all $r < s < t \in I$, $\pi_{r,t} = \pi_{r,s} \circ \pi_{s,t}$.

Then there is a κ -closed, κ^+ -Knaster po-set P such that in V^P the following holds: There is a Boolean algebra B such that

- (i) B is atomic with $|I| + \kappa$ atoms,
- (ii) $|B| = \kappa^+ + |I|$,
- (iii) if G^* is the inverse limit of $(G_t, \pi_{s,t} \mid s < t \in I)$, then G^* can be embedded into $\text{Aut}(B)$, $|\text{Aut}(B)| \leq \kappa^+ \cdot |I| \cdot |G^*|$ and if for all $t \in I$, G_t is the one element group, then $|\text{Aut}(B)| = \kappa \cdot |I|$.

Received by the editors January 31, 2001 and, in revised form, May 14, 2001.

2000 *Mathematics Subject Classification.* Primary 06E05; Secondary 03E35.

The first author was partially supported by the Academy of Finland, grant 40734, and the Mittag-Leffler Institute.

The research of the second author was supported by the Israel Science Foundation founded by the Israel Academy of Sciences and Humanities. Publ. 756.

Note: We will show that $Aut(B)$ is not very far from G^* . In particular, we will show that for all $f \in Aut(B)$ there is $g \in G^*$ such that excluding a small error, f is the same as the image of g under the embedding.

Proof. For $t \in I$, let $X_t = \{t\} \times G_t \times \kappa$, $X_{\leq t} = \bigcup_{s \leq t} X_s$ and $X = \bigcup_{t \in I} X_t$. We define P to be the set of all tuples $p = (U, T, V, Y) = (U^p, T^p, V^p, Y^p)$ such that

- (1) $U \subseteq X$ is of power $< \kappa$,
- (2) $T \subseteq I$ is of power $< \kappa$,
- (3) $V = (v_t \mid t \in T) = (v_t^p \mid t \in T)$, where each $v_t \subseteq \kappa^+$ is of power $< \kappa$,
- (4) $Y = (y_{t,\alpha} \mid t \in T, \alpha \in v_t) = (y_{t,\alpha}^p \mid t \in T, \alpha \in v_t)$, where $y_{t,\alpha} \subseteq U \cap X_{\leq t}$.

We order P so that $p \leq q$ (q is stronger than p) if

- (5) $U^p \subseteq U^q$,
- (6) $T^p \subseteq T^q$,
- (7) $v_t^p \subseteq v_t^q$ for $t \in T^p$,
- (8) $y_{t,\alpha}^p = y_{t,\alpha}^q \cap U^p$ for $t \in T^p$ and $\alpha \in v_t^p$.

Clearly P is κ -closed and $\kappa^{<\kappa} = \kappa$ implies by the usual Δ -lemma argument that P is κ^+ -Knaster.

Let H be P -generic over V . We let $Y_{t,\alpha} = \bigcup \{y_{t,\alpha}^p \mid p \in H\}$. Below, we will not distinguish these and other objects from their names; it will be clear from the context which we mean.

Let G^* be as in the theorem, i.e. G^* consists of all $g : I \rightarrow \bigcup_{t \in I} G_t$ such that for all $s < t \in I$, $\pi_{s,t}(g(t)) = g(s)$. If $g \in G^*$ and $x = (t, z, \alpha) \in X_t$, then we also write $g(x)$ for $(t, g(t)z, \alpha)$ and define $Y_{t,\alpha,g} = \{g(x) \mid x \in Y_{t,\alpha}\}$.

By a Boolean term $\delta(Y_0, \dots, Y_{n-1})$ taken inside Y we mean a term of the form $\bigcap_{i < n} Z_i$, where each Z_i is either $Y \cap Y_i$ or $Y - Y_i$. If $Z_i = Y \cap Y_i$ we say that Y_i appears in δ positively. A general Boolean term is a finite union of Boolean terms.

1.1. Claim. *Let $\delta(Y_0, \dots, Y_{n-1})$ be a Boolean term taken inside $X_{\leq t}$, $t \in I$, and for $i < n$, $t_i \in I$, $\alpha_i < \kappa^+$ and $g_i \in G^*$. Assume that there is $t' \in I$ such that*

- (a) $t' \leq t$,
- (b) *if Y_i appears in δ positively, then $t' \leq t_i$,*
- (c) *if Y_i appears in δ positively and Y_j does not appear positively in δ , then $(t_i, \alpha_i, g_i(t')) \neq (t_j, \alpha_j, g_j(t'))$.*

Then $\delta(Y_{t_0, \alpha_0, g_0}, \dots, Y_{t_{n-1}, \alpha_{n-1}, g_{n-1}}) \cap X_{\leq t'}$ has power κ . On the other hand, if such a t' does not exist, then $\delta(Y_{t_0, \alpha_0, g_0}, \dots, Y_{t_{n-1}, \alpha_{n-1}, g_{n-1}}) = \emptyset$.

Proof. An easy density argument using the fact that the actions of two distinct elements of a group differ on each element of the group proves Claim 1.1. □

Now we can define the Boolean algebra B . We let B be the subalgebra of $\mathcal{P}(X)$ generated by $[X]^{<\kappa} \cup \{Y_{t,\alpha,g} \mid t \in I, \alpha < \kappa^+, g \in G^*\}$, where $[X]^{<\kappa}$ is the set of all subsets of X of power $< \kappa$.

1.2. Claim. *Items (i) and (ii) from Theorem 1 hold and G^* can be embedded into $Aut(B)$.*

Proof. (i) is clear. For (ii), notice that $Y_{t,\alpha,g}$ is determined by $Y_{t,\alpha}$ and $g(t)$. Finally, it is easy to check that the function which takes $g \in G^*$ to \bar{g} is the required embedding of G^* into $Aut(B)$, where \bar{g} is such that it maps each $A \in B$ to $\{g(z) \mid z \in A\}$. This proves Claim 1.2. □

So we are “left” to prove the second half of (iii) from Theorem 1: For all $0 < m < \omega$ and $t \in I$, let $\delta_i^{t,m}(Y_0, \dots, Y_{m-1})$, $i < 2^m$, list all Boolean terms in m variables taken inside $X_{\leq t}$ so that $\delta_0^{t,m}(Y_0, \dots, Y_{m-1}) = \bigcap_{k < m} (X_{\leq t} - Y_k)$ and for $m = 0$, we let $\delta_0^{t,m} = X_{\leq t}$.

Let $h \in \text{Aut}(B)$. By the definition of B , h is determined by $h \upharpoonright \{\{x\} \mid x \in X\}$. In order to simplify the notation, for $x \in X$, we write $h(x)$ for the unique $z \in X$ such that $h(\{x\}) = \{z\}$. We will frequently consider such an h as if it is a permutation of X .

1.3. Claim. *In V^P , if $h \in \text{Aut}(B)$ and $t \in I$, then there are $u \subseteq X_{\leq t}$, $m < \omega$, for $i < m$, $t_i \in I$, $\alpha_i < \kappa^+$ and $g_i \in G^*$ and for $j < 2^m$, $f_j \in G^*$ such that*

- (i) $|u| < \kappa$,
- (ii) *either for all $j < 2^m$ and $x \in \delta_j^{t,m}(Y_{t_0, \alpha_0, g_0}, \dots, Y_{t_{m-1}, \alpha_{m-1}, g_{m-1}}) - u$, $h(x) = f_j(x)$ or for all $j < 2^m$ and $x \in \delta_j^{t,m}(Y_{t_0, \alpha_0, g_0}, \dots, Y_{t_{m-1}, \alpha_{m-1}, g_{m-1}}) - u$, $h^{-1}(x) = f_j(x)$.*

Proof. For a contradiction, assume that p forces that h and t are a counterexample. For all $x \in X_{\leq t}$, let $(p_{x,i})_{i < \kappa}$ be a maximal antichain such that for all $i < \kappa$, $p_{x,i} \geq p$ or $p_{x,i}$ is incompatible with p and $p_{x,i}$ decides both $h(x)$ and $h^{-1}(x)$. Choose $\alpha^* < \kappa^+$ so that

$$\alpha^* \notin \bigcup \{v_t^{p_{x,i}} \mid x \in X_{\leq t}, i < \kappa\}.$$

Choose $q \geq p$ so that it decides both $h(Y_{t, \alpha^*})$ and $h^{-1}(Y_{t, \alpha^*})$, i.e. it forces $h(Y_{t, \alpha^*})$ to be, say, $(u_0 \cup \delta(Y_{t'_0, \alpha'_0, g'_0}, \dots, Y_{t'_{k-1}, \alpha'_{k-1}, g'_{k-1}})) - u_1$ and $h^{-1}(Y_{t, \alpha^*})$ to be $(u_0^- \cup \delta^-(Y_{t'_0, \alpha'_0, g'_0}, \dots, Y_{t'_{k-1}, \alpha'_{k-1}, g'_{k-1}})) - u_1^-$, where $u_0, u_1, u_1^-, u_2^- \in [X]^{< \kappa}$ and δ and δ^- are general Boolean terms taken inside X . (Notice that $Y_{t, \alpha^*} = Y_{t, \alpha^*, 1}$.) In particular, q is chosen so that it decides values for $g'_i(t'_i)$ and $g'_i(t)$, $i < k$. Clearly we may assume that (q forces that) in $((t'_0, \alpha'_0, g'_0(t'_0)), \dots, (t'_{k-1}, \alpha'_{k-1}, g'_{k-1}(t'_{k-1})))$ there is no repetition, and for some $k^* \leq k$, $\{i < k \mid (t'_i, \alpha'_i) = (t, \alpha^*)\} = k^*$. We write $\bar{Y}_{< k^*}$ for $(Y_{t'_0, \alpha'_0, g'_0}, \dots, Y_{t'_{k^*-1}, \alpha'_{k^*-1}, g'_{k^*-1}})$ and $\bar{Y}_{\geq k^*}$ for $(Y_{t'_{k^*}, \alpha'_{k^*}, g'_{k^*}}, \dots, Y_{t'_{k-1}, \alpha'_{k-1}, g'_{k-1}})$.

If we can find $x \in X_{\leq t} - U^q$ and $q' \geq q$ such that q' forces that for all $i < k^*$, $g'_i(x) \neq h(x)$ and that $h(x) \notin u_0 \cup u_1$, we get an easy contradiction. The contradiction is obtained the same way as we obtain a contradiction after the proof of Subclaim 1.3.1 (we can find $q' \geq q$ so that it forces $h(x) \in h(Y_{t, \alpha^*}) \leftrightarrow h(x) \notin \delta(\bar{Y})$). Similarly, if we can find $x \in X_{\leq t} - U^q$ and $q' \geq q$ such that q' forces that for all $i < k^*$, $g'_i(x) \neq h^{-1}(x)$ and that $h^{-1}(x) \notin u_0^- \cup u_1^-$, we get an easy contradiction. So we assume the following:

(*) For all $x \in X_{\leq t} - U^q$, q forces that $h(x) \in u_0 \cup u_1$ or for some $i < k^*$, $h(x) = g'_i(x)$ and that $h^{-1}(x) \in u_0^- \cup u_1^-$ or for some $i < k^*$, $h^{-1}(x) = g'_i(x)$.

Notice that (*) implies that for all $t' \leq t$, q forces that excluding a small error, h is a permutation of $X_{t'}$.

By (*) above and the choice of p , h and t , and by making q stronger if necessary, we may assume that there are Boolean terms δ' and δ'' taken inside $X_{\leq t}$ such that

- (α) q forces that for all $g \in G^*$ there are κ many $x \in X_{\leq t}$ such that $g(x) \neq h(x) \in \delta'(\bar{Y}_{< k^*}) \cap \delta''(\bar{Y}_{\geq k^*})$,
- (β) q forces that $\delta'(\bar{Y}_{< k^*}) \cap \delta''(\bar{Y}_{\geq k^*}) \neq \emptyset$,
- (γ) every condition forces that $\delta'(\bar{Y}_{< k^*}) \cap \delta''(\bar{Y}_{\leq k^*}) \subseteq \delta(\bar{Y})$.

Instead of item (γ) , we may have $\delta'(\overline{Y}_{<k^*}) \cap \delta''(\overline{Y}_{\leq k^*}) \subseteq X_{\leq t} - \delta(\overline{Y})$. This is a symmetric case and we skip it.

1.3.1. Subclaim. *There are $x_1, x_2 \in X_{\leq t}$, $d_1, d_2 \in X_{\leq t}$, $q^* \geq q$ and non-empty $W_1, W_2 \subseteq k^*$ such that*

- (1) $W_1 \cap W_2 = \emptyset$,
- (2) q^* forces that for all $n \in \{1, 2\}$, $h(x_n) = d_n$,
- (3) q^* forces that for all $n \in \{1, 2\}$, $d_n \in \delta''(\overline{Y}_{\geq k^*}) - (u_0 \cup u_1)$,
- (4) q^* forces that for all $n \in \{1, 2\}$ and $i < k^*$, $d_n = g'_i(x_n)$ iff $i \in W_n$,
- (5) if $d_1 \in X_{v_1}$ and $d_2 \in X_{v_2}$, then $v_1 \leq v_2$,
- (6) for all $n \in \{1, 2\}$ and $i < k^*$, $(g'_i)^{-1}(d_n) \notin U^q$ and $\{(g'_i)^{-1}(d_1) \mid i < k^*\} \cap \{(g'_i)^{-1}(d_2) \mid i < k^*\} = \emptyset$.

Proof. Let $I' \subseteq I$ be the set of those $t' \leq t$ such that $X_{t'} \cap \delta''(\overline{Y}_{\leq k^*}) \neq \emptyset$. Notice that by Lemma 1.1, q decides I' and I' is closed under the least upper bounds. By (β) above, $I' \neq \emptyset$. For $t' \in I'$, we write $I'_{\geq t'}$ for $\{v \in I' \mid v \geq t'\}$. Then for all $t^* \in I'$ there are $W^* \subseteq k^*$ and $q' \geq q$, such that q' forces that excluding $< \kappa$ many x if $h(x) \in \delta''(\overline{Y}_{\geq k^*}) - (u_0 \cup u_1)$ and $x \in X_v$ for some $v \in I'_{\geq t^*}$, then $h(x) = g'_i(x)$ for some $i \in W^*$. We choose these so that $|W^*|$ is minimal. There are two cases:

(I) $|W^*| > 1$: Now it is easy to choose $v_1, v_2 \in I'_{\geq t^*}$, $x_1 \in X_{v_1}$, $x_2 \in X_{v_2}$, $d_1, d_2 \in X_{\leq t}$, $i \in W^*$ and $q^* \geq q'$ so that $v_2 \geq v_1$, for $n \in \{1, 2\}$, q^* forces that $h(x_n) = d_n$, $h(x_n) \in \delta''(\overline{Y}_{\geq k^*}) - (u_0 \cup u_1)$, $h(x_1) \neq g'_i(x_1)$ and $h(x_2) = g'_i(x_2)$. Then we let $W_1 = \{j < k^* \mid g'_j(x_1) = d_1\}$ and $W_2 = \{j < k^* \mid g'_j(x_2) = d_2\}$. By the choice of W^* , these can be chosen so that in addition (6) holds. These are as wanted ($v_2 \geq v_1$ and $h(x_1) \neq g'_i(x_1)$ imply (1)).

(II) $|W^*| = 1$: Let i^* be the only element of W^* . By (α) above, we can find $q'' \geq q'$, $x_1 \in X_{\leq t} - U^q$ and $d_1 \in X_{\leq t}$, so that q'' forces that $h(x_1) = d_1$, $h(x_1) \in \delta''(\overline{Y}_{\geq k^*}) - (u_0 \cup u_1)$ and $h(x_1) \neq g'_{i^*}(x_1)$. Let $v_1 \in I$ be such that $x_1 \in X_{v_1}$ and let v_2 be the least upper bound of v_1 and t^* . By the assumption $(*)$ above, $d_1 \in X_{v_1}$ and so $v_2 \in I'_{\geq t^*}$. Then by Claim 1.1 and the assumption that $|W^*| = 1$, there are $x_2 \in X_{v_2}$, $d_2 \in X_{\leq t}$ and $q^* \geq q''$ such that q^* forces that $h(x_2) = d_2 = g'_{i^*}(x_2)$ and $h(x_2) \in \delta''(\overline{Y}_{\geq k^*}) - (u_0 \cup u_1)$. Then we can let $W_1 = \{j < k^* \mid g'_j(x_1) = d_1\}$ and $W_2 = \{j < k^* \mid g'_j(x_2) = d_2\}$. By (α) and the choice of W^* , these can be chosen so that in addition (6) holds. These are as wanted and proves Subclaim 1.3.1. \square

By the choice of the conditions $p_{x,i}$, we can find $r' \geq q^*$ and $i, i' < \kappa$ so that $p_{x_1,i}, p_{x_2,i'} \leq r'$.

We define a condition r as follows:

- (a) $U^r = U^{r'} \cup \{(g'_i)^{-1}(d_n) \mid i < k, n \in \{1, 2\}\}$,
- (b) $T^r = T^{r'} \cup \{t\}$,
- (c) $v_s^r = v_s^{r'} \cup \{\alpha^*\}$,
- (d) if $(s, \alpha) \neq (t, \alpha^*)$, then $y_{s,\alpha}^r = y_{s,\alpha}^{r'}$,
- (e) $y_{t,\alpha^*}^r = y_{t,\alpha^*}^{r'} \cup \{(g'_i)^{-1}(d_n) \mid i \in W_1, n \in \{1, 2\}\}$.

Then $r \geq q, p_{x_1,i}, p_{x_2,i'}$; in particular r forces that $h(x_1) = d_1$ and $h(x_2) = d_2$. Let $\delta^*(Y_0, \dots, Y_{k^*-1})$ be a boolean term taken inside $X_{\leq t}$ so that Y_i appears in δ^* positively iff $i \in W_1$. Then using Subclaim 1.3.1, it is easy to check that r forces that $d_1 \in \delta^*(\overline{Y}_{<k^*}) \cap \delta''(\overline{Y}_{\geq k^*}) - (u_0 \cup u_1)$ and $d_1 \in h(Y_{t,\alpha^*})$. Since $r \geq q$, this implies that r forces that $\delta^*(\overline{Y}_{<k^*}) \cap \delta''(\overline{Y}_{\geq k^*}) - u_1 \subseteq \delta(\overline{Y}) - u_1 \subseteq h(Y_{t,\alpha^*})$. On

the other hand, using Subclaim 1.3.1 (1), (4) and (5), r forces that $d_2 \in \delta^*(\overline{Y}_{<k^*}) \cap \delta''(\overline{Y}_{\geq k^*}) - (u_0 \cup u_1)$ and $d_2 \notin h(Y_{t,\alpha^*})$, a contradiction which proves Claim 1.3. \square

From now on we work in V^P .

1.4. Claim. *Suppose $h \in \text{Aut}(B)$. There are $h^*, s^* \in I$ $u^* \subseteq X_{\leq s^*}$, $m^* < \omega$, for $i < m^*$, $t_i^* \in I$, $\alpha_i^{s^*} < \kappa^+$, $g_i^{s^*} \in G^*$, for $0 < j < 2^{m^*}$, $f_j^{s^*} \in G^*$ and $f^* \in G^*$ such that if we write $\overline{Y} = (Y_{t_0^{s^*}, \alpha_0^{s^*}, g_0^{s^*}}, \dots, Y_{t_{m^*-1}^{s^*}, \alpha_{m^*-1}^{s^*}, g_{m^*-1}^{s^*}})$, then the following hold:*

- (i) $h^* = h$ or $h^* = h^{-1}$,
- (ii) for all $0 < j < 2^{m^*}$ and $x \in \delta_j^{s^*, m^*}(\overline{Y}) - u^*$, $h^*(x) = f_j^{s^*}(x)$,
- (iii) for all $x \in (X - X_{\leq s^*}) \cup (\delta_0^{s^*, m^*}(\overline{Y}) - u^*)$, $h^*(x) = f^*(x)$,
- (iv) for all $i < m^*$, $t_i^{s^*} \leq s^*$,
- (v) $|u^*| < \kappa$.

Proof. For every $s \in I$, choose $u^s, m^s, t_i^s, \alpha_i^s, g_i^s$ and $f_j^s, i < m^s$ and $j < 2^{m^s}$, as in Claim 1.3. We let I^* to be the set of all $s \in I$ such that for all $j < 2^{m^s}$ and $x \in \delta_j^{s, m^s}(Y_{t_0^s, \alpha_0^s, g_0^s}, \dots, Y_{t_{m^s-1}^s, \alpha_{m^s-1}^s, g_{m^s-1}^s}) - u^s$, $h(x) = f_j^s(x)$. We may assume that I^* is unbounded in I (the case when $I - I^*$ is unbounded is similar) and we let $h^* = h$. We may also assume that these are chosen so that in addition m^s and u^s are minimal (in this order).

We write $E(s), s \in I^*$, for the set of all (t, α) such that for some $i < m^s, t = t_i^s$ and $\alpha = \alpha_i^s$.

1.4.1. Subclaim. *If $r, s \in I^*$ and $r < s$, then*

- (a) $E(r) \subseteq E(s)$,
- (b) for all $j < 2^{m^r}, k < 2^{m^s}$ and $t \leq r$, if

$$\delta_j^{r, m^r}(Y_{t_0^r, \alpha_0^r, g_0^r}, \dots, Y_{t_{m^r-1}^r, \alpha_{m^r-1}^r, g_{m^r-1}^r}) \cap \delta_k^{s, m^s}(Y_{t_0^s, \alpha_0^s, g_0^s}, \dots, Y_{t_{m^s-1}^s, \alpha_{m^s-1}^s, g_{m^s-1}^s}) \cap X_t \neq \emptyset,$$

then $f_j^r(t) = f_k^s(t)$.

Proof. We will prove (a), (b) is immediate by Lemma 1.1. For a contradiction, assume that

- (1) $(t_0^r, \alpha_0^r) \notin E(s)$.

We will write $\overline{Y}^r = (Y_{t_0^r, \alpha_0^r, g_0^r}, \dots, Y_{t_{m^r-1}^r, \alpha_{m^r-1}^r, g_{m^r-1}^r})$ and $\overline{Y}_*^r = (Y_{t_1^r, \alpha_1^r, g_1^r}, \dots, Y_{t_{m^r-1}^r, \alpha_{m^r-1}^r, g_{m^r-1}^r})$ and similarly for s instead of r .

We will show that we can find $a, b < 2^{m^r}, t \leq r$ and a Boolean term δ taken inside $X_{\leq r}$ such that

- (2) $f_a^r(t) \neq f_b^r(t)$,
- (3) $\delta_a^{r, m^r}(\overline{Y}^r) = (X_{\leq r} \cap Y_{t_0^r, \alpha_0^r, g_0^r}) \cap \delta(\overline{Y}_*^r)$ and $X_t \cap \delta_a^{r, m^r}(\overline{Y}^r) \neq \emptyset$,
- (4) $\delta_b^{r, m^r}(\overline{Y}^r) = (X_{\leq r} - Y_{t_0^r, \alpha_0^r, g_0^r}) \cap \delta(\overline{Y}_*^r)$ and $X_t \cap \delta_b^{r, m^r}(\overline{Y}^r) \neq \emptyset$.

If not, then for all $i < 2^{m^r-1}$, we define $f_i \in G^*$ so that for all $x \in \delta_i^{r, m^r-1}(\overline{Y}_*^r) - u^r$, $h(x) = f_i(x)$. This contradicts the minimality of m^r .

Let J be the set of all $t \leq t_0^r$ such that $X_t \cap \delta_i^{r, m^r-1}(\overline{Y}_*^r) \neq \emptyset$. Notice that by Lemma 1.1, J is closed under the least upper bounds. Let $c, d < 2^{m^r}$ be such that $\delta_c^{r, m^r}(\overline{Y}^r) = (X_{\leq r} \cap Y_{t_0^r, \alpha_0^r, g_0^r}) \cap \delta_c^{r, m^r-1}(\overline{Y}_*^r)$ and $\delta_d^{r, m^r}(\overline{Y}^r) = (X_{\leq r} - Y_{t_0^r, \alpha_0^r, g_0^r}) \cap \delta_d^{r, m^r-1}(\overline{Y}_*^r)$. Let J_c be the set of all $t \in J$ such that $X_t \cap \delta_c^{r, m^r}(\overline{Y}^r) \neq \emptyset$. J_d is

defined similarly. Then either J_c or J_d is cofinal in J . If J_d is cofinal, then we let $f_i = f_d^r$ and otherwise we let $f_i = f_c^r$.

We show that f_i is as wanted. We may assume that J_d is not cofinal; the other case is easy. Then since J is closed under the least upper bounds, $J - J_d$ is cofinal. By Lemma 1.1, this means that there is $0 < k < m^r$ such that for all $t \in J - J_d$, $(t_k^r, \alpha_k^r, g_k^r(t)) = (t_0^r, \alpha_0^r, g_0^r(t))$ and $Y_{t_k^r, \alpha_k^r, g_k^r}$ appears positively in $\delta_i^{r, m^r-1}(\overline{Y}_*^r)$. But then $\delta_i^{r, m^r-1}(\overline{Y}_*^r) \subseteq Y_{t_0^r, \alpha_0^r, g_0^r}$ and so f_i is as wanted.

We have shown the existence of a, b, t and δ .

By Claim 1.1 and the fact that $s \geq r$, there is $c < 2^{m^s}$ such that $\delta_c^{s, m^s}(\overline{Y}^s) \cap \delta_a^{r, m^r}(\overline{Y}^r) \cap X_t \not\subseteq u^s$. But then, by (1) above and Claim 1.1, $\delta_c^{s, m^s}(\overline{Y}^s) \cap \delta_b^{r, m^r}(\overline{Y}^r) \cap X_{\leq t} \not\subseteq u^s$. This contradicts (2) and proves Subclaim 1.4.1. \square

Since I is κ^+ -directed, I^* is also κ^+ -directed and so Subclaim 1.4.1 implies that there is s^* such that $E(s^*)$ is maximal and for all $i < m^* = m^{s^*}$, $t_i^{s^*} \leq s^*$. By the choice of u^s , $s \in I^*$, it is easy to see that if $r, s \in I^*$ and $s^* \leq r \leq s$, then $u^r \subseteq u^s$. So s^* can be chosen so that, in addition, for all $s^* \leq r \in I^*$, $u^r = u^{s^*} = u^*$. By Claim 1.1 and Subclaim 1.4.1, there is $f^* \in G^*$ such that for all $s \in I^*$, $f_0^s(s) = f^*(s)$. These are as wanted and completes the the proof of Claim 1.4. \square

Now it follows immediately from Claim 1.4 that $|Aut(B)| \leq \kappa^+ \cdot |I| \cdot |G^*|$ and if for all $t \in I$, G_t is the one element group, then $|Aut(B)| \leq \kappa \cdot |I|$. Finally, for $|Aut(B)| \geq \kappa \cdot |I|$, notice that every permutation of X which is the identity on every element except on $< \kappa$ many, can be lifted to an automorphism of B . \square

2. *Remark.* Notice that in the proof of Theorem 1 one can add $\lambda > \kappa^+$ subsets for the sets $X_{\leq t}$ making the Boolean algebra larger, and the proof still works giving $|G^*| \leq |Aut(B)| \leq \lambda \cdot |I| \cdot |G^*|$.

Our aim was to use Theorem 1 to solve questions on the number of automorphisms of Boolean algebras. We start by first proving a result from [Ro].

3. Conclusion ([Ro]). *Assume that $\kappa^{<\kappa} = \kappa$. Then there is a κ -closed, κ^+ -Knaster po-set P of power κ^+ such that in V^P there is an atomic Boolean algebra B of power κ^+ with $|Aut(B)| = \kappa$.*

Proof. Just choose $I = \{0\}$ and G_0 to be the one element group and apply Theorem 1 (with these choices of I and G_0 , our proof is essentially the same as the related proof in [Ro]). \square

We say that T is a κ -tree if it is a tree of power κ and of height κ . We write T_α for $\{t \in T \mid otp(\{s \in T \mid s < t\}) = \alpha\}$ and say that a κ -tree T is narrow if for all $\alpha < \kappa$, $|T_\alpha| < \kappa$. We say that b is a branch in a tree T if b is a linearly ordered downwards closed subtree of T and the height of b is the same as that of T .

4. Corollary. *Assume that $\kappa^{<\kappa} = \kappa$ and that there is a narrow κ^+ -tree with $\lambda > 0$ branches. Then there is a κ -closed, κ^+ -Knaster po-set P of power κ^+ with the following property: In V^P there is an atomic Boolean algebra B of power κ^+ such that $\lambda \leq |Aut(B)| \leq \kappa^+ \cdot \lambda$.*

Proof. We apply Theorem 1: Clearly we may assume that every element of T belongs to some branch. Then we let $I = \kappa^+$ and for each $\alpha \in I$, let G_α be the free group generated by T_α . Finally, let $\pi_{\alpha, \beta}$ be the natural projection along the branches of T . Then by Theorem 1, it is enough to show the following:

4.1. Claim. *The cardinality of the inverse limit of the system is $\omega \cdot \lambda$ (in V^P).*

Proof. It is enough to show that the forcing does not add branches to T (since $cf(\kappa^+) > \omega$, the inverse limit is the free group generated by the “branches” of T). For a contradiction assume that p forces that b is a new branch in T . For all $\alpha < \kappa^+$, choose $p_\alpha \geq p$ and $t_\alpha \in T_\alpha$ so that p_α forces that $t_\alpha \in b$. Since P is κ^+ -Knaster, there is $Z \subseteq \kappa^+$ of power κ^+ such that for all $\alpha, \beta \in Z$, p_α and p_β are compatible (and so if $\alpha < \beta$, then $t_\alpha < t_\beta$). Then there are no $\alpha \in Z$ so that p_α forces that there are κ^+ many $\beta \in Z$ such that p_β is in the generic set. So for every $\gamma < \kappa^+$ we can find $i_\gamma < \kappa^+$ and $q_\gamma \geq p_{i_\gamma}$ such that $i_\gamma \in Z$, if $\gamma < \gamma'$, then $i_\gamma < i_{\gamma'}$ and for all $j \geq i_{\gamma+1}$, if $j \in Z$, then q_γ forces that p_j does not belong to the generic set. But then $\{q_\gamma \mid \gamma < \kappa^+\}$ is an antichain, a contradiction. \square

5. Conclusion. *Con(ZFC) implies the consistency of ZFC together with the following: There is an atomic Boolean algebra B of power \aleph_1 such that $|Aut(B)| = \aleph_\omega$ (and if one wants $2^{\aleph_0} > \aleph_\omega$).*

Proof. By Corollary 4 it is enough to show that Con(ZFC) implies the consistency of ZFC together with the following: ($2^{\aleph_0} > \aleph_\omega$, if wanted, and) there is a narrow ω_1 -tree T with \aleph_ω branches. This is standard but let us sketch the proof: Assume CH. Let P consist of four-tuples $p = (\alpha^p, T^p, u^p, \{\eta_i^p \mid i \in u^p\})$ such that $\alpha^p < \omega_1$, T^p is a downwards closed countable subtree of $\omega^{\leq \alpha^p}$, u^p is a countable non-empty subset of \aleph_ω and for all $i \in u^p$, η_i^p is a branch in T^p . The order is the obvious one: $p < q$ if $\alpha^p < \alpha^q$, $T^q \upharpoonright \alpha^p + 1 = T^p$, etc. Then P is ω_1 -closed and by CH, it has ω_2 -cc. P also adds the required tree T . (There are no branches other than $\eta_i = \bigcup_{p \in G} \eta_i^p$, $i < \aleph_\omega$, because if p forces that b is a linearly ordered subtree of T and for all $i < \aleph_\omega$, $b \neq \eta_i$, then by using the fact that P is ω_1 -closed, we can find $q > p$ such that q decides $b \upharpoonright \alpha^q$, $b \upharpoonright \alpha^q \neq \eta_i^q \upharpoonright \alpha^q$ for all $i \in u^q$ and no $t \in T^q$ is on top of $b \upharpoonright \alpha^q$. Then q forces that the height of b is $\leq \alpha^q$.) Now, if we want, we can add $> \aleph_\omega$ Cohen reals. As in the proof of Claim 4.1, we can see that this does not add branches to T . \square

REFERENCES

- [Ro] J. Roitman, *The number of automorphism of an atomic Boolean algebra*, Pacific Journal of Mathematics, vol. 94 (1981), 231-242. MR **82j**:03064
 [Mo] J. D. Monk, *Automorphism groups*, J. D. Monk and R. Bonnet (ed.) Handbook of Boolean Algebras, vol. 2, North-Holland, Amsterdam, 1989, 517-546. CMP 21:10

DEPARTMENT OF MATHEMATICS, P.O. BOX 4, 00014 UNIVERSITY OF HELSINKI, FINLAND
E-mail address: thyttine@helsinki.fi

INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY, JERUSALEM, ISRAEL – AND – RUTGERS UNIVERSITY, DEPARTMENT OF MATHEMATICS, NEW BRUNSWICK, NEW JERSEY
E-mail address: shelah@math.huji.ac.il