

## ON ANTOSIK'S LEMMA AND THE ANTOSIK-MIKUSINSKI BASIC MATRIX THEOREM

QU WENBO AND WU JUNDE

(Communicated by Jonathan M. Borwein)

ABSTRACT. That Antosik's Lemma is not a special case of the Antosik-Mikusinski Basic Matrix Theorem will be shown and, an equivalent form of the Antosik-Mikusinski Basic Matrix Theorem will also be presented in this paper.

In [1] and [2], Antosik proved two results which are called the Antosik-Mikusinski Basic Matrix Theorem and Antosik's Lemma, respectively. The theorem and lemma have been proven to be quite effective in treating various topics in Functional Analysis and Set Function Theory [1]–[6]. In [6, 2.2], Swartz thought that Antosik's Lemma is a special case of the Antosik-Mikusinski Basic Matrix Theorem. In [7], Li Ronglu presented the Uniform Convergent Principle. Now, we will show that Swartz's conclusion is incorrect and, the Uniform Convergent Principle is an equivalent form of the Antosik-Mikusinski Basic Matrix Theorem.

**Lemma 1** (Antosik). *Let  $G$  be an abelian topological group and  $x_{ij} \in G$  for  $i, j \in \mathbf{N}$ . Suppose that each strictly increasing sequence  $\{m_i\}$  in  $\mathbf{N}$  has a subsequence  $\{n_i\}$  such that*

(i)  $\lim_i x_{n_i n_j} = 0$  for all  $j \in \mathbf{N}$ , and

(ii)  $\lim_i \sum_{j=1}^{\infty} x_{n_i n_j} = 0$ .

Then  $\lim x_{ii} = 0$ .

Antosik observed that assumption (i) can be dropped if  $G$  is a locally convex space and posed the problem ([2]) of whether (i) can also be dropped in general. In [8], Weber solved this problem showing that assumption (i) is in fact superfluous.

A direct consequence of Lemma 1 is as follows:

**Corollary 1.** *Let  $G$  be an abelian topological group and  $z_{ij} \in G$  for  $i, j \in \mathbf{N}$ . Suppose that*

(I)  $\lim_i z_{ij} = 0$  for each  $j \in \mathbf{N}$ , and

(II) for each strictly increasing sequence of positive integers  $\{m_j\}$  in  $\mathbf{N}$  there is a subsequence  $\{n_j\}$  such that  $\lim_i \sum_{j=1}^{\infty} z_{in_j} = 0$ .

Then  $\lim_i z_{ii} = 0$ .

---

Received by the editors June 20, 2000 and, in revised form, June 11, 2001.

2000 *Mathematics Subject Classification.* Primary 40C05.

*Key words and phrases.* Antosik lemma, basic matrix theorem, uniform convergence principle.

This work was supported by the NSF of China.

Now, we show that the following Antosik-Mikusinski Basic Matrix Theorem can be obtained from Corollary 1.

**Theorem 1** (Antosik-Mikusinski). *Let  $G$  be an abelian topological group,  $x_{ij} \in G$  for  $i, j \in \mathbf{N}$ . Suppose*

(III)  $\lim_i x_{ij} = x_j$  exists for each  $j$ , and

(IV) for each strictly increasing sequence of positive integers  $\{m_j\}$  there is a subsequence  $\{n_j\}$  such that  $\{\sum_j x_{in_j}\}_{i=1}^\infty$  is a Cauchy sequence.

Then  $\lim_i x_{ij} = x_j$  uniformly for  $j \in \mathbf{N}$ . In particular,  $\lim_i x_{ii} = 0$ .

*Proof.* If the conclusion fails, then there exist a closed, symmetric neighbourhood  $V_0$  of 0 in  $G$  and strictly increasing sequences of positive integers  $\{p_k\}$  and  $\{q_k\}$  such that

$$(1) \quad x_{p_k q_k} - x_{q_k} \notin V_0$$

for all  $k$ . Pick a closed, symmetric neighbourhood  $V_1$  of 0 such that  $V_1 + V_1 \subseteq V_0$ . Note that  $x_{p_i q_j} - x_{q_j} \rightarrow 0$  as  $i \rightarrow \infty$  for  $j \in \mathbf{N}$ . Therefore, there exists a subsequence  $\{m_i\}$  of  $\{p_i\}$  such that

$$(2) \quad x_{m_i q_i} - x_{q_i} \in V_1$$

for  $i \in \mathbf{N}$ . We have

$$(3) \quad x_{p_i q_i} - x_{q_i} = (x_{p_i q_i} - x_{m_i q_i}) + (x_{m_i q_i} - x_{q_i}).$$

Consider the matrix  $(x_{p_i q_j} - x_{m_i q_j})$  and note that the matrix satisfies conditions of Corollary 1. Consequently,

$$x_{p_i q_i} - x_{m_i q_i} \rightarrow 0$$

as  $i \rightarrow \infty$ , and

$$x_{p_i q_i} - x_{m_i q_i} \in V_1$$

for sufficiently large  $i$ . Hence, by (3) and (2)

$$x_{p_i q_i} - x_{q_i} \in V_1 + V_1 \subseteq V_0$$

for sufficiently large  $i$ . Which contradicts (1) and we established the result.

Thus, we have Lemma 1  $\Rightarrow$  Corollary 1  $\iff$  Theorem 1. Now, we show that Lemma 1 is not a special case of Theorem 1.  $\square$

**Example 1.** Consider the matrix  $(x_{ij})$  such that  $x_{ij} = 0$  if  $i \neq j+1$  and  $x_{ii-1} = 1$ . Then  $(x_{ij})$  satisfies the assumptions of Lemma 1. If Lemma 1 was a special case of Theorem 1, then the columns should converge to 0 uniformly. But they do not converge to 0 uniformly.

In this way, we have corrected the incorrect statement in ([6, 2.2]).

Now, we use Theorem 1 to prove Theorem 2 below.

**Theorem 2** (Uniform Convergent Principle). *Let  $G$  be an abelian topological group and let  $\Omega$  be a sequentially compact topological space. Let  $\{f_i\}$  be a sequence of sequentially continuous  $G$ -valued functions defined on  $\Omega$ . If each strictly increasing sequence  $\{m_j\}$  in  $\mathbf{N}$  has a subsequence  $\{n_j\}$  such that for each  $\omega \in \Omega$ , the series  $\sum_j f_{n_j}(\omega)$  converges and  $\sum_j f_{n_j} : \Omega \rightarrow G$  is sequentially continuous, then  $\lim_j f_j(\omega) = 0$  uniformly with respect to  $\omega \in \Omega$ .*

*Proof.* We will show that  $f_j(\omega) \rightarrow 0$  uniformly for  $\omega$  in  $\Omega$  or, equivalently, for each sequence  $\{\omega_i\}$  in  $\Omega$ ,

$$(4) \quad f_i(\omega_i) \rightarrow 0$$

as  $i \rightarrow \infty$ . Let  $\{\omega_i\}$  be a sequence of  $\Omega$ . Since  $\Omega$  is sequentially compact, there exists a subsequence  $\{\omega_{n_i}\}$  of  $\{\omega_i\}$  and  $\omega_0$  such that  $\omega_{n_i} \rightarrow \omega_0$ . Consider the matrix  $(f_{n_j}(\omega_{n_i}) - f_{n_j}(\omega_0))$ . Note that the matrix satisfies conditions of Theorem 1. Therefore,

$$f_{n_i}(\omega_{n_i}) - f_{n_i}(\omega_0) \rightarrow 0.$$

Since  $f_{n_i}(\omega_0) \rightarrow 0$ , we get  $f_{n_i}(\omega_{n_i}) \rightarrow 0$ . Hence, by assumption conditions, (4) holds. Thus, Theorem 1  $\implies$  Theorem 2.  $\square$

**Theorem 3.** *Theorem 2 is an equivalent form of Theorem 1.*

*Proof.* Since Corollary 1  $\iff$  Theorem 1  $\implies$  Theorem 2, we only need to show that Theorem 2  $\implies$  Corollary 1.

Indeed, assume the conditions of Corollary 1 are satisfied. Let  $\Omega = \{\frac{1}{i}, 0\}_{i=1}^{\infty}$ , for  $x, y \in \Omega$ , put  $d(x, y) = |x - y|$ . Then  $(\Omega, d)$  is a sequentially compact topological space. Let  $f_j : \Omega \rightarrow G$  satisfy that if  $\omega = \frac{1}{i}$ ,  $f_j(\omega) = z_{ij}$ ; if  $\omega = 0$ ,  $f_j(\omega) = 0$ . It is easily shown that each  $f_j$  is continuous, and for each strictly increasing sequence  $\{m_j\}$  in  $\mathbf{N}$  has subsequence  $\{n_j\}$  such that for  $i \in \mathbf{N}$ , the series  $\sum_j z_{in_j}$  is convergent and  $\lim_i \sum_j z_{in_j} = 0$ . Thus, for each  $\omega \in \Omega$ , the series  $\sum_j f_{n_j}(\omega)$  is convergent and  $\sum_j f_{n_j} : \Omega \rightarrow G$  is continuous. It follows from Theorem 2 that  $\lim_j f_j(\omega) = 0$  uniformly with respect to  $\omega \in \Omega$ . In particular,  $\lim_j f_j(\frac{1}{j}) = \lim_j z_{jj} = 0$ . So Corollary 1 is proved.  $\square$

#### ACKNOWLEDGEMENT

The authors would like to thank Professor Antosik and Professor Li Ronglu for their useful remarks and suggestions.

#### REFERENCES

1. P. Antosik, C. Swartz, Matrix Methods in Analysis, Lecture Notes in Mathematics, 1113, Springer-Verlag, Berlin, 1985. MR **87b**:46079
2. P. Antosik, A lemma on matrices and its applications, Contemp. Math. 52(1986), 89-95. MR **87f**:46022
3. Wu Junde, Li Ronglu, An Orlicz-Pettis theorem with applications to  $\mathcal{A}$ -spaces, Studia Sci. Math. Hungary. 35(1999), 353-358. MR **2001g**:46004
4. Wu Junde, Li Ronglu, Unconditional convergent series on locally convex spaces, Taiwanese J. Math. (4)2000, 253-259. MR **2001b**:46009
5. Wu Junde, Li Ronglu, Hypocontinuity and uniform boundedness for bilinear maps, Studia Sci. Math. Hungary. (35)1999, 133-138. MR **2000a**:46003
6. C. Swartz, Infinite Matrices and the Gliding Hump, World Sci. Publ., Singapore, 1996. MR **98b**:46002
7. Li Ronglu, Cho Minhyung, A uniform convergent principle, J. Harbin Institute of Technology 24(1992), 107-108. MR **94a**:46016
8. H. Weber, A diagonal theorem. Answer to a question of Antosik, Bull. Polish Acad. Sci. Math., 41(1993), 95-102. MR **98c**:22001

DEPARTMENT OF MATHEMATICS, HARBIN INSTITUTE OF TECHNOLOGY, HARBIN 150006, PEOPLE'S REPUBLIC OF CHINA

DEPARTMENT OF MATHEMATICS, ZHEJIANG UNIVERSITY, HANGZHOU 310027, PEOPLE'S REPUBLIC OF CHINA

*E-mail address:* WJD@math.zju.edu.cn