

WHITTLE TYPE INEQUALITY FOR DEMISUBMARTINGALES

B. L. S. PRAKASA RAO

(Communicated by Claudia M. Neuhauser)

ABSTRACT. A Whittle type inequality for demisubmartingales is derived and a strong law of large numbers for functions of a demisubmartingale is obtained.

1. INTRODUCTION

Whittle ([12]) proved an inequality for real valued random variables generalizing the Kolmogorov inequality, the inequality of Hajek-Renyi ([3]) and the inequality of Dufresnoy ([2]). An application of this result for Hilbert space valued random elements $\{Z_k, k \geq 1\}$ such that the family $\{\phi_k(Z_k), k \geq 1\}$ is a real valued submartingale is given in Rao ([8]). An application of this result to obtain a lower bound for the probability of a simultaneous confidence region in multivariate analysis is given in Rao ([8]) sharpening the bound given in Sen ([10]). Recently Shixin ([11]) proved a Hajek-Renyi type inequality for Banach space valued martingales. A Whittle type inequality for Banach space valued martingales was given in Prakasa Rao ([6]) from which the results in Shixin ([11]) follow as special cases.

We now derive a Whittle type inequality for demisubmartingales. This result generalises the recent results on Hajek-Renyi type inequality for demimartingales proved by Christofides ([1]) and the Hajek-Renyi type inequality for associated sequences proved by Prakasa Rao ([5]).

2. PRELIMINARIES

Let $S_i, i \geq 1$, be a sequence of integrable random variables such that

$$(2.1) \quad E\{(S_{j+1} - S_j)f(S_1, \dots, S_j)\} \geq 0, j \geq 1,$$

for every componentwise nondecreasing function f such that the expectation is defined. Then the sequence $\{S_j, j \geq 1\}$ is called a *demimartingale* (cf. Newman and Wright ([4])). If condition (2.1) holds for every componentwise nonnegative nondecreasing function f such that the expectation is defined, then the sequence $\{S_j, j \geq 1\}$ is called a *demisubmartingale*.

A collection of random variables $X_i, 1 \leq i \leq n$, is said to be *associated* if

$$(2.2) \quad Cov(f(X_1, \dots, X_n), g(X_1, \dots, X_n)) \geq 0$$

Received by the editors June 15, 2001 and, in revised form, August 3, 2001.

2000 *Mathematics Subject Classification*. Primary 60E15.

Key words and phrases. Whittle's inequality, Kolmogorov's inequality, maximal inequality, Hajek-Renyi inequality, Dufresnoy's inequality, demisubmartingale, demimartingale, associated sequences, strong law of large numbers.

for any two componentwise nondecreasing functions f and g such that the covariance exists. An infinite sequence of random variables $\{X_n, n \geq 1\}$ is said to be *associated* if every finite subset of $\{X_n, n \geq 1\}$ is associated.

If $X_i, 1 \leq i \leq n$, is an associated sequence of random variables with $E(X_i) = 0, 1 \leq i \leq n$, then the sequence of partial sums $S_i = X_1 + \cdots + X_i, 1 \leq i \leq n$, forms a demimartingale (cf. Newman and Wright [4]).

For an extensive review of the probabilistic properties of associated sequences of random variables and related statistical inference problems, see Prakasa Rao and Dewan ([7]) and Roussas ([9]).

3. WHITTLE TYPE INEQUALITY

Let $S_n, n \geq 1$, be a demisubmartingale and $\phi(\cdot)$ be a nondecreasing convex function. Then the sequence $\phi(S_n), n \geq 1$, is a demisubmartingale by Lemma 2.1 of Christofides ([1]).

We now state our main theorem.

Theorem 3.1. *Let the sequence of random variables $\{S_n, n \geq 1\}$ be a demisubmartingale and $\phi(\cdot)$ be a nonnegative nondecreasing convex function such that $\phi(S_0) = 0$. Let $\psi(u)$ be a positive nondecreasing function for $u > 0$. Let A_n be the event that $\phi(S_k) \leq \psi(u_k), 1 \leq k \leq n$, where $0 = u_0 < u_1 \leq \cdots \leq u_n$. Then*

$$(3.1) \quad P(A_n) \geq 1 - \sum_{k=1}^n \frac{E[\phi(S_k)] - E[\phi(S_{k-1})]}{\psi(u_k)}.$$

If, in addition, there exist nonnegative real numbers $\Delta_k, 1 \leq k \leq n$, such that

$$\begin{aligned} 0 &\leq E[(\phi(S_k) - \phi(S_{k-1}))f(\phi(S_1), \dots, \phi(S_{k-1}))] \\ &\leq \Delta_k E[f(\phi(S_1), \dots, \phi(S_{k-1}))], 1 \leq k \leq n, \end{aligned}$$

for all componentwise nonnegative nondecreasing functions f such that the expectation is defined and

$$\psi(u_k) \geq \psi(u_{k-1}) + \Delta_k, 1 \leq k \leq n,$$

then

$$(3.2) \quad P(A_n) \geq \prod_{k=1}^n \left(1 - \frac{\Delta_k}{\psi(u_k)}\right).$$

Remarks. The above result is an analogue of the inequality in Whittle ([12]) for real valued random variables. A version of Theorem 3.1 for a sequence of Hilbert space valued random elements was proved in Rao ([8]) and an application to Banach space valued martingales is given in Prakasa Rao ([6]).

Proof. Since the sequence $\{S_n, n \geq 1\}$ is a demisubmartingale by hypothesis and the function $\phi(\cdot)$ is a nondecreasing convex function, it follows that the sequence $\{\phi(S_n), n \geq 1\}$ forms a demisubmartingale by Lemma 2.1 of Christofides ([1]). Hence

$$(3.3) \quad E\{(\phi(S_{n+1}) - \phi(S_n))f(\phi(S_1), \dots, \phi(S_n))\} \geq 0, n \geq 1,$$

for every nonnegative componentwise nondecreasing function f such that the expectation is defined.

Let χ_j be the indicator function of the event $[\phi(S_j) \leq \psi(u_j)]$ for $1 \leq j \leq n$.

Note that

$$\chi_n \geq \left(1 - \frac{\phi(S_n)}{\psi(u_n)}\right)$$

and hence

$$\begin{aligned} P(A_n) &= E\left(\prod_{i=1}^n \chi_i\right) = E\left(\left\{\prod_{i=1}^{n-1} \chi_i\right\} \chi_n\right) \\ &\geq E\left(\left\{\prod_{i=1}^{n-1} \chi_i\right\} \left(1 - \frac{\phi(S_n)}{\psi(u_n)}\right)\right). \end{aligned}$$

Therefore

$$\begin{aligned} E\left[\left\{\prod_{i=1}^{n-1} \chi_i\right\} \left\{\left(1 - \frac{\phi(S_n)}{\psi(u_n)}\right) - \left(1 - \frac{\phi(S_{n-1})}{\psi(u_n)}\right)\right\} + \frac{\phi(S_n) - \phi(S_{n-1})}{\psi(u_n)}\right] \\ = E\left[\left(1 - \prod_{i=1}^{n-1} \chi_i\right) \left(\frac{\phi(S_n) - \phi(S_{n-1})}{\psi(u_n)}\right)\right] \geq 0 \end{aligned}$$

since the function $1 - \prod_{i=1}^{n-1} \chi_i$ is a nonnegative componentwise nondecreasing function of $\phi(S_i)$, $1 \leq i \leq n-1$. Hence

$$\begin{aligned} P(A_n) &\geq E\left(\left\{\prod_{i=1}^{n-1} \chi_i\right\} \left(1 - \frac{\phi(S_{n-1})}{\psi(u_n)}\right)\right) - \frac{E\{\phi(S_n)\} - E\{\phi(S_{n-1})\}}{\psi(u_n)} \\ &\geq E\left(\left\{\prod_{i=1}^{n-2} \chi_i\right\} \left(1 - \frac{\phi(S_{n-1})}{\psi(u_{n-1})}\right)\right) - \frac{E\{\phi(S_n)\} - E\{\phi(S_{n-1})\}}{\psi(u_n)}. \end{aligned}$$

The last inequality follows from the observation that the sequence $\psi(u_n)$, $n \geq 1$, is positive and nondecreasing.

Applying this inequality repeatedly, we get that

$$(3.4) \quad P(A_n) \geq 1 - \sum_{k=1}^n \frac{E[\phi(S_k)] - E[\phi(S_{k-1})]}{\psi(u_k)},$$

completing the proof of the first part of the theorem.

Note that

$$\begin{aligned} E\left\{\prod_{i=1}^{n-1} \chi_i \left(1 - \frac{\phi(S_n)}{\psi(u_n)}\right) - \left(1 - \frac{\Delta_n}{\psi(u_n)}\right) \left(1 - \frac{\phi(S_{n-1})}{\psi(u_{n-1})}\right) \prod_{i=1}^{n-1} \chi_i\right\} \\ \geq E\left\{\frac{\phi(S_{n-1})}{\psi(u_n)\psi(u_{n-1})} [\psi(u_n) - \psi(u_{n-1}) - \Delta_n] \prod_{i=1}^{n-1} \chi_i\right\} \end{aligned}$$

and the last term is nonnegative by hypothesis. Hence

$$(3.5) \quad P(A_n) \geq \left(1 - \frac{\Delta_n}{\psi(u_n)}\right) E\left(\left\{\prod_{i=1}^{n-2} \chi_i\right\} \left(1 - \frac{\phi(S_{n-1})}{\psi(u_{n-1})}\right)\right).$$

Applying this inequality repeatedly, we obtain that

$$(3.6) \quad P(A_n) \geq \prod_{k=1}^n \left(1 - \frac{\Delta_k}{\psi(u_k)}\right).$$

4. APPLICATIONS

Suppose $\{S_n, n \geq 1\}$ is a demisubmartingale. Then $\{(S_n^+)^p, n \geq 1\}$ and $\{(S_n^-)^p, n \geq 1\}$ are demisubmartingales by Corollary 2.1 of Christofides ([1]). Furthermore $|S_n|^p = (S_n^+)^p + (S_n^-)^p$ for all $p \geq 1$.

(1) Let $\psi(u) = u^p, p \geq 1$. Applying Theorem 3.1, we get that

$$(4.1) \quad P(S_j^+ \leq u_j, 1 \leq j \leq n) \geq 1 - \sum_{j=1}^n \frac{E(S_j^+)^p - E(S_{j-1}^+)^p}{u_j^p}$$

and

$$(4.2) \quad P(S_j^- \leq u_j, 1 \leq j \leq n) \geq 1 - \sum_{j=1}^n \frac{E(S_j^-)^p - E(S_{j-1}^-)^p}{u_j^p}.$$

Hence, for every $\varepsilon > 0$,

$$\begin{aligned} P\left(\sup_{1 \leq j \leq n} \frac{|S_j|}{u_j} \geq \varepsilon\right) &= P\left(\sup_{1 \leq j \leq n} \frac{|S_j|^p}{u_j^p} \geq \varepsilon^p\right) \\ &= P\left(\sup_{1 \leq j \leq n} \frac{(S_j^+)^p + (S_j^-)^p}{u_j^p} \geq \varepsilon^p\right) \\ &= P\left(\sup_{1 \leq j \leq n} \frac{(S_j^+)^p}{u_j^p} \geq \frac{1}{2}\varepsilon^p\right) \\ &\quad + P\left(\sup_{1 \leq j \leq n} \frac{(S_j^-)^p}{u_j^p} \geq \frac{1}{2}\varepsilon^p\right) \\ &\leq 2\varepsilon^{-p} \sum_{j=1}^n \frac{E(S_j^+)^p - E(S_{j-1}^+)^p}{u_j^p} \\ &\quad + 2\varepsilon^{-p} \sum_{j=1}^n \frac{E(S_j^-)^p - E(S_{j-1}^-)^p}{u_j^p} \\ &\leq 2\varepsilon^{-p} \sum_{j=1}^n \frac{E|S_j|^p - E|S_{j-1}|^p}{u_j^p}. \end{aligned}$$

In particular for $p=2$, we have

$$(4.3) \quad P\left(\sup_{1 \leq j \leq n} \frac{|S_j|}{u_j} \geq \varepsilon\right) \leq 2\varepsilon^{-2} \sum_{j=1}^n \frac{ES_j^2 - ES_{j-1}^2}{u_j^2},$$

which is the Hajek-Renyi type inequality for associated sequences derived in Corollary 2.3 of Christofides ([1]).

Suppose $p = 1$. Let $\phi(x) = \max(0, x)$. Then $\phi(x)$ is a nonnegative nondecreasing convex function and it is clear that $S_n \leq S_n^+ = \phi(S_n)$ for every $n \geq 1$. Let $\psi(u) = u$. Then

$$\begin{aligned} P\left(\sup_{1 \leq j \leq n} \frac{S_j}{u_j} \geq \varepsilon\right) &\leq P\left(\sup_{1 \leq j \leq n} \frac{S_j^+}{u_j} \geq \varepsilon\right) \\ &\leq \varepsilon^{-1} \sum_{j=1}^n \frac{ES_j^+ - ES_{j-1}^+}{u_j} \end{aligned}$$

by Theorem 3.1 which is the Chow type maximal inequality derived in Theorem 2.1 of Christofides ([1]).

(2) Let $p=2$ again in the above discussion. If

$$E(S_j^2 - S_{j-1}^2) \leq u_j^2 - u_{j-1}^2$$

for $1 \leq j \leq n$, then

$$P(A_n) \geq \prod_{j=1}^n \left(1 - \frac{E(S_j^2) - E(S_{j-1}^2)}{u_j^2}\right)$$

which is an analogue of the Dufresnoy's inequality.

(3) Let $\{S_n, n \geq 1\}$ be a demisubmartingale and $\phi(\cdot)$ be a nonnegative nondecreasing convex function such that $\phi(S_0) = 0$. Let $\psi(u)$ be a positive nondecreasing function for $u > 0$. Then, for any nondecreasing sequence $u_n, n \geq 1$ with $u_0 = 0$,

$$(4.4) \quad P\left(\sup_{1 \leq j \leq n} \frac{\phi(S_j)}{\psi(u_j)} \geq \varepsilon\right) \leq \varepsilon^{-1} \sum_{k=1}^n \frac{E[\phi(S_k)] - E[\phi(S_{k-1})]}{\psi(u_k)}.$$

In particular, for any fixed $n \geq 1$,

$$(4.5) \quad P\left(\sup_{k \geq n} \frac{\phi(S_k)}{\psi(u_k)} \geq \varepsilon\right) \leq \varepsilon^{-1} \left[E\left(\frac{\phi(S_n)}{\psi(u_n)}\right) + \sum_{k=n+1}^{\infty} \frac{E[\phi(S_k)] - E[\phi(S_{k-1})]}{\psi(u_k)} \right].$$

We now derive a strong law of large numbers for functions of demisubmartingales.

Theorem 4.1. *Let $\{S_n, n \geq 1\}$ be a demisubmartingale and $\phi(\cdot)$ be a nonnegative nondecreasing convex function such that $\phi(S_0) = 0$. Let $\psi(u)$ be a positive nondecreasing function for $u > 0$ such that $\psi(u) \rightarrow \infty$ as $u \rightarrow \infty$. Further suppose that*

$$(4.6) \quad \sum_{k=1}^{\infty} \frac{E[\phi(S_k)] - E[\phi(S_{k-1})]}{\psi(u_k)} < \infty$$

for a nondecreasing sequence $u_n \rightarrow \infty$ as $n \rightarrow \infty$. Then

$$(4.7) \quad \frac{\phi(S_n)}{\psi(u_n)} \xrightarrow{a.s} 0 \text{ as } n \rightarrow \infty.$$

Proof of this result follows by the standard arguments following the inequality (4.5) given above. We omit the details.

REFERENCES

1. Christofides, T.C.: Maximal inequalities for demimartingales and a strong law of large numbers, *Statist. Probab. Lett.*, 50 (2000), 357-363. MR **2001m**:60032
2. Dufresnoy, J.: Autour de l'inegalite de Kolmogorov, *C. R. Acad. Sci. Paris*, 264A (1967), 603. MR **37**:5909
3. Hajek, J. and Renyi, A.: A generalization of an inequality of Kolomogorov, *Acta. Math. Acad. Sci. Hung.*, 6 (1955), 281-284. MR **17**:864a
4. Newman, C.M. and Wright, A.L.: Associated random variables and martingale inequalities, *Z. Wahrsch. Verw. Geb.*, 59 (1982), 361-371. MR **85d**:60088
5. Prakasa Rao, B.L.S.: Hajek-Renyi type inequality for associated sequences, *Statist. Probab. Lett.*, 2001 (to appear).
6. Prakasa Rao, B.L.S.: Application of Whittle's inequality for Banach space valued martingales, *Statist. Probab. Lett.*, 2001 (to appear).
7. Prakasa Rao, B.L.S. and Dewan, I.: Associated sequences and related inference problems, In *Handbook of Statistics: Stochastic Processes: Theory and Methods*, (Edited by D.N. Shanbhag and C.R.Rao), North Holland, Amsterdam, 19 (2001), 693-728. CMP 2002:03

8. Rao, P.: Whittle's inequality in Hilbert space, *Theory of Probability and Mathematical Statistics*, 16 (1978), 111-116. MR **55**:13517
9. Roussas, G.G.: Positive and negative dependence with some statistical applications, In *Asymptotics, Nonparametrics and Time Series*, (Edited by S.Ghosh), Marcel Dekker, New York, 1999, 757-788. CMP 2000:05
10. Sen, P.K.: A Hajek-Renyi inequality for stochastic vectors with applications to simultaneous confidence regions, *Ann. Math. Statist.*, 42 (1971), 1132-1134.
11. Shixin, Gan.: The Hajek-Renyi inequality for Banach space valued martingales and the p smoothness of Banach spaces, *Statist. Probab. Lett.*, 32 (1997), 245-248. MR **97k**:60006
12. Whittle, P.: Refinements of Kolmogorov's inequality, *Theory Probab. Appl.*, 14 (1969), 310-311. MR **39**:7657

INDIAN STATISTICAL INSTITUTE, 7, SJSS MARG, NEW DELHI 110 016, INDIA
E-mail address: blsp@isid.ac.in