

## ON A PROBLEM OF J. P. WILLIAMS

EDWARD KISSIN AND VICTOR S. SHULMAN

(Communicated by David R. Larson)

ABSTRACT. Let  $B(H)$  be the algebra of all bounded operators on a Hilbert space  $H$ . Let  $g$  be a continuous function on the closed disk  $D$  and let

$$\|g(A)X - Xg(A)\| \leq C\|AX - XA\|,$$

for some  $C > 0$ , for all  $X \in B(H)$  and all  $A \in B(H)$  with  $\|A\| \leq 1$ . Then  $g$  is differentiable on  $D$ . The paper shows that the function  $g$  may have a discontinuous derivative.

### 1. INTRODUCTION

Let  $B(H)$  be the algebra of all bounded operators on a Hilbert space  $H$  and  $\mathbf{B}_1$  be the unit ball of  $B(H)$ . For  $A, B \in B(H)$ , we denote by  $[A, B]$  their commutator  $AB - BA$ . Let  $D = \{z \in \mathbb{C} : |z| \leq 1\}$  be the closed unit disk. In his paper [7] Williams raised the following problem. If  $g$  is a continuous complex-valued function on  $D$ , possessing the property

$$(1) \quad \|[g(A), X]\| \leq C\|[A, X]\|,$$

for some  $C > 0$ , for any  $X \in B(H)$  and any normal operator  $A$  in  $\mathbf{B}_1$ , must  $g$  always be continuously differentiable on  $D$ ?

It should be noted that Johnson and Williams proved earlier [2, Theorem 4.1] that  $g$  must be differentiable on  $D$  and therefore analytic in the interior  $D^\circ$  of  $D$ , and its derivative must be bounded on  $D$ .

We will show that the answer to Williams's problem is negative. Moreover, we will show that the function on  $D$  may have a discontinuous derivative even if it satisfies (1) for *all* (not necessarily normal) contractions  $A$ .

The authors are grateful to the referee for his useful suggestions.

### 2. FULLY OPERATOR LIPSCHITZ FUNCTIONS

We denote by  $\mathcal{A}(D)$  the *disk algebra*: the algebra of all continuous complex-valued functions on  $D$  which are analytic on  $D^\circ$ . The algebra  $\mathcal{A}(D)$  is a closed subalgebra of the algebra  $C(D)$  of all continuous complex-valued functions on  $D$  with the norm  $\|g\| = \sup_{z \in D} |g(z)|$ . The subalgebra  $P(D)$  of all polynomials on  $D$  is dense in  $\mathcal{A}(D)$  (see, for example, [4, §3.2.13]).

---

Received by the editors March 19, 2001 and, in revised form, July 6, 2001.  
2000 *Mathematics Subject Classification*. Primary 47A56.

By von Neumann’s theorem (see [6, Proposition I.8.3]),  $\|p(A)\| \leq \|p\|$  for any polynomial  $p$  and any  $A \in \mathbf{B}_1$ . Therefore functions from  $\mathcal{A}(D)$  act on  $\mathbf{B}_1$  and

$$(2) \quad \|g(A)\| \leq \|g\|, \quad \text{for any } g \in \mathcal{A}(D) \text{ and } A \in \mathbf{B}_1.$$

We call a function  $g \in \mathcal{A}(D)$  *Fully Operator Lipschitzian* if there is  $C > 0$  such that

$$(3) \quad \|g(A) - g(B)\| \leq C\|A - B\|, \quad \text{for } A, B \in \mathbf{B}_1.$$

The class of Fully Operator Lipschitz functions is contained in the wider class of Operator Lipschitz functions on  $D$  which consists of all continuous functions on  $D$  satisfying inequality (3) for all normal operators in  $\mathbf{B}_1$  (see [3]). The function  $g(z) = \bar{z}$ , for example, is Operator Lipschitzian on  $D$ , since  $\|A^* - B^*\| = \|A - B\|$ , for all normal  $A, B \in \mathbf{B}_1$ . However, it is not Fully Operator Lipschitzian. Both classes of functions are important for applications in mathematical physics and have attracted much attention (see, for example, Bibliography in [1]).

**Proposition 1.** *A function  $g \in \mathcal{A}(D)$  is Fully Operator Lipschitzian if and only if there is  $C > 0$  such that*

$$(4) \quad \|[g(A), X]\| \leq C\|[A, X]\|, \quad \text{for } A \in \mathbf{B}_1 \text{ and } X \in B(H).$$

*Proof.* If  $A, B \in \mathbf{B}_1$ , the operator  $L = \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$  on  $H \oplus H$  belongs to the unit ball of  $B(H \oplus H)$ . Let  $X = \begin{pmatrix} 0 & \mathbf{1} \\ 0 & 0 \end{pmatrix}$ . Clearly, (4) holds for operators on  $H \oplus H$ . Hence  $\|[g(L), X]\| \leq C\|[L, X]\|$  which implies (3).

Conversely, let  $\|A\| < 1$ . For  $X \in B(H)$ , the operators  $A(t) = e^{tX}Ae^{-tX}$  belong to  $\mathbf{B}_1$  for sufficiently small  $t$ . If (3) holds then, taking into account that  $g(e^{tX}Ae^{-tX}) = e^{tX}g(A)e^{-tX}$ , we obtain

$$\|g(A) - e^{tX}g(A)e^{-tX}\| = \|g(A) - g(e^{tX}Ae^{-tX})\| \leq C\|A - e^{tX}Ae^{-tX}\|.$$

Dividing through by  $t$  and taking the limit as  $t \rightarrow 0$ , we have that (4) holds.

Let  $\|A\| = 1, X \in B(H)$ . For  $r < 1, \|[g(rA), X]\| \leq C\|[rA, X]\|$ . Taking the limit as  $r \rightarrow 1$ , we obtain that (4) holds. □

It follows from Proposition 1 that our aim is to construct a Fully Operator Lipschitz function with discontinuous derivative.

### 3. FULLY OPERATOR LIPSCHITZ FUNCTIONS WITH DISCONTINUOUS DERIVATIVE

Consider the following function on  $D$ :

$$h(1) = 0 \quad \text{and} \quad h(z) = (z - 1)^2 \exp((z - 1)^{-1}), \quad \text{for } z \in D, z \neq 1.$$

Since  $\frac{x-1}{(x-1)^2+y^2} < 0$ , if  $z = x + iy \in D \setminus 1$ , we have that

$$(5) \quad \begin{aligned} \sup_{z \in D \setminus 1} |\exp((z - 1)^{-1})| &= \sup_{z \in D \setminus 1} \left| \exp\left(\frac{(x - 1) - iy}{(x - 1)^2 + y^2}\right) \right| \\ &= \sup_{z \in D \setminus 1} \left| \exp\left(\frac{x - 1}{(x - 1)^2 + y^2}\right) \right| \left| \exp\left(\frac{iy}{(x - 1)^2 + y^2}\right) \right| \\ &= \sup_{z \in D \setminus 1} \left| \exp\left(\frac{x - 1}{(x - 1)^2 + y^2}\right) \right| < 1. \end{aligned}$$

The function  $h$  is analytic on  $D^\circ$  and continuous on  $D$ , since, by (5),

$$|h(z)| = |h(x + iy)| = |z - 1|^2 |\exp((z - 1)^{-1})| \leq |z - 1|^2 \rightarrow 0,$$

as  $z \rightarrow 1$ . Thus  $h \in \mathcal{A}(D)$ . We obtain similarly that

$$\left| \frac{h(z) - h(1)}{z - 1} \right| = |z - 1| |\exp((z - 1)^{-1})| \leq |z - 1| \rightarrow 0,$$

as  $z \rightarrow 1$ , so  $h'(1) = 0$ . We also obtain that

$$h'(z) = 2(z - 1) \exp((z - 1)^{-1}) - \exp((z - 1)^{-1}), \quad \text{for } z \in D, z \neq 1.$$

We have, as above, that  $(z - 1) \exp((z - 1)^{-1}) \rightarrow 0$ , as  $z \rightarrow 1$ , while  $\exp((z - 1)^{-1})$  does not have limit as  $z \rightarrow 1$ . Therefore  $h'$  is discontinuous at  $z = 1$ .

**Theorem 2.** *The function  $h$  is Fully Operator Lipschitzian.*

*Proof.* By Proposition 1, we only need to prove that (4) holds for  $h$ . For  $0 < \lambda < 1$ , set  $h_\lambda(z) = h(\lambda z)$ . Every  $h_\lambda$  is analytic in a neighbourhood of  $D$ , so it belongs to  $\mathcal{A}(D)$ , and  $\|h - h_\lambda\| \rightarrow 0$ , as  $\lambda \rightarrow 1$ . Hence it follows from (2) that

$$(6) \quad \begin{aligned} \|[h(A), X] - [h_\lambda(A), X]\| &= \|[(h(A) - h_\lambda(A)), X]\| \\ &\leq 2\|h(A) - h_\lambda(A)\| \|X\| \leq 2\|h - h_\lambda\| \|A\| \|X\| \rightarrow 0. \end{aligned}$$

For any  $A \in \mathbf{B}_1$  and  $X \in B(H)$ ,

$$\begin{aligned} \|[h_\lambda(A), X]\| &= \|[(\lambda A - \mathbf{1}) \exp((\lambda A - \mathbf{1})^{-1})(\lambda A - \mathbf{1}), X]\| \\ &= \|[(\lambda A - \mathbf{1}), X] \exp((\lambda A - \mathbf{1})^{-1})(\lambda A - \mathbf{1}) \\ &\quad + (\lambda A - \mathbf{1})[\exp((\lambda A - \mathbf{1})^{-1}), X](\lambda A - \mathbf{1}) \\ &\quad + (\lambda A - \mathbf{1}) \exp((\lambda A - \mathbf{1})^{-1})[(\lambda A - \mathbf{1}), X]\| \\ &\leq 2\lambda\|[A, X]\| \|\exp((\lambda A - \mathbf{1})^{-1})\| \|\lambda A - \mathbf{1}\| \\ &\quad + \|(\lambda A - \mathbf{1})[\exp((\lambda A - \mathbf{1})^{-1}), X](\lambda A - \mathbf{1})\|. \end{aligned}$$

We have that  $\|\lambda A - \mathbf{1}\| < 2$  and that the function  $\exp((\lambda z - 1)^{-1})$  belongs to  $\mathcal{A}(D)$ . We obtain from (2) and (5) that

$$(7) \quad \|\exp((\lambda A - \mathbf{1})^{-1})\| \leq \|\exp((\lambda z - 1)^{-1})\| \leq \sup_{z \in D \setminus 1} |\exp((z - 1)^{-1})| < 1.$$

Therefore

$$(8) \quad \|[h_\lambda(A), X]\| \leq 4\lambda\|[A, X]\| + \|(\lambda A - \mathbf{1})[\exp((\lambda A - \mathbf{1})^{-1}), X](\lambda A - \mathbf{1})\|.$$

It follows from Lemma 2 of [5] that, for any  $B \in B(H)$ ,

$$[\exp(B), X] = \int_0^1 \exp(tB)[B, X] \exp((1 - t)B) dt.$$

If  $B$  is invertible, then  $B[B^{-1}, X]B = [X, B]$ . Hence

$$\begin{aligned} &\|(\lambda A - \mathbf{1})[\exp((\lambda A - \mathbf{1})^{-1}), X](\lambda A - \mathbf{1})\| \\ &= \left\| \int_0^1 \exp(t(\lambda A - \mathbf{1})^{-1})(\lambda A - \mathbf{1})[(\lambda A - \mathbf{1})^{-1}, X](\lambda A - \mathbf{1}) \right. \\ &\quad \left. \exp((1 - t)(\lambda A - \mathbf{1})^{-1}) dt \right\| \\ &\leq \|[X, \lambda A - \mathbf{1}]\| \int_0^1 \|\exp(t(\lambda A - \mathbf{1})^{-1})\| \|\exp((1 - t)(\lambda A - \mathbf{1})^{-1})\| dt. \end{aligned}$$

As in (7), we have that

$$\|\exp(t(\lambda A - \mathbf{1})^{-1})\| < 1 \quad \text{and} \quad \|\exp((1-t)(\lambda A - \mathbf{1})^{-1})\| < 1.$$

Therefore

$$\|(\lambda A - \mathbf{1})[\exp((\lambda A - \mathbf{1})^{-1}), X](\lambda A - \mathbf{1})\| \leq \lambda\| [A, X] \|.$$

Hence we obtain from (8) that  $\| [h_\lambda(A), X] \| \leq 5\lambda\| [A, X] \|$ . Combining this with (6), we conclude that  $\| [h(A), X] \| \leq 5\| [A, X] \|$ .  $\square$

#### REFERENCES

1. D. R. Jocić, *Integral representation formula for generalized normal derivation*, Proc. Amer. Math. Soc. **127** (1999), 2303–2314. MR **99j**:47026
2. B. E. Johnson and J. P. Williams, *The range of a normal derivation*, Pacific J. Math. **58** (1975), 105–122. MR **52**:1390
3. E. Kissin and V. S. Shulman, *Classes of Operator-smooth Functions. I. Operator Lipschitz Functions*, preprint, (2000).
4. T. W. Palmer, *Banach Algebras and the General Theory of \*-algebras*, vol. I, CUP, 1994. MR **95c**:46002
5. R. Powers, *A remark on the domain of an unbounded derivation of a C\*-algebra*, J. Funct. Anal. **18** (1975), 85–95. MR **52**:1334
6. B. Sz.-Nagy and C. Foias, *Analyse Harmonique des Operateurs de l'espace de Hilbert*, Academiai Kiado, Budapest, 1967. MR **37**:778
7. J. P. Williams, *Derivation ranges: open problems*, Topics in Modern Operator Theory, (Timisoara/Herculane, 1980), 319–328, Operator Theory: Adv. Appl., 2, Birkhäuser, Basel-Boston, MA, 1981. MR **83k**:47027

SCHOOL OF COMMUNICATIONS TECHNOLOGY AND MATHEMATICAL SCIENCES, UNIVERSITY OF NORTH LONDON, HOLLOWAY, LONDON N7 8DB, GREAT BRITAIN

*E-mail address*: e.kissin@unl.ac.uk

SCHOOL OF COMMUNICATIONS TECHNOLOGY AND MATHEMATICAL SCIENCES, UNIVERSITY OF NORTH LONDON, HOLLOWAY, LONDON N7 8DB, GREAT BRITAIN – AND – DEPARTMENT OF MATHEMATICS, VOLOGDA STATE TECHNICAL UNIVERSITY, VOLOGDA, RUSSIA

*E-mail address*: shulman\_v@yahoo.com