

**A CORRECTION TO
 “ULTRADIFFERENTIABLE FUNCTIONS ON LINES IN \mathbb{R}^n ”**

TEJINDER NEELON

(Communicated by Christopher D. Sogge)

ABSTRACT. The proof of Theorem 1 in Proc. Amer. Math. Soc. **127** (1999), no. 7, 2099–2104, is revised.

The proof of Theorem 1 in [1] uses Lemma 3(ii) which turns out to be valid only for the case $n = 2$. In this note the proof of Theorem 1 is revised by replacing Lemma 3(ii) by the proposition stated below.

For $x \in \mathbb{R}^n$, $0 \neq \xi \in \mathbb{S}^{n-1}$, let $H_{x\xi}$ denote the hyperplane in \mathbb{R}^n that passes through x with ξ as its normal vector. If $\varphi : \mathbb{R}^{n-1} \rightarrow \mathbb{R}^n$ is a linear map such that $H_{x\xi} = \varphi(\mathbb{R}^{n-1})$, then the restriction of a function $u \in C^\infty(\mathbb{R}^n)$ to $H_{x\xi}$ is defined by $u_{x\xi}(\mathbf{t}) = u(x + \varphi(\mathbf{t}))$, $\mathbf{t} \in \mathbb{R}^{n-1}$.

Proposition. *For any $u \in C^\infty(\mathbb{R}^n)$, the following inequality holds:*

$$(0.1) \quad \max_{|\alpha|=k} |\partial^\alpha u(x)| \leq \max_{\beta \in \mathbb{Z}_+^{n-1}, |\beta|=k} \max_{|\xi|=1} |\partial^\beta (u_{x\xi})(0)|, \forall k \geq 0.$$

Proof. Conforming to the notation in [1], let $a_1, \varepsilon \in \mathbb{R}, \varepsilon > 0, x \in \mathbb{R}^n, k \in \mathbb{Z}, k \geq 0$, and $(\alpha_1, \alpha_2, \dots, \alpha_n) \in \mathbb{Z}_+^n, |\alpha| = k$, be fixed. Put $a_{1j} = a_1 + \frac{\varepsilon j}{k}, j = 0, \dots, k$. Consider the hyperplanes given by the images of the linear maps

$$\begin{aligned} \varphi_j & : \mathbb{R}^{n-1} \rightarrow \mathbb{R}^n, \\ \varphi_j(t_2, \dots, t_n) & = (x_1 + a_{1j}t_2, x_2 + t_2, \dots, x_n + t_n), \forall j, 0 \leq j \leq k. \end{aligned}$$

Then,

$$\partial_2^{\alpha_1 + \alpha_2} \partial^{\alpha''} (u \circ \varphi_j)(0) = \sum_{l=0}^{\alpha_1 + \alpha_2} a_{1j}^{\alpha_1} \left(\partial_1^l \partial_2^{\alpha_1 + \alpha_2 - l} \partial^{\alpha''} u \right) (x),$$

where $\partial^{\alpha''} = \partial_3^{\alpha_3} \dots \partial_n^{\alpha_n}$. By applying Lemma 3(i), we have for $\varepsilon = 8e^3$ and $l = \alpha_1$,

$$|\partial^\alpha u(x)| \leq \max_{|\beta|=|\alpha|} \max_{0 \leq r \leq k} |\partial^\beta (u \circ \varphi_r)(0)|.$$

□

Let V be an n -dimensional vector space. By using a linear isomorphism between V and \mathbb{R}^n , the classes $C^\infty(V)$ and $C^M(V)$ can be identified with $C^\infty(\mathbb{R}^n)$ and

Received by the editors June 20, 2002.

2000 *Mathematics Subject Classification.* Primary 30D60, 46F05.

Key words and phrases. Ultradifferentiable functions, transfinite diameter.

©2002 American Mathematical Society

$C^M(\mathbb{R}^n)$, respectively. When $n = 2$, the inequality (0.1) reduces to the last inequality in the proof of Theorem 1. In particular, the conclusion of Theorem 1 is valid for any 2-dimensional vector space V . Hence, by the above proposition and by induction on n , we see that Theorem 1 holds for all n .

REFERENCE

1. Tejinder Neelon, *Ultradifferentiable functions on lines in \mathbb{R}^n* , Proc. Amer. Math. Soc. **127** (1999), no. 7, 2099–2104. MR **99j**:46044

DEPARTMENT OF MATHEMATICS, 333 S. TWIN OAKS VALLEY ROAD, CALIFORNIA STATE UNIVERSITY, SAN MARCOS, CALIFORNIA 92056
E-mail address: `neelon@csusm.edu`