

UNIFORM STABILITY OF RESOLVENT FAMILIES

CARLOS LIZAMA AND VICENTE VERGARA

(Communicated by Joseph A. Ball)

ABSTRACT. In this article we study uniform stability of resolvent families associated to an integral equation of convolution type. We give sufficient conditions for the uniform stability of the resolvent family in Hilbert and Banach spaces. Our main result can be viewed as a substantial generalization of the Gearhart-Greiner-Prüss characterization of exponential stability for strongly continuous semigroups.

1. INTRODUCTION

Let X be a Banach space. In this paper we are concerned with the study of the asymptotic behavior for the following integral Volterra equation of scalar type:

$$(1) \quad u(t) = \int_0^t a(t-s)Au(s)ds + f(t), \quad t \geq 0,$$

where A is a closed and linear operator with domain $D(A)$ dense in X , $a \in L^1_{loc}(\mathbb{R}_+)$ is a scalar kernel and $f \in W^{1,1}(\mathbb{R}_+, X)$.

Equation (1) has been extensively studied in recent years, mainly because of its applications in the theory of linear viscoelasticity. See, for instance, the monograph [5].

It is well known that equation (1) is well-posed if and only if it admits a *resolvent family*, that is, there is a strongly continuous family $S(t)$, $t \geq 0$, of bounded and linear operators defined in X , which commutes with A and satisfies the resolvent equation

$$(2) \quad S(t)x = x + \int_0^t a(t-s)AS(s)xds, \quad t \geq 0, \quad x \in D(A).$$

In particular, the resolvent family for (1) in the case $a(t) \equiv 1$ corresponds to the C_0 -semigroup generated by A .

Due to the special feature of a convolution in (1), it is appropriate to employ the Laplace transform for its study. Formally, the Laplace transform $H(\lambda) = \hat{S}(\lambda)$ of the resolvent family is represented by

$$(3) \quad H(\lambda) = (\lambda - \lambda \hat{a}(\lambda)A)^{-1}.$$

Received by the editors August 7, 2001 and, in revised form, September 5, 2002.

2000 *Mathematics Subject Classification*. Primary 45D05, 45N05; Secondary 47D06.

Key words and phrases. Uniform stability, resolvent family, abstract integral equations.

The authors were supported in part by FONDECYT Grant #1010675.

This work is part of the M.Sc. thesis for the second author.

The problem of finding necessary and sufficient conditions for the stability of resolvent families is, in general, difficult to handle, essentially due to the complex structure of (3). In 1992, Arendt and Prüss [1] studied the existence of $\lim_{t \rightarrow \infty} S(t) = P$ in various senses. In particular, they gave sufficient conditions for the strong stability of $S(t)$. This result extends to the well-known ABLP-theorem on stability for C_0 -semigroups (see [4]).

To our knowledge, there is no further study on the stability of resolvent. Our purpose in this article is to give some advances in the analysis of sufficient conditions in terms of (3) for the uniform stability of resolvent families in Hilbert and Banach spaces. The arguments used in [5, Theorem 10.2 and Theorem 10.5] to prove integrability give tools to prove two general results on uniform stability of resolvents. The first one (Theorem 1) concerning the hyperbolic case and valid in Hilbert spaces, generalizes the Gearhart-Greiner-Prüss characterization of exponential stability for strongly continuous semigroups. The second one (Theorem 2), is concerned with the parabolic case and is true in general Banach spaces.

Our basic assumption to get the above-mentioned results is only 1-regularity of the kernel $a(t)$. We remark that in general it is more difficult to prove integrability properties than stability for resolvents. However, the kernel $a(t)$ need not have the property that $1/\lambda\hat{a}(\lambda)$ is locally analytic and therefore the results in [5, Theorems 10.2 and Theorem 10.5] cannot be applied. We show in this paper that this condition is not needed for stability of resolvents.

Some immediate consequences of our results in the study of the asymptotic behavior of (1) can be obtained; for instance, suppose $f \in W^{1,1}(\mathbb{R}_+; X)$. Then, by the variation of parameters formula, we can conclude that $\|u(t)\| \rightarrow 0$ as $t \rightarrow \infty$.

Another application is the connection between the solutions of (1) and the solutions of the equation on the line

$$(4) \quad v(t) = \int_0^\infty a(s)Av(t-s)ds + g(s), \quad t \in \mathbb{R},$$

where $g \in W^{1,1}(\mathbb{R}; X)$. If (1) admits a resolvent $S(t)$ which is uniformly stable and uniformly integrable (see Corollary 3), then the solution of (4) is given by

$$v(t) = \int_0^\infty S(t)\dot{g}(t-s)ds, \quad t \in \mathbb{R}.$$

Assume that $\|\dot{f}(t) - \dot{g}(t)\| \rightarrow 0$ as $t \rightarrow +\infty$. From the variation of parameters formula we have

$$u(t) - v(t) = S(t)f(0) + \int_0^t S(\tau)[\dot{f}(t-\tau) - \dot{g}(t-\tau)]d\tau - \int_t^\infty S(\tau)\dot{g}(t-\tau)d\tau.$$

Hence $\|u(t) - v(t)\| \rightarrow 0$ as $t \rightarrow \infty$. This shows that the solutions $u(t)$ of (1) and $v(t)$ of (4) are asymptotic to each other as $t \rightarrow \infty$.

2. CONDITIONS FOR UNIFORM STABILITY

Recall that a resolvent family $\{S(t)\}_{t \geq 0}$ defined in a Banach space X is called *uniformly stable* if

$$\lim_{t \rightarrow +\infty} \|S(t)\| = 0.$$

In the following, we will suppose that $a \in L^1_{loc}(\mathbb{R}_+)$ is of subexponential growth, that is, $\int_0^\infty e^{-\varepsilon t} |a(t)| dt < \infty$, for all $\varepsilon > 0$. In this situation, the Laplace transform, $\hat{a}(\lambda)$, exists for all $Re\lambda > 0$.

We also recall that $a(t)$ is called k -regular ($k \in \mathbb{N}$) if there is a constant $c > 0$ such that

$$|\lambda^n \hat{a}^{(n)}(\lambda)| \leq c |\hat{a}(\lambda)|$$

for all $\operatorname{Re}\lambda > 0$ and $1 \leq n \leq k$.

Observe that if $a \in L^1_{loc}(\mathbb{R}_+)$ is of subexponential growth and 1-regular, then $\hat{a}(is) := \lim_{\lambda \rightarrow is} \hat{a}(\lambda)$ exists for all $s \neq 0$. Moreover, $\hat{a}(\lambda) \neq 0$ for $\operatorname{Re}\lambda \geq 0, \lambda \neq 0$ (see [5], Lemma 8.1).

Theorem 1. *Suppose $a(t) \in L^1_{loc}(\mathbb{R}_+)$ is 1-regular; assume that (1) admits a resolvent $S(t)$ with finite growth bound $\omega_0(S) < \infty$ in a Hilbert space H , and the following conditions:*

- (H1) $\frac{1}{\hat{a}(\lambda)} \in \rho(A)$ for all $\operatorname{Re}\lambda \geq 0, \lambda \neq 0$.
- (H2) $\lambda \hat{a}(\lambda) \rightarrow a(\infty) \neq 0$ as $\lambda \rightarrow 0$ and $0 \in \rho(A)$.
- (H3) $H(\lambda)$ is uniformly bounded in $\mathbb{C}_+ := \{\lambda \in \mathbb{C} : \operatorname{Re}\lambda > 0\}$.

Then $S(t)$ is uniformly stable.

Proof. By hypothesis there are constants $M > 0$ and $\omega_0 \in \mathbb{R}$ such that $\|S(t)\| \leq Me^{\omega_0 t}$. We may suppose that $\omega_0 \geq 0$. Let $\omega > \omega_0 + 1$ be given and define $R(t) := e^{-\omega t} S(t)$. Then $\|R(t)\| \leq Me^{-(\omega - \omega_0)t}$. Let $x \in H$ be fixed, and observe that $\chi_{[0, +\infty)}(\cdot)R(\cdot)x$ is in $L^2(\mathbb{R}; H)$, where $\chi_{[0, +\infty)}(\cdot)$ denotes the characteristic function. In fact,

$$\begin{aligned} \|\chi_{[0, +\infty)}(\cdot)R(\cdot)x\|_2^2 &= \int_0^\infty \|R(t)x\|^2 dt \leq M^2 \int_0^\infty (e^{-(\omega - \omega_0)t} \|x\|)^2 dt \\ &\leq \frac{M^2 \cdot \|x\|^2}{2(\omega - \omega_0)}; \end{aligned}$$

hence

$$\|\chi_{[0, +\infty)}(\cdot)R(\cdot)x\|_2 \leq \frac{M \cdot \|x\|}{\sqrt{2} \sqrt{\omega - \omega_0}}.$$

Because H is a Hilbert space, the Plancherel theorem shows us that the Fourier transform \mathcal{F} satisfies $\|\mathcal{F}f\|_2 = \sqrt{2\pi} \|f\|_2$ for all $f \in L^2(\mathbb{R}, H)$. On the other hand, because $S(t)$ is an exponentially bounded resolvent for (1), its Laplace transform $\hat{S}(\lambda)$ is then well-defined, holomorphic and satisfies $H(\lambda) = \hat{S}(\lambda)$ for all $\operatorname{Re}\lambda > 0$. Hence, we have for all $x \in H$ and $s \in \mathbb{R}$,

$$\begin{aligned} H(\omega + is)x &= \hat{S}(\omega + is)x = \int_0^\infty e^{-(\omega + is)t} S(t)x dt = \int_0^\infty e^{-ist} e^{-\omega t} S(t)x dt \\ &= \int_0^\infty e^{-ist} R(t)x dt = \int_{-\infty}^{+\infty} e^{-ist} \chi_{[0, +\infty)}(t) R(t)x dt \\ &= \mathcal{F}(\chi_{[0, +\infty)}(\cdot)R(\cdot)x)(s). \end{aligned}$$

It follows from the Plancherel theorem that $H(\omega + i \cdot)x \in L^2(\mathbb{R}; H)$ and

$$(5) \quad \|H(\omega + i \cdot)x\|_2 = \sqrt{2\pi} \cdot \|\chi_{[0, +\infty)}(\cdot)R(\cdot)x\|_2 \leq M \cdot \sqrt{\frac{\pi}{\omega - \omega_0}} \cdot \|x\|.$$

Writing $H(\lambda) = (\lambda - \lambda \hat{a}(\lambda)A)^{-1} = \frac{1}{\lambda \hat{a}(\lambda)} (\frac{\lambda}{\lambda \hat{a}(\lambda)} - A)^{-1}$ we observe that $\lim_{\lambda \rightarrow 0} H(\lambda)$ exist in $B(H)$ due to (H1) and (H2). Hence, from 1-regularity of $a(t)$, conditions (H1) and (H3) and the Banach-Steinhaus theorem we obtain that $H(i\rho)$ is bounded,

for each $\rho \in \mathbb{R}$. It follows from the uniform boundedness principle that $H(\lambda)$ is also uniformly bounded in the imaginary axis $i\mathbb{R}$.

On the other hand, from the identity

$$\lambda H(\lambda) - \lambda \hat{a}(\lambda) A H(\lambda) = I$$

valid for all $Re\lambda \geq 0$, $\lambda \neq 0$, we obtain

$$(6) \quad H(i\rho)x = H(\omega + i\rho)x + \frac{\omega}{\omega + i\rho} H(i\rho)x + h(\rho)H(i\rho)\left(H(\omega + i\rho)x - \frac{x}{\omega + i\rho}\right)$$

for all $\rho \neq 0$, where $h(\rho) = i\rho\left(\frac{\hat{a}(i\rho)}{\hat{a}(\omega + i\rho)} - 1\right)$.

It follows from the proof of Theorem 10.5 in [5] that $h(\rho)$ is bounded for $|\rho| \geq 1$.

Choose a function $\varphi(\rho)$ in $C_0^\infty(\mathbb{R})$, defined by $\varphi(\rho) = 1$ for $|\rho| < 1$ and $\varphi(\rho) = 0$ for $|\rho| \geq 2$. Define $\psi(\rho) = 1 - \varphi(\rho)$ for all $\rho \in \mathbb{R}$. Then using the uniform boundedness of $H(i\cdot)$ in \mathbb{R} and (5) in (6), we conclude that $\psi(\cdot)H(i\cdot)x \in L^2(\mathbb{R}; H)$ and

$$\begin{aligned} \|\psi(\cdot)H(i\cdot)x\|_2^2 &= \int_{-\infty}^{+\infty} \|\psi(\rho)H(i\rho)x\|^2 d\rho \\ &= \int_{|\rho| \geq 2} \|H(i\rho)x\|^2 d\rho + \int_{1 \leq |\rho| \leq 2} \|\psi(\rho)H(i\rho)x\|^2 d\rho \\ &\leq M_0 \cdot \|x\|^2. \end{aligned}$$

Analogously, we can prove that $H(\omega + i\cdot)^*x \in L^2(\mathbb{R}; H)$ and following the same argument as above we conclude that $\psi(\cdot)H(i\cdot)^*x \in L^2(\mathbb{R}, H)$.

By Parseval's theorem, there exists a function $u \in L^2(\mathbb{R}; H)$ such that

$$\mathcal{F}(u(\cdot))(\rho) = \psi(\rho)H(i\rho)x \text{ for a.a. } \rho \in \mathbb{R}.$$

It follows that

$$\begin{aligned} \mathcal{F}(u(\cdot))'(\rho) &= \psi'(\rho)H(i\rho)x + i\psi(\rho)H'(i\rho)x \\ &= \psi'(\rho)H(i\rho)x + i\psi(\rho)\left(-\frac{H(i\rho)}{i\rho}x + H(i\rho)i\rho\frac{\hat{a}'(i\rho)}{\hat{a}(i\rho)}\left(H(i\rho)x - \frac{x}{i\rho}\right)\right), \end{aligned}$$

hence, by 1-regularity of $a(t)$ and the fact that $\psi(\cdot)H(i\cdot)x$, $\psi(\cdot)H(i\cdot)^*x^*$ are in $L^2(\mathbb{R}; H)$ for each $x, x^* \in H$, we get

$$(7) \quad \int_{-\infty}^{+\infty} |\langle \mathcal{F}(u(\cdot))'(\rho), x^* \rangle| d\rho \leq M_0 \cdot \|x\| \cdot \|x^*\|.$$

On the other hand, again from the uniform boundedness of $H(i\cdot)$ in \mathbb{R} we have that for each $t > 0$

$$(8) \quad \begin{aligned} S_0(t) &:= \int_{-\infty}^{+\infty} \varphi(\rho)H(i\rho)e^{i\rho t} d\rho \\ &= \int_{-2}^2 \varphi(\rho)H(i\rho)e^{i\rho t} d\rho \in B(H). \end{aligned}$$

Hence, by the Riemman-Lebesgue lemma it follows that $S_0(t) \rightarrow 0$ in $B(H)$ as $t \rightarrow +\infty$.

Finally, for $x, x^* \in H$ we have that

$$\begin{aligned} \langle S(t)x, x^* \rangle &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} \langle H(i\rho)x, x^* \rangle e^{i\rho t} d\rho \\ &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} \langle \varphi(\rho)H(i\rho)x, x^* \rangle e^{i\rho t} d\rho \\ &\quad + \frac{1}{2\pi} \int_{-\infty}^{+\infty} \langle \psi(\rho)H(i\rho)x, x^* \rangle e^{i\rho t} d\rho. \end{aligned}$$

Integrating by parts in the second integral, we get

$$\begin{aligned} \langle S(t)x, x^* \rangle &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} \langle \varphi(\rho)H(i\rho)x, x^* \rangle e^{i\rho t} d\rho \\ &\quad + \frac{1}{2\pi it} \int_{-\infty}^{+\infty} \langle (\psi(\rho)H(i\rho))'x, x^* \rangle e^{i\rho t} d\rho \\ &= \frac{1}{2\pi} \langle S_0(t)x, x^* \rangle + \frac{1}{2\pi it} \int_{-\infty}^{+\infty} \langle \mathcal{F}(u(\cdot))'(\rho), x^* \rangle e^{i\rho t} d\rho. \end{aligned}$$

It follows from (7) and (8) that

$$|\langle S(t)x, x^* \rangle| \leq \frac{1}{2\pi} \|S_0(t)\| \cdot \|x\| \cdot \|x^*\| + \frac{1}{2\pi t} M_0 \cdot \|x\| \cdot \|x^*\|.$$

Therefore, $\|S(t)\| \leq \frac{1}{2\pi} \|S_0(t)\| + \frac{1}{2\pi t} M_0$, from which we obtain the result.

Corollary 1. *Suppose that the equation $u = a * Au + f$, where $a(t)$ is 1-regular, admits a resolvent $S(t)$ with finite growth bound $\omega_0(S) < \infty$ in a Hilbert space H . If $S(t)$ is strongly integrable, then $S(t)$ is uniformly stable.*

Proof. The proof follows from Theorem 1 and Theorem 10.5 in [5].

The special case $a(t) \equiv 1$ gives us the following result on stability of C_0 -semigroups due to Gearhart, Greiner and Prüss (see [2], Theorem 1.11).

Corollary 2. *Let A be the generator of a C_0 -semigroup $\{T(t)\}_{t \geq 0}$ with finite growth bound $\omega_0(S) < \infty$ defined in a Hilbert space H . The following conditions are equivalent:*

- (a) *The semigroup $T(t)$ is uniformly stable.*
- (b) $\{\lambda \in \mathbb{C} : \operatorname{Re} \lambda \geq 0\} \subset \rho(A)$ and $\sup_{\operatorname{Re} \lambda > 0} \|R(\lambda; A)\| < \infty$.

Proof. Assume (b). Is easy to see that $a(t) \equiv 1$ is 1-regular and the hypotheses (H1)-(H3) are clearly satisfied. Hence (a) follows by Theorem 1.

Assume (a). Since the semigroup is uniformly stable, we conclude from ([2], Proposition 1.2) that it is also exponentially stable, that is, there is a constant $\omega_0 > 0$ such that $\|S(t)\| \leq M e^{-\omega_0 t}$. Hence $\{\lambda \in \mathbb{C} : \operatorname{Re} \lambda \geq 0\} \subseteq \rho(A)$ and $\|R(\tau + is; A)\| \leq \frac{M}{\tau + \omega_0} < \frac{M}{\omega_0}$ for all $\tau \geq 0$ and $s \in \mathbb{R}$ from which we obtain (b).

We recall the following:

Definition 1. Equation (1) is called *parabolic* if the following conditions are satisfied:

- (P1) $\hat{a}(\lambda) \neq 0$, $1/\hat{a}(\lambda) \in \rho(A)$ for all $Re\lambda > 0$.
(P2) There exists a constant $M \geq 1$ such that

$$(9) \quad \|H(\lambda)\| \leq \frac{M}{|\lambda|} \text{ for all } Re\lambda > 0.$$

Theorem 2. *Let X be a Banach space. Suppose that $a(t)$ is 1-regular and (H2) of Theorem 1 holds. If (1) is parabolic, then (1) admits a resolvent $S(t)$ uniformly stable.*

Proof. From parabolicity and 1-regularity of $a(t)$ we have by Theorem 3.1 in [5] the existence of a resolvent family $S(t)$, continuous and bounded for $t \geq 0$. Hence $S(\cdot) \in L^1_{loc}(\mathbb{R}_+, B(X))$ and moreover by the identity $\lambda H(\lambda) - \lambda \hat{a}(\lambda) A H(\lambda) = I$, for $Re\lambda > 0$, we have that

$$\begin{aligned} H'(\lambda) &= -\frac{H(\lambda)}{\lambda} + \hat{a}'(\lambda) A (I - \hat{a}(\lambda) A)^{-1} H(\lambda) \\ &= -\frac{H(\lambda)}{\lambda} + \frac{\hat{a}'(\lambda)}{\hat{a}(\lambda)} \cdot \hat{a}(\lambda) A \cdot \lambda H(\lambda) \cdot H(\lambda) \\ &= -\frac{H(\lambda)}{\lambda} + \frac{\hat{a}'(\lambda)}{\hat{a}(\lambda)} \cdot (\lambda H(\lambda) - I) \cdot H(\lambda). \end{aligned}$$

It follows from (9) and 1-regularity of $a(t)$ that $\|\lambda H'(\lambda)\| \leq M_1 \|H(\lambda)\|$, for $Re\lambda > 0$. So, we are in the conditions of Lemma 8.1 in [5]. It follows that $H(\lambda)$ admits a $B(X)$ -continuous extension to $\overline{\mathbb{C}}_+ \setminus \{0\}$. We will prove that $H(\lambda)$ is, in addition, continuous at $\lambda = 0$. Indeed, from (H2) and $0 \in \rho(A)$ we have that

$$H(0) = \lim_{\lambda \rightarrow 0} H(\lambda) = \lim_{\lambda \rightarrow 0} \frac{1}{\lambda \hat{a}(\lambda)} \left(\frac{\lambda}{\lambda \hat{a}(\lambda)} - A \right)^{-1} = \frac{-A^{-1}}{a(\infty)}$$

exists in $B(X)$. Hence $\|H(\lambda)\| \leq \frac{M_2}{1+|\lambda|}$ for $Re\lambda \geq 0$. It follows that $H(\lambda)$ is uniformly bounded on $\overline{\mathbb{C}}_+$. Let $x \in X$ and $x^* \in X^*$ be fixed. We observe that $\langle H(\omega + i\cdot)x, x^* \rangle \in L^2(\mathbb{R})$ for all $\omega \geq 0$. It follows from Lemma 6.1 in [3] that $\langle S(t)x, x^* \rangle \in L^2(\mathbb{R}_+)$, therefore by L^2 -theory we have the next representation,

$$(10) \quad \langle S(t)x, x^* \rangle = \frac{1}{2\pi} \int_{-\infty}^{\infty} \langle H(i\rho)x, x^* \rangle e^{i\rho t} d\rho, \text{ a.a. } t \in \mathbb{R}_+.$$

Now we consider $N \in \mathbb{N}$ fixed and $\varphi \in C_0^\infty(\mathbb{R})$ defined by $\varphi(\rho) = 1$ if $|\rho| \leq N$, 0 if $|\rho| \geq N+2$ and $0 \leq \varphi \leq 1$ in another case. Then by (10) we have that

$$(11) \quad \begin{aligned} \langle S(t)x, x^* \rangle &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \langle \varphi(\rho) H(i\rho)x, x^* \rangle e^{i\rho t} d\rho \\ &\quad + \frac{1}{2\pi} \int_{-\infty}^{\infty} \langle (1 - \varphi(\rho)) H(i\rho)x, x^* \rangle e^{i\rho t} d\rho. \end{aligned}$$

Hence, integrating by parts in the second integral in (11), this becomes

$$(12) \quad \begin{aligned} \langle S(t)x, x^* \rangle &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \langle \varphi(\rho) H(i\rho)x, x^* \rangle e^{i\rho t} d\rho \\ &\quad + \frac{1}{2\pi i t} \int_{-\infty}^{\infty} \langle [(1 - \varphi(\rho)) H(i\rho)]' x, x^* \rangle e^{i\rho t} d\rho. \end{aligned}$$

It follows from boundedness of $H(\lambda)$ on $\overline{\mathbb{C}}_+$ that the first integral is absolutely integrable. For the second integral, using the estimate $\|\rho H'(i\rho)\| \leq M_2 \|H(i\rho)\|$ a.a. $\rho \in \mathbb{R}$ and the Cauchy-Schwarz and Hölder inequalities, it follows that it is also absolutely integrable. Therefore, x and x^* can be dropped in (12) and $S(t)$ can be written as

$$(13) \quad \begin{aligned} S(t) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \varphi(\rho) H(i\rho) e^{i\rho t} d\rho + \frac{1}{2\pi i t} \int_{-\infty}^{\infty} [(1 - \varphi(\rho)) H(i\rho)]' e^{i\rho t} d\rho \\ &:= S_1(t) + S_2(t). \end{aligned}$$

It follows from the Riemman-Lebesgue lemma that $S_1(t) \rightarrow 0$ as $t \rightarrow +\infty$. On the other hand, $S_2(t)$ satisfies an estimate of the form $\|S_2(t)\| \leq \frac{M_2}{t}$. It follows that $S(t) \rightarrow 0$ as $t \rightarrow +\infty$.

As an immediate consequence of Theorem 2 and Theorem 10.2 in [5], we obtain the following.

Corollary 3. *Suppose that $u = a * Au + f$ is parabolic in Banach space X , where $a(t)$ is 1-regular. If $S(t)$ is uniformly integrable, then $S(t)$ is uniformly stable.*

ACKNOWLEDGMENTS

The authors thank Professor J. Prüss for valuable suggestions in order to improve a previous version of this paper.

REFERENCES

- [1] W. Arendt and J. Prüss, *Vector-valued tauberian theorem and asymptotic behaviour of linear Volterra equations*, SIAM J. Math. Anal. **23** (2), (1992), 412–448. MR **92m**:47150
- [2] K. J. Engel and R. Nagel, *One-Parameter Semigroups for Linear Evolution Equations*, Springer-Verlag, Berlin, Heidelberg, New York, 2000. MR **2000i**:47075
- [3] G. Gripenberg, S. O. Londen, and O. J. Staffans. *Volterra Integral and Functional Equations*, Cambridge Univ. Press, Cambridge, 1990. MR **91c**:45003
- [4] J.M.A.A. van Neerven, *The Asymptotic Behaviour of Semigroups of Linear Operators*, Oper. Theory Adv. Appl. **88**, Birkhäuser-Verlag, 1996. MR **98d**:47001
- [5] J. Prüss, *Evolutionary Integral Equations and Applications*, Birkhäuser-Verlag, Basel-Boston-Berlin, 1993. MR **94h**:45010

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDAD DE SANTIAGO DE CHILE, CASILLA 307-CORREO 2, SANTIAGO, CHILE

E-mail address: `clizama@usach.cl`

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDAD DE SANTIAGO DE CHILE, CASILLA 307-CORREO 2, SANTIAGO, CHILE

E-mail address: `vvergara@usach.cl`