

ON TWO PROBLEMS OF ERDŐS AND HECHLER: NEW METHODS IN SINGULAR MADNESS

MENACHEM KOJMAN, WIESŁAW KUBIŚ, AND SAHARON SHELAH

(Communicated by Carl G. Jockusch, Jr.)

ABSTRACT. For an infinite cardinal μ , $\text{MAD}(\mu)$ denotes the set of all cardinalities of *nontrivial maximal almost disjoint families* over μ .

Erdős and Hechler proved in 1973 the consistency of $\mu \in \text{MAD}(\mu)$ for a singular cardinal μ and asked if it was ever possible for a singular μ that $\mu \notin \text{MAD}(\mu)$, and also whether $2^{\text{cf } \mu} < \mu \implies \mu \in \text{MAD}(\mu)$ for every singular cardinal μ .

We introduce a new method for controlling $\text{MAD}(\mu)$ for a singular μ and, among other new results about the structure of $\text{MAD}(\mu)$ for singular μ , settle both problems affirmatively.

1. INTRODUCTION

1.1. Background. Let μ be an infinite cardinal. A family of sets \mathcal{A} is μ -almost disjoint (μ -ad for short) if $|A| = \mu = |\bigcup \mathcal{A}|$ for every $A \in \mathcal{A}$ and $|A \cap B| < \mu$ for every distinct $A, B \in \mathcal{A}$. \mathcal{A} is *maximal μ -almost disjoint* (μ -mad) if there is no $C \subseteq \bigcup \mathcal{A}$ such that $\mathcal{A} \cup \{C\}$ is μ -almost disjoint; in this case we also say that \mathcal{A} is *mad in μ* . It is clear that every μ -almost disjoint family consisting of fewer than $\text{cf } \mu$ sets is mad in μ ; such a family will be called *trivial*. We denote by $\text{MAD}(\mu)$ the set of all cardinalities of nontrivial mad families in μ . A standard diagonalization argument shows that $\text{cf } \mu \notin \text{MAD}(\mu)$. Therefore, $\text{MAD}(\mu)$ is contained in the interval of cardinals $[\text{cf}(\mu)^+, 2^\mu]$.

W. W. Comfort asked (see [7]) under what conditions it follows that $\mu \in \text{MAD}(\mu)$ for a singular cardinal μ . P. Erdős and S. Hechler [7] proved that $\mu \in \text{MAD}(\mu)$ if $\lambda^{\text{cf } \mu} < \mu$ for every $\lambda < \mu$. Thus, if $2^{\aleph_0} < \aleph_\omega$, then the interval $[2^{\aleph_0}, \aleph_\omega]$ of cardinals is contained in $\text{MAD}(\aleph_\omega)$.

Erdős and Hechler asked in [7] whether it is consistent that $\mu \notin \text{MAD}(\mu)$ for some singular cardinal μ and, more concretely, whether Martin's axiom together with $2^{\aleph_0} > \aleph_\omega$ implies that $\aleph_\omega \notin \text{MAD}(\aleph_\omega)$. They also asked whether $2^{\text{cf } \mu} < \mu$ implies $\mu \in \text{MAD}(\mu)$ for singular cardinals μ other than \aleph_ω .

Received by the editors June 10, 2002 and, in revised form, September 10, 2002.

2000 *Mathematics Subject Classification*. Primary 03E10, 03E04, 03E17, 03E35; Secondary 03E55, 03E50.

Key words and phrases. Almost disjoint family, singular cardinal, bounding number, smooth pcf scales.

The first author's research partially supported by an Israeli Science Foundation grant no. 177/01.

The third author's research was supported by The Israel Science Foundation, Publication 793.

Both problems are settled affirmatively by the general results below on $\text{MAD}(\mu)$ for a singular μ .

1.2. Notation. Let $\mathfrak{a}_\mu = \min \text{MAD}(\mu)$, and let $\mathfrak{a} = \mathfrak{a}_{\aleph_0}$. For a singular μ it follows that $\text{MAD}(\text{cf } \mu) \subseteq \text{MAD}(\mu)$; therefore $\mathfrak{a}_\mu \leq \mathfrak{a}_{\text{cf } \mu}$.

A crucial role in the results is played by two *bounding numbers*: \mathfrak{b}_μ and $\mathfrak{b}_{\text{cf } \mu}$.

For every quasi-ordering (P, \leq) with no maximum, the *bounding number* $\mathfrak{b}(P, \leq)$ is the least cardinality of a subset of P with no upper bound. For a regular cardinal κ , let \mathfrak{b}_κ denote the bounding number of (κ^κ, \leq^*) , where $f \leq^* g$ means that $|\{i < \kappa : f(i) > g(i)\}| < \kappa$; let $\mathfrak{b} = \mathfrak{b}_{\aleph_0}$. It is well known that $\kappa < \mathfrak{b}_\kappa \leq \mathfrak{a}_\kappa$ for a regular cardinal κ (for $\kappa = \aleph_0$ see [6]; the general case is similar) and that under Martin's axiom, $\mathfrak{b} = 2^{\aleph_0}$.

Suppose that μ is a singular cardinal of cofinality κ and that $\langle \mu_i : i < \kappa \rangle$ is a strictly increasing sequence of regular cardinals with supremum μ . Standard diagonalization shows that $\mathfrak{b}(\prod_{i < \kappa} \mu_i, \leq^*) > \mu$. Denote by \mathfrak{b}_μ the *supremum of $\mathfrak{b}(\prod \mu_i, \leq^*)$ over all strictly increasing sequences of regular cardinals $\langle \mu_i : i < \kappa \rangle$ with supremum μ* .

Each of the following three relations is consistent with ZFC: $\mathfrak{b} < \mathfrak{b}_{\aleph_\omega}$, $\mathfrak{b} = \mathfrak{b}_{\aleph_\omega}$ and $\mathfrak{b} > \mathfrak{b}_{\aleph_\omega}$.

1.3. The results. We prove that for every singular cardinal μ :

- (1) $\mathfrak{a}_\mu \geq \min\{\mathfrak{b}_\mu, \mathfrak{b}_{\text{cf } \mu}\}$;
- (2) $\mathfrak{a}_\mu \leq \lambda < \mathfrak{b}_\mu \implies \lambda \in \text{MAD}(\mu)$.

Thus, if $\mathfrak{b}_{\text{cf } \mu} > \mu$, it follows from (1) that $\mathfrak{a}_\mu > \mu$, and hence $\mu \notin \text{MAD}(\mu)$; and if $\mathfrak{a}_{\text{cf } \mu} < \mu$, it follows from (2) that $\mu \in \text{MAD}(\mu)$. In particular,

- (b) $MA + 2^{\aleph_0} > \aleph_\omega \implies \aleph_\omega \notin \text{MAD}(\aleph_\omega)$,
- (a) $2^{\text{cf } \mu} < \mu \implies \mu \in \text{MAD}(\mu)$ for every singular μ ,

which, respectively, settle in the affirmative both problems of Erdős and Hechler from [7].

If one assumes the consistency of large cardinals, $\mathfrak{b}_{\aleph_\omega}$ can be shifted up arbitrarily high below \aleph_{ω_1} . Following this with a ccc forcing for controlling \mathfrak{b} proves the following:

- (3) for every regular $\lambda \in (\aleph_\omega, \aleph_{\omega_1})$ and regular uncountable $\theta \leq \lambda^+$, it is consistent that

$$\text{MAD}(\aleph_\omega) = [\theta, \lambda^+].$$

So, e.g., the following are consistent:

- $\text{MAD}(\aleph_\omega) = \{\aleph_1, \aleph_2, \dots, \aleph_{\omega+\beta+2} = 2^{\aleph_\omega}\}$ for an arbitrary $\beta < \omega_1$,
- $\text{MAD}(\aleph_\omega) = \{\aleph_{\omega+\beta+2}\}$ for an arbitrary $\beta < \omega_1$,
- $\text{MAD}(\aleph_\omega) = [\aleph_{\omega+\alpha+1}, \aleph_{\omega+\beta+2}]$ for arbitrary $\alpha \leq \beta < \omega_1$,

and so on.

We refer the reader to the comprehensive list of references in D. Monk's recent paper [12], in which maximal almost disjoint families are viewed as partitions of unity in the Boolean algebra $\mathcal{P}(\mu)/_{[\mu] < \mu}$.

1.4. Preliminary facts. We will use the following facts from [7]:

- (1) $\text{MAD}(\text{cf } \mu) \subseteq \text{MAD}(\mu)$, and
- (2) $\text{MAD}(\mu)$ is closed under singular suprema.

The latter fact is stated in [7] in a less general form. So we give a proof here.

Lemma 1.1. *Assume that $\lambda = \sup_{i < \theta} \lambda_i$, where $\{\lambda_i : i < \theta\} \subseteq \text{MAD}(\mu)$ and $\theta < \lambda$. Then $\lambda \in \text{MAD}(\mu)$.*

Proof. We may assume that $\theta \leq \lambda_0$. Let \mathcal{A} be a mad family in μ with $|\mathcal{A}| = \lambda_0$. Write $\mathcal{A} = \{A_i : i < \lambda_0\}$ and for each $i < \theta$ choose a mad family \mathcal{B}_i with $\bigcup \mathcal{B}_i = A_i$ and $|\mathcal{B}_i| = \lambda_i$. Set

$$\mathcal{C} = \bigcup_{i < \theta} \mathcal{B}_i \cup \{A_j : \theta \leq j < \lambda_0\}.$$

Then $|\mathcal{C}| = \lambda$ and \mathcal{C} is mad in μ . \square

The following fact will also be used in some proofs.

Lemma 1.2. *Let $\kappa = \text{cf } \mu$, and let \mathcal{A} be a μ -almost disjoint family of size κ . Then there exists a mad family $\mathcal{A}' \supseteq \mathcal{A}$ such that $|\mathcal{A}'| = \mathfrak{a}_\mu$ and $\bigcup \mathcal{A}' = \bigcup \mathcal{A}$.*

Proof. Fix a μ -mad family \mathcal{B} with $|\mathcal{B}| = \mathfrak{a}_\mu$. Choose $\mathcal{B}_0 = \{B_i : i < \kappa\} \subseteq \mathcal{B}$. Let $X = \bigcup_{i < \kappa} B_i$ and define

$$\mathcal{B}' = \{B \cap X : X \in \mathcal{B} \setminus \mathcal{B}_0 \text{ \& } |X \cap B| = \mu\}.$$

Let $\langle A_i : i < \kappa \rangle$ be a one-to-one enumeration of \mathcal{A} . Define a bijection $f : \bigcup \mathcal{A} \rightarrow X$ so that $f[A_i \setminus \bigcup_{j < i} A_j] = B_i \setminus \bigcup_{j < i} B_j$. Finally, set $\mathcal{A}' = \mathcal{A} \cup \{f^{-1}[B] : B \in \mathcal{B}'\}$. Observe that \mathcal{A}' is mad and $|\mathcal{A}'| = \mathfrak{a}_\mu$. \square

2. INEQUALITIES

From now on, μ will always denote a singular cardinal whose cofinality is denoted by κ .

2.1. Bounding numbers and madness in singular cardinals.

Theorem 2.1. *For every singular cardinal μ ,*

$$(1) \quad \mathfrak{a}_\mu \geq \min\{\mathfrak{b}_\mu, \mathfrak{b}_{\text{cf } \mu}\}.$$

Proof. Let $\kappa = \text{cf } \mu$. Suppose to the contrary that $\mathfrak{a}_\mu < \min\{\mathfrak{b}_\mu, \mathfrak{b}_\kappa\}$, and fix a strictly increasing sequence of regular cardinals $\langle \mu_i : i < \kappa \rangle$ with supremum μ such that $\mathfrak{b}(\prod_{i < \kappa} \mu_i, \leq^*) > \mathfrak{a}_\mu$.

Let $\mathcal{A} = \{\{i\} \times \mu : i < \kappa\}$. By Lemma 1.2, there exists a family $\mathcal{B} \subseteq [\kappa \times \mu]^\mu$ such that $\mathcal{B} \cup \mathcal{A}$ is mad in μ , $\mathcal{B} \cap \mathcal{A} = \emptyset$ and $|\mathcal{B}| = \mathfrak{a}_\mu$.

For each $B \in \mathcal{B}$, define a function $f_B : \kappa \rightarrow \kappa$ by $f_B(i) = \min\{j < \kappa : |B \cap (\{i\} \times \mu)| < \mu_j\}$. This function is well defined, since $|B \cap (\{i\} \times \mu)| < \mu$ for each $i < \kappa$.

Since $|\mathcal{B}| = \mathfrak{a}_\mu < \mathfrak{b}_\kappa$, there exists a function $f : \kappa \rightarrow \kappa$ so that $f_B <^* f$ for all $B \in \mathcal{B}$. Without loss of generality, we may assume that f is strictly increasing.

For each $B \in \mathcal{B}$, for all but boundedly many $i < \kappa$, it follows that $\sup\{\alpha < \mu_{f(i)} : (i, \alpha) \in B\} < \mu_{f(i)}$. Let $g_B(i)$ be defined by

$$g_B(i) = \begin{cases} 0 & \text{if } \sup\{\alpha < \mu_{f(i)} : (i, \alpha) \in B\} = \mu_{f(i)}, \\ \sup\{\alpha < \mu_{f(i)} : (i, \alpha) \in B\} & \text{otherwise.} \end{cases}$$

For each $B \in \mathcal{B}$ the function g_B belongs to $\prod_{i < \kappa} \mu_{f(i)}$. Since

$$\mathfrak{b}(\prod_{i < \kappa} \mu_{f(i)}, \leq^*) \geq \mathfrak{b}(\prod_{i < \kappa} \mu_i) > \mathfrak{a}_\mu,$$

we can fix a function $g \in \prod_{i < \kappa} \mu_{f(i)}$ so that $g_B <^* g$ for all $B \in \mathcal{B}$.

Define

$$C = \bigcup_{i < \kappa} \{i\} \times [g(i), \mu_{f(i)}).$$

Clearly, $|C| = \mu$. For each $B \in \mathcal{B}$ there exists $j_B < \kappa$ such that $g_B(i) < g(i)$ for all $i > j_B$. This implies that $\{i\} \times [g(i), \mu_{f(i)})$ is disjoint from B for all $i > j_B$. Hence $|B \cap C| \leq \mu_{f(j_B)} < \mu$. Clearly, $|C \cap (\{i\} \times \mu)| \leq \mu_{f(i)} < \mu$ for all $i < \kappa$; so $\mathcal{A} \cup \mathcal{B} \cup \{C\}$ is μ -almost disjoint, contrary to the maximality of $\mathcal{A} \cup \mathcal{B}$. \square

A positive answer to the first question of Comfort, Erdős and Hechler follows now as a corollary:

Corollary 2.2. *If Martin's Axiom holds and $2^{\aleph_0} > \mu > \text{cf } \mu = \aleph_0$, then $\mu \notin \text{MAD}(\mu)$.*

2.2. Between \mathfrak{a}_μ and \mathfrak{b}_μ . In this section we shall show that $\text{MAD}(\mu)$ contains the interval of cardinals $[\mathfrak{a}_\mu, \mathfrak{b}_\mu)$ and even $[\mathfrak{a}_\mu, \mathfrak{b}_\mu]$ in the case that \mathfrak{b}_μ is a successor of a regular cardinal.

Theorem 2.3. *For every singular cardinal μ and every cardinal λ ,*

$$(2) \quad \mathfrak{a}_\mu \leq \lambda < \mathfrak{b}_\mu \implies \lambda \in \text{MAD}(\mu).$$

If \mathfrak{b}_μ is a successor of a regular cardinal, then $\mathfrak{a}_\mu \leq \mathfrak{b}_\mu \implies \mathfrak{b}_\mu \in \text{MAD}(\mu)$.

To prove the theorem it suffices, by Lemma 1.1, to show that every regular $\lambda \in [\mathfrak{a}_\mu, \mathfrak{b}_\mu)$ belongs to $\text{MAD}(\mu)$.

The proof of this will now be divided into two cases. First we prove that every regular $\mathfrak{a}_\mu < \lambda < \mu$ belongs to $\text{MAD}(\mu)$. The proof in this case does not require any specialized techniques. Then we prove the same for regular $\mu < \lambda < \mathfrak{b}_\mu$ and for \mathfrak{b}_μ itself when it is the successor of a regular cardinal. In this case the proof requires some machinery from pcf theory.

Despite the technical differences between both proofs, they are similar, and could, in fact, be combined to a single proof. Both follow the same scheme of gluing together λ different μ -mad families, each of size \mathfrak{a}_μ , to a single μ -mad family of size λ . In the case $\lambda < \mu$, a simple presentation of μ as a disjoint union of λ parts works; in the second part we need to rely on smooth pcf scales to get a presentation of μ as an *almost increasing* and *continuous* union of length λ of sets of size μ .

2.2.1. The case $\lambda < \mu$.

Lemma 2.4. *Suppose $\mu > \text{cf } \mu = \kappa$. Then for every regular cardinal λ ,*

$$\mathfrak{a}_\mu \leq \lambda < \mu \implies \lambda \in \text{MAD}(\mu).$$

Proof. Suppose λ is regular and $\mathfrak{a}_\mu \leq \lambda < \mu$. Since $\mathfrak{a}_\mu > \kappa = \text{cf } \mu$, $\lambda > \kappa$.

Fix a strictly increasing sequence of regular cardinals $\langle \mu_i : i < \kappa \rangle$ such that $\sup_{i < \kappa} \mu_i = \mu$ and $\lambda < \mu_0$. We will work in $\mu \times \lambda$ instead of μ . Let $S = \{\delta < \lambda : \text{cf } \delta = \kappa\}$. For each $\delta \in S$ fix a strictly increasing, continuous sequence $D_\delta = \langle \gamma_i^\delta : i < \kappa \rangle$ with limit δ such that $\gamma_0^\delta = 0$. Define

$$F_j^\delta = \bigcup \{ \mu \times \{ \beta \} : \gamma_j^\delta \leq \beta < \gamma_{j+1}^\delta \}.$$

Thus $\mathcal{F}_\delta = \{F_j^\delta : j < \kappa\}$ is a disjoint family of sets, each set of size μ , which covers $\mu \times \delta$. Let $\mathcal{A}_\delta \subseteq [\mu \times \delta]^\mu$ be such that $\mathcal{A}_\delta \cup \mathcal{F}_\delta$ is mad in $\mu \times \delta$, $\mathcal{A}_\delta \cap \mathcal{F}_\delta = \emptyset$ and $|\mathcal{A}_\delta| = \mathfrak{a}_\mu$.

Define

$$\mathcal{B} = \{\mu \times \{\alpha\} : \alpha < \lambda\} \cup \bigcup_{\delta \in S} \mathcal{A}_\delta.$$

Then $|\mathcal{B}| = \lambda$ and $\mathcal{B} \subseteq [\mu \times \lambda]^\mu$. We will show that \mathcal{B} is μ -mad.

First, observe that \mathcal{B} is almost disjoint: clearly each element of \mathcal{A}_δ is almost disjoint from any set of the form $\mu \times \{\alpha\}$, because if $\alpha < \delta$, then $\mu \times \{\alpha\} \subseteq F_j^\delta$ for $j < \kappa$ such that $\gamma_j^\delta \leq \alpha < \gamma_{j+1}^\delta$. Finally, consider $A_i \in \mathcal{A}_{\delta_i}$, $i < 2$, with $\delta_0 < \delta_1$. Then $A_0 \subseteq \bigcup_{j < j_0} F_j^{\delta_1}$, where $j_0 < \kappa$ is such that $\delta_0 < \gamma_{j_0}^{\delta_1}$. Thus $|A_0 \cap A_1| < \mu$.

To see that \mathcal{B} is mad, fix an arbitrary $Z \in [\mu \times \lambda]^\mu$. There exists a sequence $\langle \alpha_i : i < \kappa \rangle$ in λ such that

$$|Z \cap (\mu \times \{\alpha_i\})| \geq \mu_i.$$

If $|\{\alpha_i : i < \kappa\}| < \kappa$, then $|Z \cap (\mu \times \{\alpha\})| = \mu$ for some α . So suppose that $|Z \cap (\mu \times \{\alpha_i\})| < \mu$ for every $i < \kappa$. Taking a subsequence, we may assume that $\langle \alpha_i : i < \kappa \rangle$ is strictly increasing. Let δ be its supremum. By regularity of λ , $\delta \in S$ and therefore $Z \in [\mu \times \delta]^\mu$. Shrinking Z if necessary, assume that $Z \subseteq \bigcup_{i < \kappa} \mu \times \{\alpha_i\}$. Then $|Z \cap F_j^\delta| < \mu$ for every $j < \kappa$. Thus, $|Z \cap A| = \mu$ for some $A \in \mathcal{A}_\delta$. This completes the proof. \square

Corollary 2.5. *Let $\mu > \text{cf } \mu = \kappa$. If $\mathfrak{a}_\kappa \leq \mu$, then $[\mathfrak{a}_\kappa, \mu] \subseteq \text{MAD}(\mu)$. In particular, if $2^\kappa < \mu$, then $\mu \in \text{MAD}(\mu)$.*

Corollary 2.5 answers affirmatively the second question of Erdős and Hechler in [7].

2.2.2. The case $\lambda > \mu$. A (μ, λ) -scale, for a regular cardinal $\lambda > \mu$, is a sequence $\bar{f} = \langle f_\alpha : \alpha < \lambda \rangle \subseteq \prod_{i < \kappa} \mu_i$ such that $\langle \mu_i : i < \kappa \rangle$ is a strictly increasing sequence of regular cardinals with limit μ , and so that $\alpha < \beta < \lambda \implies f_\alpha <^* f_\beta$ and for every $g \in \prod_{i < \kappa} \mu_i$ there is $\alpha < \lambda$ with $g <^* f_\alpha$. The relation $f <^* g$ means that the set $\{i < \kappa : f(i) \geq g(i)\}$ is bounded in κ . When μ is fixed, “ (μ, λ) -scale” will be abbreviated by “ λ -scale”. A λ -scale \bar{f} is *smooth* if for every $\delta < \lambda$ with $\text{cf } \delta > \kappa$, the sequence $\bar{f} \upharpoonright \delta = \langle f_\alpha : \alpha < \delta \rangle$ is cofinal in $(\prod_{i < \kappa} f_\delta(i), <^*)$. In this case we say that f_δ is an *exact upper bound* of $\bar{f} \upharpoonright \delta$. We will denote by $[f, g]$ the set $\{(i, \alpha) : i < \kappa \wedge f(i) \leq \alpha < g(i)\}$.

The proof in the present case goes through two steps. First, it is shown that whenever a smooth (μ, λ) -scale exists and $\mathfrak{a}_\mu < \lambda$, it follows that $\lambda \in \text{MAD}(\mu)$. Then it is shown that for every $\mu < \lambda < \mathfrak{b}_\mu$ there is a smooth (μ, λ) -scale and that in case \mathfrak{b}_μ is a successor of a regular cardinal there is also a smooth (μ, \mathfrak{b}_μ) -scale.

Lemma 2.6. *Assume $\lambda > \mu > \text{cf } \mu = \kappa$ and there exists a smooth (μ, λ) -scale. If $\mathfrak{a}_\mu \leq \lambda$, then $\lambda \in \text{MAD}(\mu)$.*

Proof. Suppose there exists a smooth λ -scale $\langle g_\xi : \xi < \lambda \rangle \subseteq \prod_{i < \kappa} \mu_i$. Let $S = \{\delta < \lambda : \text{cf } \delta = \kappa\}$, and for each $\delta \in S$ fix a strictly increasing, continuous, sequence $\langle \gamma_i^\delta : i < \kappa \rangle$ with limit δ such that $\gamma_0^\delta = 0$ and put $D_\delta = \{\gamma_i^\delta : i < \kappa\}$.

By induction on $\xi < \lambda$ we construct a smooth λ -scale $\bar{f} = \langle f_\xi : \xi < \lambda \rangle \subseteq \prod_{i < \kappa} \mu_i$ that satisfies the following two conditions:

- (1) If $\delta < \lambda$ is a limit and $\text{cf } \delta \leq \kappa$, then $f_\delta(i) = \sup_{\xi \in D_\delta} f_\xi(i)$.
- (2) For each $\xi < \lambda$ the set $[f_\xi, f_{\xi+1}] = \{(i, \alpha) : f_\xi(i) \leq \alpha < f_{\xi+1}(i)\}$ has cardinality μ .

By induction on $\xi < \lambda$ we define an increasing and continuous sequence of ordinals $\zeta(\xi) < \lambda$ and a $<^*$ -increasing sequence of functions $f_\xi \in \prod_{i < \kappa} \mu_i$ so that $f_\xi = g_{\zeta(\xi)}$ for all $\xi < \lambda$ *except* when ξ is limit of cofinality $\leq \kappa$. Then $\bar{f} := \langle f_\xi : \xi < \lambda \rangle$ will be a smooth λ -scale as required.

At a limit stage ξ of cofinality $\leq \kappa$, let $\zeta(\xi) = \bigcup_{\xi' < \xi} \zeta(\xi')$ and use condition (1) to define f_ξ ; at successor $\xi + 1$ choose $\zeta(\xi + 1)$ so that $\max\{f_\xi, g_{\zeta(\xi)}\} <^* g_{\zeta(\xi+1)}$ and (2) holds, and let $f_{\xi+1} = g_{\zeta(\xi+1)}$. Suppose now that ξ is a limit of cofinality $> \kappa$. By the smoothness of \bar{g} , and since $\langle g_{\zeta(\xi')} : \xi' < \xi \rangle$ is $<^*$ -increasing, after defining $\zeta(\xi) = \bigcup_{\xi' < \xi} \zeta(\xi')$ we get that $g_{\zeta(\xi)}$ is an exact upper bound of $\langle g_{\zeta(\xi')} : \xi' < \xi \rangle$. But then $g_{\zeta(\xi)}$ is also an exact upper bound of $\langle f_{\zeta(\xi')} : \xi' < \xi \rangle$, and we let $f_\xi = g_{\zeta(\xi)}$.

Let f_λ be defined on κ by $f_\lambda(i) = \mu_i$.

Claim 2.7. Suppose $\delta \leq \lambda$ and $A \subseteq [0, f_\delta)$ has cardinality μ . If $\text{cf } \delta > \kappa$, there is some $\delta' < \delta$ so that $|A \cap [0, f_{\delta'})| = \mu$.

Proof. Find $g < f_\delta$ so that $\sum_{i < \kappa} |A \cap (i \times g(i))| = \mu$. By smoothness there exists some $\delta' < \delta$ so that $g <^* g_{\delta'}$. □

For every $\xi < \lambda$, let $A_\xi = [f_\xi, f_{\xi+1})$ and let $\mathcal{A} = \{A_\xi : \xi < \lambda\}$. Then $\mathcal{A} \subseteq \mathcal{P}([0, f_\lambda))$ is μ -almost disjoint and $|\mathcal{A}| = \lambda$.

For each $\delta \in S$ and $i < \kappa$, let $F_i^\delta = [f_{\gamma_i^\delta}, f_{\gamma_{i+1}^\delta})$. Then $\mathcal{F}_\delta = \{F_i^\delta : i < \kappa\}$ is a μ -almost disjoint family whose union is, by condition (1) on \bar{f} , equal to $[0, f_\delta)$. Fix a μ -ad family $\mathcal{B}_\delta \subseteq \mathcal{P}([0, f_\delta))$ such that $|\mathcal{B}_\delta| = \mathfrak{a}_\mu$, $\mathcal{B}_\delta \cup \mathcal{F}_\delta$ is μ -mad and $\mathcal{B}_\delta \cap \mathcal{F}_\delta = \emptyset$ (by Lemma 1.2).

Claim 2.8. If $\delta \in S$ and $B \in \mathcal{B}_\delta$, then for all $i < \kappa$, it follows that $|B \cap [0, f_{\gamma_i^\delta})| < \mu$.

Proof. If not so, let $i_0 < \kappa$ be the largest value so that $|B \cap [0, f_{\gamma_{i_0}^\delta})| < \mu$; i_0 exists because D_δ is closed. Now $|B \cap F_{i_0}^\delta| = \mu$, a contradiction. □

Let $\mathcal{B} = \bigcup_{\delta \in S} \mathcal{B}_\delta$. Then $|\mathcal{B}| = \mathfrak{a}_\mu \cdot \lambda = \lambda$, and therefore $|\mathcal{A} \cup \mathcal{B}| = \lambda$. We will show now that $\mathcal{A} \cup \mathcal{B}$ is μ -mad.

Suppose that $A = A_\xi \in \mathcal{A}$ and $B \in \mathcal{B}_\delta$ for some $\delta \in S$. If $\xi \geq \delta$, then clearly $|A \cap B| < \mu$, and if $\xi < \delta$, there is some $i < \kappa$ so that $A_\xi \subseteq^* F_i^\delta$ and $|A \cap B| < \mu$ follows from Claim 2.8.

If $B_1 \in \mathcal{B}_{\delta_1}$ and $B_2 \in \mathcal{B}_{\delta_2}$ with $\delta_1 < \delta_2$ in S , then there is some $i < \kappa$ so that $f_{\delta_1} <^* f_{\gamma_i^{\delta_2}}$ and Claim 2.8 gives $|B_1 \cap B_2| < \mu$.

This establishes that $\mathcal{A} \cup \mathcal{B}$ is μ -mad. To verify maximality, let $Z \subseteq [0, f_\lambda)$ be arbitrary of size μ . By Claim 2.7 the first $\xi \leq \lambda$ for which $|Z \cap [0, f_\xi)| = \mu$ is either a successor or of cofinality $\leq \kappa$. Cofinality $< \kappa$ is ruled out by condition (1) on \bar{f} . The case ξ successor implies that $|Z \cap A_\xi| = \mu$. Finally, in the remaining case $\xi = \delta \in S$, there is some $B \in \mathcal{B}_\delta$ so that $|Z \cap B| = \mu$. □

Now the proof of Theorem 2.3 will be completed by the following lemma, whose proof is actually found implicitly in [15]. We shall sketch a proof here too.

Lemma 2.9. *Suppose μ is singular and $\mu < \lambda < \mathfrak{b}_\mu$, λ regular. Then there is a smooth (μ, λ) -scale. If \mathfrak{b}_μ is a successor of a regular cardinal, there is also a smooth (μ, \mathfrak{b}_μ) -scale.*

Proof. Since $\lambda < \mathfrak{b}_\mu$, there exists a product $\prod_{i < \kappa} \mu_i$, where $\kappa = \text{cf } \mu$, so that $\mathfrak{b}(\prod_{i < \kappa} \mu_i, <^*) > \lambda$.

By Claim 1.3 in [15] there exists a λ -scale $\bar{f} = \langle f_\alpha : \alpha < \lambda \rangle$ in some $\prod_{i < \kappa} \mu'_i$ such that for all regular $\theta \in (\kappa, \mu)$, every $\alpha < \lambda$ with $\text{cf } \alpha = \theta$ satisfies that $\bar{f} \upharpoonright \alpha$ is *flat*, that is, is equivalent modulo the bounded ideal on κ to a strictly increasing sequence of ordinal functions on κ .

By Lemma 15 in [10], every $\alpha < \lambda$ with $\text{cf } \alpha > \kappa$ satisfies that $\bar{f} \upharpoonright \alpha$ has an exact upper bound. Now it is clear how to replace \bar{f} by a smooth λ -scale.

Suppose now that $\mathfrak{b}_\mu = \lambda^+$, $\lambda = \text{cf } \lambda$. By [14], 4.1, the set $S_{<\lambda}^{\lambda^+} := \{\alpha : \alpha < \lambda^+ \wedge \text{cf } \alpha < \lambda\}$ is a union of λ sets, each of which carries a square sequence. Therefore, $S_{<\lambda}^{\lambda^+} \in I[\lambda]$. By 2.5 in chapter 1 of [15], there exists a (μ, \mathfrak{b}_μ) -scale in which all points of cofinality $< \mu$ are flat and therefore a smooth (μ, \mathfrak{b}_μ) -scale. \square

In contrast to the case of singular μ , let us mention the following result of A. Blass [4], which generalizes Hechler's result [8]: it is consistent that $\text{MAD}(\aleph_0) = C$, for any prescribed closed set of uncountable cardinals C that satisfies that $[\aleph_1, \aleph_1 + |C|] \subseteq C$ and $\lambda^+ \in C$ whenever $\lambda \in C$ has countable cofinality. For example, by Blass' or by Hechler's results there are universes of set theory in which $\text{MAD}(\aleph_0) = \{\aleph_1, \aleph_{\omega+1}\}$. By Corollary 2.5, in any universe that satisfies this, it follows that $[\aleph_1, \aleph_{\omega+1}] \subseteq \text{MAD}(\aleph_\omega)$.

Recently Brendle [5], using techniques from [16], proved the consistency of $\mathfrak{a} = \aleph_\omega$.

Problem 2.10. Is it consistent that $\mathfrak{a}_{\aleph_\omega} = \aleph_\omega$?

3. CONSISTENCY RESULTS ON $\text{MAD}(\aleph_\omega)$ FROM LARGE CARDINAL AXIOMS

The inequality (1) can be used to control $\text{MAD}(\aleph_\omega)$ by first increasing $\mathfrak{b}_{\aleph_\omega}$ and then increasing \mathfrak{b} . PCF theory implies that whenever the SCH fails at a singular cardinal μ , it follows that $\mathfrak{b}_\mu > \mu^+$. On the other hand, \mathfrak{b}_μ cannot be changed by a ccc forcing.

Before we state the result, let us recall some pcf terminology:

$$\text{pcf}\{\aleph_n : n < \omega\} = \left\{ \mathfrak{b} \left(\prod_n \aleph_n, \leq_I \right) : I \subseteq \mathcal{P}(\omega) \text{ is a proper ideal} \right\}.$$

The relation $<_I$ is defined by $f <_I g \Leftrightarrow \{n : f(n) \geq g(n)\} \in I$.

$\text{pcf}\{\aleph_n : n < \omega\}$ is an interval of regular cardinals and has a maximum. For every $\lambda \in \text{pcf}\{\aleph_n : n < \omega\}$ there exists a *pcf generator* $B_\lambda \subseteq \omega$ so that the following holds: denote by $J_{<\lambda}$ the ideal that is generated by $\{B_\theta : \theta \in \text{pcf}\{\aleph_n : n < \omega\} \wedge \theta < \lambda\}$; then

$$\lambda = \mathfrak{b} \left(\prod_n \aleph_n, \leq_{J_{<\lambda}} \right).$$

Finally, $(\aleph_\omega)^{\aleph_0} = \max \text{pcf}\{\aleph_n : n < \omega\} \times 2^{\aleph_0}$. Therefore, if \aleph_ω is a strong limit, $2^{\aleph_\omega} = \max \text{pcf}\{\aleph_n : n < \omega\}$.

Fact 3.1. For every $\beta < \omega_1$ it is consistent (from large cardinal axioms) that $2^{\aleph_\omega} = \mathfrak{b}_\mu = \aleph_{\omega+\beta+1}$.

Proof. Let V be any universe of set theory in which \aleph_ω is a strong limit cardinal and $2^{\aleph_\omega} = \max \text{pcf}\{\aleph_n : n < \omega\} = \aleph_{\omega+\beta+1}$ [13], [9].

In V , the ideal $J_{<\max \text{pcf}\{\aleph_n : n < \omega\}}$ is proper and is generated by countably many sets. Therefore, by simple diagonalization there exists an infinite $B \subseteq \omega$ so that

$J_{\langle \max \text{pcf}\{\aleph_n : n < \omega\} \upharpoonright B}$ is contained in the ideal of finite subsets of B . Since $\mathfrak{b}(\prod_n \aleph_n, \leq_{J_{\langle \max \text{pcf}\{\aleph_n : n < \omega\}}}) = \aleph_{\omega+\beta+1}$, it follows that $\mathfrak{b}(\prod_{n \in B} \aleph_n, \leq^*) = \aleph_{\omega+\beta+1}$; hence $\mathfrak{b}_{\aleph_\omega} = \aleph_{\omega+\beta+1}$. \square

Theorem 3.2. *For every $\beta < \omega_1$ and $\alpha \leq \omega + \beta + 2$, it is consistent (from large cardinals) that $2^{\aleph_\omega} = \aleph_{\omega+\beta+2}$ and $\text{MAD}(\aleph_\omega) = [\aleph_\alpha, \aleph_{\omega+\beta+2}]$.*

Proof. Start from a model V in which $2^{\aleph_0} = \aleph_1$, \aleph_ω is strong limit and $2^{\aleph_\omega} = \aleph_{\omega+\beta+2}$. Such a model exists by the previous Fact.

For every regular $\aleph_\omega < \lambda \leq \aleph_{\omega+\beta+2}$, there is a smooth λ -scale by Lemma 2.9. Consequently, there is also a smooth $\aleph_{\omega+\beta+2}$ -scale.

Now apply Theorem 2.3 to finish the proof. \square

By Theorem 5.4(b) in [3], after adding many Cohen subsets to ω_1 , $\max \text{MAD}(\aleph_\omega)$ does not increase by much. Therefore, it is consistent to have $\text{MAD}(\aleph_\omega) = [\aleph_1, \aleph_{\omega+\beta+2}]$ as above, and to have 2^{\aleph_ω} arbitrarily large.

ACKNOWLEDGMENTS

The second author would like to thank Uri Abraham for fruitful discussions of some of the proofs in this paper and to thank Isaac Gorelic for useful remarks.

REFERENCES

- [1] B. Balcar and P. Simon. *On collections of almost disjoint families*, Comment. Math. Univ. Carolinae 29 (1988), no 4, pp. 631–646. MR 90b:03072
- [2] B. Balcar and P. Simon. *Disjoint refinements*, in: Handbook of Boolean Algebras, eds. J. D. Monk, R. Bonnet, vol. 2. North-Holland, Amsterdam, 1989.
- [3] J. E. Baumgartner. *Almost-disjoint sets, the dense set problem and the partition calculus*, Ann. Math. Logic 9 (1976), no. 4, 401–439. MR 53:5299
- [4] A. BLASS, *Simple cardinal characteristics of the continuum*, in: Set theory of the reals, Israel Math. Conf. Proc. 6, Bar-Ilan Univ., Ramat Gan, 1993. (1991) 63–90. MR 94i:03098
- [5] J. Brendle. *The almost disjointness number may have countable cofinality*, Trans. Amer. Math. Soc. 355 (2003), no. 7, 2633–2649 (electronic). MR 2004c:03062
- [6] E. K. VAN DOUWEN, *The integers and topology*, in: Handbook of set-theoretic topology, 111–167, North-Holland, Amsterdam, 1984. MR 87f:54008
- [7] P. ERDŐS, S. HECHLER, *On maximal almost-disjoint families over singular cardinals*, Colloquia Mathematica Societatis János Bolyai 10, Infinite and finite sets, Keszthely (Hungary) 1973, pp. 597–604. MR 51:12530
- [8] S. Hechler. *Short complete nested sequences in $\beta\mathbb{N} - \mathbb{N}$ and small maximal almost disjoint families*, Gen. Top. Appl. 2 (1972), 139–149. MR 46:7028
- [9] M. Gitik and M. Magidor. *The singular cardinal hypothesis revisited*, Set theory of the continuum (Berkeley, CA, 1989), 243–279, Math. Sci. Res. Inst. Publ., 26, Springer, New York, 1992. MR 95c:03131
- [10] M. Kojman. *Exact upper bounds and their uses in set theory*, Ann. Pure Appl. Logic 92 (1998), no. 3, 267–282. MR 2000b:03163
- [11] E. MILNER, K. PRIKRY, *Almost disjoint sets*, in: Surveys in combinatorics, (New Cross 1987) pp. 155–172, Cambridge, 1987. MR 88k:04001
- [12] J. D. MONK, *The spectrum of partitions of a Boolean algebra*, Arch. Math. Logic 40 (2001), no. 4, 243–254. MR 2002j:03073
- [13] S. SHELAH, *The singular cardinals problem: independence results.*, Surveys in set theory, 116–134, London Math. Soc. Lecture Note Ser., 87, Cambridge Univ. Press, Cambridge, 1983. MR 87b:03114
- [14] S. Shelah. *Reflecting stationary sets and successors of singular cardinals*, Arch. Math. Logic 31 (1991), no. 1, 25–53. MR 93h:03072

- [15] S. Shelah. *Cardinal Arithmetic*, Oxford University Press, 1994. MR 96e:03001
- [16] S. Shelah. *Are \aleph and \beth your cup of tea?*, Acta Math., to appear.

DEPARTMENT OF MATHEMATICS, BEN-GURION UNIVERSITY OF THE NEGEV, BEER-SHEVA, ISRAEL

E-mail address: `kojman@math.bgu.ac.il`

DEPARTMENT OF MATHEMATICS, BEN-GURION UNIVERSITY OF THE NEGEV, BEER-SHEVA, ISRAEL – AND – INSTITUTE OF MATHEMATICS, UNIVERSITY OF SILESIA, KATOWICE, POLAND

E-mail address: `kubis@math.bgu.ac.il`

INSTITUTE OF MATHEMATICS, HEBREW UNIVERSITY OF JERUSALEM, ISRAEL – AND – DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEW BRUNSWICK, NEW JERSEY

E-mail address: `shelah@math.huji.ac.il`