

FIBONACCI NUMBERS THAT ARE NOT SUMS OF TWO PRIME POWERS

FLORIAN LUCA AND PANTELIMON STĂNICĂ

(Communicated by David E. Rohrlich)

ABSTRACT. In this paper, we construct an infinite arithmetic progression \mathcal{A} of positive integers n such that if $n \in \mathcal{A}$, then the n th Fibonacci number is not a sum of two prime powers.

1. INTRODUCTION

In 1849, A. de Polignac [7] asked if every odd positive integer can be represented as the sum of a power of 2 and a prime (or 1). Euler did note that 959 was not of this form. Romanoff [8] used the Brun sieve to show that a positive proportion of integers are representable in this way. Later, van der Corput [2] showed that a positive proportion of integers are not representable in this way by using covering congruences. With the same method as van der Corput's, Erdős [3] constructed a residue class of odd numbers which contains no integers of the above form. Extending Erdős's argument, Cohen and Selfridge [1] constructed a 26 digit number which is neither the sum nor the difference of two prime powers. Inspired by their work, Z.W. Sun (see [9]) constructed a residue class of odd integers consisting exclusively of numbers not of the form $\pm p^a \pm q^b$ with some primes p and q and some nonnegative integers a and b . We mention that Erdős asked if there exist infinitely many positive integers which are not representable as a sum or difference of two powers (see [5]) and a partial result can be found in [6].

In this paper, we show that there exist infinitely many positive integers which are not of the form $p^a + q^b$ with primes p and q and nonnegative integers a and b and which further can be chosen to be members of the Fibonacci sequence $(F_n)_{n \geq 0}$ given by $F_0 = 0$, $F_1 = 1$ and $F_{n+2} = F_{n+1} + F_n$ for $n \geq 0$.

In what follows, we use the Vinogradov symbols \gg and \ll with their usual meanings. We recall that given two functions A and B of the real variable x , the notations $A \ll B$ and $B \gg A$ are equivalent to the fact that the inequality $|A(x)| \leq cB(x)$ holds with some positive constant c and for all sufficiently large real numbers x .

Received by the editors February 13, 2004.

2000 *Mathematics Subject Classification*. Primary 11B39; Secondary 11B25, 11B50, 11P32.

Key words and phrases. Fibonacci numbers, arithmetic progressions, covering system of congruences.

©2005 American Mathematical Society
Reverts to public domain 28 years from publication

2. THE RESULT

Our main result is the following.

Theorem 1. *There exists a positive integer n_0 such that if $n > n_0$ and*

$$n \equiv 1807873 \pmod{3543120},$$

then $F_n \neq p^a + q^b$ with p, q prime numbers and a, b nonnegative integers.

We use the same method employed in [2, 3, 9]. However, there are additional difficulties which arise because we want our numbers to belong to the Fibonacci sequence.

Let us quickly recall how one can create a residue class of odd integers most of which are not sums of two prime powers. Well, assume that n is odd and a sum of two prime powers, say $p^a + q^b$. Since n is odd, it follows that one of p and q , say p , equals 2. Suppose now that we are given a finite set of triples $(a_i, b_i, p_i)_{i=1}^s$ where a_i and b_i are nonnegative integers and p_i are distinct primes such that the following hold:

- (i): for every $a \in \mathbb{Z}$ there exists $i \in \{1, \dots, s\}$ such that $a \equiv a_i \pmod{b_i}$;
- (ii): $p_i \mid (2^{b_i} - 1)$ for all $i = 1, \dots, s$;

We may then choose n to belong to the arithmetic progression given by $n \equiv 2^{a_i} \pmod{p_i}$. Since the p_i 's are distinct primes, the above system admits a unique solution modulo $p_1 \dots p_s$ by the Chinese Remainder Lemma. Let \mathcal{A} be this progression. Assume that n is sufficiently large in the above progression \mathcal{A} and that $n = 2^a + q^b$ holds with some prime number q . By (i) above, there exists $i \in \{1, \dots, s\}$ such that $a \equiv a_i \pmod{b_i}$. By (ii) above, $2^a \equiv 2^{a_i} \pmod{p_i}$. However, $n \equiv 2^{a_i} \pmod{p_i}$, and therefore $n \equiv 2^a \pmod{p_i}$. Hence, $p_i \mid (n - 2^a)$. However, $n - 2^a = q^b$. Since q is prime, it follows that $p_i = q$. Now let X be a very large positive real number. The number of positive integers $n \leq X$ which belong to \mathcal{A} is $\gg X$. However, the number of positive integers of the form $2^a + q^b$ with $q \in \{p_1, \dots, p_s\}$ and which are $\leq X$ is $\ll \log^2 X$. This shows that most numbers in \mathcal{A} are not of the form $p^a + q^b$.

We now modify the above construction in order to insure that our numbers can be chosen from the Fibonacci sequence. Let k be a positive integer. It is known that $(F_n)_{n \geq 0}$ is periodic modulo k . We let $h(k)$ denote this period. Moreover, for integers f and k we write $\mathcal{A}(f, k)$ for the set of residue classes n modulo $h(k)$ such that $F_n \equiv f \pmod{k}$.

Assume now that $(a_i, b_i, p_i)_{i=1}^s$ is a finite set of triples of nonnegative integers a_i and b_i and distinct odd primes p_i for $i = 1, \dots, s$ which fulfill the following conditions:

- (i): for every $a \in \mathbb{Z}$ there exists $i \in \{1, \dots, s\}$ such that $a \equiv a_i \pmod{b_i}$;
- (ii): $p_i \mid (2^{b_i} - 1)$ for all $i = 1, \dots, s$;
- (iii): the set

$$\bigcap_{i=1}^s \mathcal{A}(2^{a_i}, p_i) \neq \emptyset.$$

Moreover, if there exists $i \in \{1, \dots, s\}$ such that $3 \mid h(p_i)$, then we shall assume that the above intersection contains a class coprime to 3.

Let x be an element of the set $\bigcap_{i=1}^s \mathcal{A}(2^{a_i}, p_i)$. Note that x is defined only modulo $M = \text{lcm}[h(p_1), \dots, h(p_s)]$. Moreover, if $3 \mid M$, then x is not a multiple of 3. If M is not a multiple of 3 we replace M by $3M$ and x by the solution of

the system of congruences $x \pmod{M}$ and $1 \pmod{3}$. Assume now that $n \equiv x \pmod{M}$ and that $F_n = p^a + q^b$ with some primes p and q and nonnegative integers a and b . It then follows that F_n is an odd integer, because the only even Fibonacci numbers are those whose indices are multiples of 3. Since F_n is odd, it follows that one of p and q , say p , is 2. By (i) above, there exists $i \in \{1, \dots, s\}$ such that $a \equiv a_i \pmod{b_i}$. By (ii) above, it follows that $2^a \equiv 2^{a_i} \pmod{p_i}$. By the choice of n , we have that $F_n \equiv 2^{a_i} \pmod{p_i}$. Thus, $F_n \equiv 2^a \pmod{p_i}$. In particular, $p_i \mid (F_n - 2^a)$. However, since $F_n - 2^a = q^b$, it follows that $q = p_i$. Thus, $q \in \{p_1, \dots, p_s\}$. We now get that $F_n = 2^a + p_i^b$ for some $i = 1, \dots, s$. Since $F_n = \frac{1}{\sqrt{5}}(\alpha^n - \beta^n)$, where $\alpha = \frac{1+\sqrt{5}}{2}$ and $\beta = \frac{1-\sqrt{5}}{2}$, we may apply a well-known result from the theory of \mathcal{S} -unit equations (see [4]) to conclude that such an equation can have only finitely many solutions (n, a, b) . Thus, there exists n_0 with the property that if $n > n_0$ and $n \equiv x \pmod{M}$, then F_n is not of the form $p^a + q^b$ with primes p and q and nonnegative integers a and b .

In order to finish the proof of the theorem, it suffices to find a finite set of triples $(a_i, b_i, p_i)_{i=1}^s$ fulfilling (i)–(iii) above.

We first note that by taking $s = 7$ and

$$((a_1, b_1), \dots, (a_7, b_7)) = ((0, 2), (0, 3), (3, 4), (1, 12), (5, 36), (17, 36), (29, 36))$$

we get that (i) above is fulfilled. Indeed, every integer is congruent either to 0 (mod 2) or to 1 (mod 2) or to 0 (mod 3) or to 3 (mod 4) or to 1 (mod 12) or to 5 (mod 12), and in this last case it is congruent to either 5, 17 or 29 modulo 36. We now take $(p_1, \dots, p_7) = (3, 7, 5, 13, 19, 37, 73)$ and note that condition (ii) above is fulfilled. It is easy to check that $(h(p_1), \dots, h(p_7)) = (8, 16, 20, 28, 18, 76, 148)$. Finally, it is easy to see that

$$\begin{aligned} \mathcal{A}(2^{a_1}, p_1) &= \mathcal{A}(2^0, 3) = \{1, 2, 7\} \pmod{8}, \\ \mathcal{A}(2^{a_2}, p_2) &= \mathcal{A}(2^0, 7) = \{1, 2, 6, 15\} \pmod{16}, \\ \mathcal{A}(2^{a_3}, p_3) &= \mathcal{A}(2^3, 5) = \{4, 6, 7, 13\} \pmod{20}, \\ \mathcal{A}(2^{a_4}, p_4) &= \mathcal{A}(2^1, 13) = \{3, 25\} \pmod{28}, \\ \mathcal{A}(2^{a_5}, p_5) &= \mathcal{A}(2^5, 19) = \{7, 11\} \pmod{18}, \\ \mathcal{A}(2^{a_6}, p_6) &= \mathcal{A}(2^{17}, 37) = \{10, 15, 28, 61\} \pmod{76}, \\ \mathcal{A}(2^{a_7}, p_7) &= \mathcal{A}(2^{29}, 73) = \{53, 95\} \pmod{148}. \end{aligned}$$

One now checks that the system of congruences $x \equiv 1 \pmod{8}$, $x \equiv 1 \pmod{16}$, $x \equiv 13 \pmod{20}$, $x \equiv 25 \pmod{28}$, $x \equiv 7 \pmod{18}$, $x \equiv 61 \pmod{76}$ and $x \equiv 53 \pmod{148}$ has a positive integer solution x . In fact, the above system is equivalent to the system of congruences $x \equiv 1 \pmod{16}$, $x \equiv 3 \pmod{5}$, $x \equiv 4 \pmod{7}$, $x \equiv 7 \pmod{9}$, $x \equiv 4 \pmod{19}$ and $x \equiv 16 \pmod{37}$. Solving this system, we get $x \equiv 1807873 \pmod{3543120}$, which, together with the above arguments completes the proof of Theorem 1.

We would like to conclude by offering the following problem.

Problem. Find an arithmetic progression of positive integers $A \pmod{B}$ such that if $n > n_0$ satisfies $n \equiv A \pmod{B}$, then $F_n \neq \pm p^a \pm q^b$ for any primes p, q and nonnegative integers a, b .

ACKNOWLEDGMENTS

This work was done during a visit of F. L. at the Mathematics Department of the Auburn University at Montgomery. He thanks the people of this department for their warm hospitality.

REFERENCES

- [1] F. Cohen and J. L. Selfridge, ‘Not every number is the sum or difference of two prime powers’, *Math. Comp.* **29** (1975), 79–81. MR0376583 (51:12758)
- [2] J. G. van der Corput, ‘On de Polignac’s conjecture’, *Simon Stevin* **27** (1950), 99–105. MR0035298 (11:714e)
- [3] P. Erdős, ‘On integers of the form $2^k + p$ and some related problems’, *Summa Brasil. Math.* **2** (1950), 113–123. MR0044558 (13:437i)
- [4] J.-H. Evertse, K. Győry, C. L. Stewart and R. Tijdeman, ‘ S -unit equations and their applications’, in *New advances in transcendence theory* (Durham, 1986), 110–174, Cambridge Univ. Press, Cambridge, 1988. MR0971998 (89j:11028)
- [5] R. K. Guy, *Unsolved Problems in Number Theory*, Springer, 1994. MR1299330 (96e:11002)
- [6] F. Luca, ‘On integers which are not differences of two powers’, *Mathematika*, **49** (2002), no. 1–2. 231–243 (2004). MR2059057 (2004c:11041)
- [7] A. de Polignac, ‘Recherches nouvelles sur les nombres premiers’, *C.R. Acad. Sci. Paris Math.* **29** (1949), 397–401, 738–739.
- [8] N. P. Romanoff, ‘Über einige Sätze der additiven Zahlentheorie’, *Math. Ann.* **57** (1934), 668–678.
- [9] Z. W. Sun, ‘On integers not of the form $\pm p^a \pm q^b$ ’, *Proc. Amer. Math. Soc.* **128** (2000), 997–1002. MR1695111 (2000i:11157)

IMATE, UNAM, AP. POSTAL 61-3 (XANGARI), CP. 58 089, MORELIA, MICHOACÁN, MEXICO
E-mail address: fluca@matmor.unam.mx

DEPARTMENT OF MATHEMATICS, AUBURN UNIVERSITY MONTGOMERY, MONTGOMERY, ALABAMA
 36124-4023
E-mail address: pstanica@mail.aum.edu