

UNIFORMLY BOUNDED LIMIT OF FRACTIONAL HOMOMORPHISMS

PEDRO J. MIANA

(Communicated by Jonathan M. Borwein)

ABSTRACT. We show that a bounded homomorphism $T : L_\omega^1(\mathbb{R}^+) \rightarrow \mathcal{A}$ is equivalent to a uniformly bounded family of fractional homomorphisms $T_\alpha : AC_\omega^{(\alpha)}(\mathbb{R}^+) \rightarrow \mathcal{A}$ for any $\alpha > 0$. We add this characterization to the Widder-Arendt-Kisyański theorem and relate it to α -times integrated semigroups.

1. INTRODUCTION

Let \mathcal{A} be a Banach algebra (with or without identity). For $\Omega \subset \mathbb{C}$, a family $(r_\lambda)_{\lambda \in \Omega}$ of elements of \mathcal{A} is called a *pseudo-resolvent* if the equation $r_\lambda - r_\mu = (\mu - \lambda)r_\lambda r_\mu$ holds for $\lambda, \mu \in \Omega$. An example of a pseudo-resolvent is the following. Let \mathbb{R}, \mathbb{R}^+ and \mathbb{C} be the sets of real, positive real and complex numbers, respectively. For each $\lambda \in \mathbb{R}$, denote by ϵ_λ the function

$$\epsilon_\lambda(t) = e^{\lambda t}, \quad t \in \mathbb{R}^+.$$

Take $\omega \in \mathbb{R}^+ \cup \{0\}$ and let $L_\omega^1(\mathbb{R}^+)$ be the usual Banach algebra with norm given by

$$\|f\|_\omega := \int_0^\infty |f(t)|e^{\omega t} dt < +\infty,$$

and the convolution $f * g(t) := \int_0^t f(t-s)g(s)ds$, with $t \geq 0$, as its product. Then $(\epsilon_{-\lambda})_{\lambda \in (\omega, +\infty)}$ is a pseudo-resolvent in $L_\omega^1(\mathbb{R}^+)$ and verifies

$$\| \underbrace{\epsilon_{-\lambda} * \cdots * \epsilon_{-\lambda}}_{n \text{ times}} \|_\omega = \frac{1}{(\lambda - \omega)^n}, \quad \lambda \in (\omega, +\infty), \quad n \in \mathbb{N}.$$

Moreover, the set $(\epsilon_{-\lambda})_{\lambda \in (\omega, +\infty)}$ is linearly dense in $L_\omega^1(\mathbb{R}^+)$.

The next result shows the equivalence between a homomorphism $T : L_\omega^1(\mathbb{R}^+) \rightarrow \mathcal{A}$ and a class of pseudo-resolvents; in other words, the family $(\epsilon_{-\lambda})_{\lambda \in (\omega, +\infty)}$ is universal for this class of pseudo-resolvents. We present here an early version; see [8] and [3] which is included in [9, Theorem 5.1]. Recently, this result has been called the Widder-Arendt-Kisyański theorem; see for example [5, Theorem 1.1].

Received by the editors February 1, 2003.

2000 *Mathematics Subject Classification*. Primary 47D62; Secondary 26A33, 46J25.

Key words and phrases. Pseudo-resolvents, homomorphisms, integrated semigroups.

This work was supported by a grant from Programa Europa, CAI, 2002. This paper was made during a visit to the Charles University in Prague. The author thanks Dr. Eva Fasangova and the Analysis Mathematical Department for the stay in Prague.

©2005 American Mathematical Society
Reverts to public domain 28 years from publication

Theorem 1.1 ([8]). *Let \mathcal{A} be a Banach algebra, let $\omega \geq 0$, let $(r_\lambda)_{\lambda \in (\omega, +\infty)}$ be a pseudo-resolvent in \mathcal{A} and let*

$$M = \sup\{(\lambda - \omega)^n \|r_\lambda^n\| : n \in \mathbb{N}, \lambda \in (\omega, +\infty)\}.$$

Then the following conditions are equivalent:

- (i) $M < +\infty$.
- (ii) *There exists a continuous homomorphism $T : L_\omega^1(\mathbb{R}^+) \rightarrow \mathcal{A}$ such that $T(\epsilon_{-\lambda}) = r_\lambda$ for each $\lambda \in (\omega, +\infty)$.*

Furthermore, if a continuous homomorphism $T : L_\omega^1(\mathbb{R}^+) \rightarrow \mathcal{A}$ satisfying $T(\epsilon_{-\lambda}) = r_\lambda$ for each $\lambda \in (\omega, +\infty)$ exists, then it is unique and $\|T\| = M$.

This theorem has many interesting applications, as W. Chojnacki pointed out, and it is equivalent to Hille-Yosida theorem; see [3] and [4]. Another different proof of this result is given in [2], where the author uses Yosida approximation in a more general setting. A generalization of [9, Theorem 5.1] is presented in [5] improving Kiszyński's arguments.

In this paper, we arrive at Theorem 1.1 with an original point of view. We show that a bounded homomorphism $T : L_\omega^1(\mathbb{R}^+) \rightarrow \mathcal{A}$ is equivalent to a uniformly bounded family of homomorphisms $T_\alpha : AC_\omega^{(\alpha)}(\mathbb{R}^+) \rightarrow \mathcal{A}$ for any $\alpha > 0$; see Theorem 3.1. Fractional Banach algebras $AC_\omega^{(\alpha)}(\mathbb{R}^+)$ are included in $L_\omega^1(\mathbb{R}^+)$ and defined in the second section. The proof of this main result is based on Theorem 1.1.

A family of fractional algebras similar to $AC_\omega^{(\alpha)}(\mathbb{R}^+)$ was introduced in [10], working with α -times integrated semigroups. Integrated semigroups are considered in the fourth section to complete a different approach to the Widder-Arendt-Kiszyński theorem in which Lipschitz and Hölder continuous functions appear; see [1, Theorem 1.1], [6, Theorem 2.6], [9, Corollary 7.2], or [7, Theorem 3.2]. Lipschitz functions are considered in the integrated version of the Widder's characterization of Laplace transforms on arbitrary Banach spaces.

Actually, α -times integrated semigroups (in the case $\alpha = 1$) were also used to prove Theorem 1.1 with $\omega = 0$ in [8, Theorem 4.2]. However, we present here a different technique since we consider α -times integrated semigroups for any $\alpha > 0$ and the limit when α goes to zero.

2. FRACTIONAL BANACH ALGEBRAS

Suppose $f \in \mathcal{D}_+$, where $C_c^\infty[0, \infty) \equiv \mathcal{D}_+$ is the set of infinitely differentiable functions with compact support on $[0, \infty)$. The *Weyl fractional integral* of f of order $\alpha > 0$, $W^{-\alpha}f$, is defined by

$$W^{-\alpha}f(u) = \frac{1}{\Gamma(\alpha)} \int_u^\infty (t-u)^{\alpha-1} f(t) dt, \quad u \geq 0,$$

with $\alpha > 0$; see for example [11]. This operator $W^{-\alpha} : \mathcal{D}_+ \rightarrow \mathcal{D}_+$ is one-to-one and onto. Its inverse, W^α , is called a *Weyl fractional derivate* (of f) of order α and $W^\alpha = (-1)^n \frac{d^n}{dt^n} W^{-(n-\alpha)}$ with $n \in \mathbb{N}$ and $n > \alpha$.

Theorem 2.1. *Suppose $\alpha > 0$ and $\omega \geq 0$. If $f \in \mathcal{D}_+$, then the formula*

$$\|f\|_{(\alpha, \omega)} := \frac{1}{\Gamma(\alpha+1)} \int_0^\infty t^\alpha e^{\omega t} |W^\alpha f(t)| dt$$

defines a norm in \mathcal{D}_+ such that

$$\|f * g\|_{(\alpha, \omega)} \leq (2^{\alpha+1} - 1) \|f\|_{(\alpha, \omega)} \|g\|_{(\alpha, \omega)},$$

with $f, g \in \mathcal{D}_+$. We denote by $AC_\omega^{(\alpha)}(\mathbb{R}^+)$ the Banach algebra obtained by the completion of \mathcal{D}_+ in this norm.

Proof. The proof is similar to [10, Theorem 1.2]. □

Remark 2.2. For $\alpha = 0$, it is clear that $AC_\omega^{(0)}(\mathbb{R}^+) = L_\omega^1(\mathbb{R}^+)$ and $AC_\omega^{(\beta)}(\mathbb{R}^+) \hookrightarrow AC_\omega^{(\alpha)}(\mathbb{R}^+)$ holds with $0 \leq \alpha \leq \beta$.

It is easy to check that $W^\alpha(\epsilon_{-\lambda}) = \lambda^\alpha \epsilon_{-\lambda}$ with $\alpha \in \mathbb{R}$ and $\lambda > 0$. Functions $(\epsilon_{-\lambda})_{\lambda \in (\omega, +\infty)}$ belong to $AC_\omega^{(\alpha)}(\mathbb{R}^+)$ with $\alpha \geq 0$ and

$$(2.1) \quad \|\epsilon_{-\lambda}\|_{(\alpha, \omega)} = \frac{\lambda^\alpha}{(\lambda - \omega)^{\alpha+1}}, \quad \lambda \in (\omega, +\infty).$$

Note that $\|\epsilon_{-\lambda}\|_{(\alpha, \omega)} \rightarrow \|\epsilon_{-\lambda}\|_\omega$ when $\alpha \rightarrow 0^+$ and $\lambda \in (\omega, +\infty)$. Moreover, $(\epsilon_{-\lambda})_{\lambda \in (\omega, \infty)}$ is a pseudo-resolvent in $AC_\omega^{(\alpha)}(\mathbb{R}^+)$ and $(n\epsilon_{-n})_{n > \omega}$ is a bounded approximate identity in $AC_\omega^{(\alpha)}(\mathbb{R}^+)$ for any $\alpha \geq 0$.

Proposition 2.3. *The set $(\epsilon_{-\lambda})_{\lambda \in (\omega, +\infty)}$ is linearly dense in $AC_\omega^{(\alpha)}(\mathbb{R}^+)$ with $\alpha, \omega \geq 0$.*

Proof. It is a consequence of $W^\alpha(\epsilon_{-\lambda}) = \lambda^\alpha \epsilon_{-\lambda}$ with $\alpha \geq 0$, $\lambda > \omega$ and [5, Proposition 2.2]. □

3. MAIN RESULT

Theorem 3.1. *Let \mathcal{A} be a Banach algebra.*

(i) *If there exists a continuous homomorphism $T : L_\omega^1(\mathbb{R}^+) \rightarrow \mathcal{A}$, then there exists a family of continuous homomorphisms $T_\alpha : AC_\omega^{(\alpha)}(\mathbb{R}^+) \rightarrow \mathcal{A}$ for each $\alpha > 0$ such that $T_\alpha(\epsilon_{-\lambda}) = T(\epsilon_{-\lambda})$ for each $\lambda \in (\omega, +\infty)$ and $\|T_\alpha\| \leq \|T\|$ for any $\alpha > 0$.*

(ii) *Conversely, if for each $\alpha > 0$ there exist a continuous homomorphisms $T_\alpha : AC_\omega^{(\alpha)}(\mathbb{R}^+) \rightarrow \mathcal{A}$ such that $T_\alpha(\epsilon_{-\lambda})$ does not depend on α for each $\lambda \in (\omega, +\infty)$ and $\limsup_{\alpha \rightarrow 0^+} \|T_\alpha\| < +\infty$, then there exists a unique continuous homomorphism $T : L_\omega^1(\mathbb{R}^+) \rightarrow \mathcal{A}$ such that $T(\epsilon_{-\lambda}) = T_\alpha(\epsilon_{-\lambda})$ for each $\lambda \in (\omega, +\infty)$ and $\|T\| \leq \limsup_{\alpha \rightarrow 0^+} \|T_\alpha\|$.*

Proof. The first part follows from Remark 2.2. To prove (ii), we use Theorem 1.1. We define $r_\lambda := T_\alpha(\epsilon_\lambda)$ for each $\lambda > \omega$. The family $(r_\lambda)_{\lambda \in (\omega, \infty)}$ is well defined and is a pseudo-resolvent in \mathcal{A} . For $n \in \mathbb{N}$ and $\lambda \in (\omega, +\infty)$,

$$\|r_\lambda^n\| \leq \|T_\alpha\| \underbrace{\|\epsilon_\lambda * \dots * \epsilon_\lambda\|}_{n \text{ times}}_{(\alpha, \omega)}.$$

We apply Theorem 2.1 and get that

$$\|\underbrace{\epsilon_\lambda * \dots * \epsilon_\lambda}_{n \text{ times}}\|_{(\alpha, \omega)} \leq (2^{\alpha+1} - 1)^{n-1} \|\epsilon_\lambda\|_{(\alpha, \omega)}^n = (2^{\alpha+1} - 1)^{n-1} \frac{\lambda^{n\alpha}}{(\lambda - \omega)^{n(\alpha+1)}},$$

where we use (1). Now we obtain

$$\|r_\lambda^n\| \leq \lim_{\alpha \rightarrow 0} \left((2^{\alpha+1} - 1)^{n-1} \frac{\lambda^{n\alpha}}{(\lambda - \omega)^{n(\alpha+1)}} \right) \limsup_{\alpha \rightarrow 0^+} \|T_\alpha\|,$$

and we get

$$\sup\{(\lambda - \omega)^n \|r_\lambda^n\| : n \in \mathbb{N}, \lambda \in (\omega, +\infty)\} \leq \limsup_{\alpha \rightarrow 0^+} \|T_\alpha\| < +\infty.$$

By Theorem 1.1, there exists a unique continuous homomorphism $T : L^1_\omega(\mathbb{R}^+) \rightarrow \mathcal{A}$ such that $T(\epsilon_{-\lambda}) = r_\lambda = T_\alpha(\epsilon_{-\lambda})$ for each $\lambda \in (\omega, +\infty)$ and $\|T\| \leq \limsup_{\alpha \rightarrow 0^+} \|T_\alpha\|$. \square

Remark 3.2. Since the set $(\epsilon_{-\lambda})_{\lambda \in (\omega, +\infty)}$ is linearly dense in $AC^{(\alpha)}_\omega(\mathbb{R}^+)$ (Proposition 2.3), $T_\alpha(f) = T(f)$ for any $f \in AC^{(\alpha)}_\omega(\mathbb{R}^+)$ and $\alpha > 0$. Also, we may combine (i) and (ii) to get

$$\|T\| = \limsup_{\alpha \rightarrow 0^+} \|T_\alpha\| = \sup_{\alpha > 0} \|T_\alpha\|,$$

where we use $\|T_\beta\| \leq \|T_\alpha\|$ with $\beta \geq \alpha > 0$.

4. CONNECTION WITH INTEGRATED SEMIGROUPS

Widder’s characterization of Laplace transforms of real-valued bounded functions which states that for $r \in C^{(\infty)}(0, \infty)$ there exists $f \in L^\infty(0, \infty)$ such that

$$r(\lambda) = \int_0^\infty e^{-\lambda t} f(t) dt, \quad \lambda > 0,$$

if and only if

$$\sup\{\lambda^{n+1} \frac{|r^{(n)}(\lambda)|}{n!} : \lambda > 0, n \in \mathbb{N}\} < +\infty,$$

(see [12]) was considered by W. Arendt on arbitrary Banach space X . An integrated version of this result is given in terms of a function $F : [0, \infty) \rightarrow X$ such that $F(0) = 0$, $\|F(t+h) - F(t)\| \leq Mh$, ($t, h \geq 0$) and

$$r(\lambda) = \lambda \int_0^\infty e^{-\lambda t} F(t) dt, \quad \lambda > 0;$$

see [1, Theorem 1.1]. This result was improved by M. Hieber who showed that for any $\alpha \in (0, 1]$ there exists a function $F_\alpha : [0, \infty) \rightarrow X$ such that $F_\alpha(0) = 0$, $\|F_\alpha(t+h) - F_\alpha(t)\| \leq Mh^\alpha$, ($t, h \geq 0$) and

$$r(\lambda) = \lambda^\alpha \int_0^\infty e^{-\lambda t} F(t) dt, \quad \lambda > 0;$$

see [7, Theorem 3.2]. If r is a pseudo-resolvent in a Banach algebra \mathcal{A} , then the function F_α is an α -times integrated semigroup in \mathcal{A} . We now give here the definition of α -times integrated semigroups in a Banach algebra \mathcal{A} , since they are usually considered in the setting of linear and bounded operators on a Banach space X ; see for example [7].

Definition 4.1. For any $\alpha > 0$, an α -times integrated semigroup $s_\alpha(\cdot) : [0, \infty) \rightarrow \mathcal{A}$ is a continuous mapping with $s_\alpha(0) = 0$ and

$$(4.1) \quad s_\alpha(t)s_\alpha(s) = \frac{1}{\Gamma(\alpha)} \left(\int_t^{t+s} (t+s-r)^{\alpha-1} s_\alpha(r) dr - \int_0^s (t+s-r)^{\alpha-1} s_\alpha(r) dr \right),$$

with $t, s \geq 0$.

If $\|s_\alpha(t)\| \leq Ce^{\omega t}$ for $C, t \geq 0$ and $\omega \in \mathbb{R}$, then condition (2) is equivalent (by Laplace transform) to $r_\lambda := \lambda^\alpha \int_0^\infty e^{-\lambda t} s_\alpha(t) dt$, with $\lambda > \omega$ a pseudo-resolvent in \mathcal{A} ; see the analogous proof in Banach space in [7]. If $(s_\alpha(t))_{t \geq 0}$ is an α -times integrated semigroup in \mathcal{A} , then $(s_\nu(t))_{t \geq 0}$ with

$$(4.2) \quad s_\nu(t) := \frac{1}{\Gamma(\nu - \alpha)} \int_0^t (t - s)^{\nu - \alpha - 1} s_\alpha(s) ds, \quad t \geq 0,$$

is a ν -times integrated semigroup with $\nu > \alpha$ in \mathcal{A} . The set of Bochner-Riesz functions $(R_t^\nu)_{t \geq 0}$, i.e.,

$$R_t^\nu(s) := \frac{(t - s)^{\nu - 1}}{\Gamma(\nu)} \chi_{(0,t)}(s), \quad t \geq 0,$$

is an example of a ν -times integrated semigroup in $AC_\omega^{(\alpha)}(\mathbb{R}^+)$ for $\nu > \alpha, \omega \geq 0$ and

$$\|R_t^\nu\|_{(\alpha, \omega)} \leq \frac{t^\nu e^{t\omega}}{\Gamma(\nu + 1)}, \quad t \geq 0.$$

Now, we may add in the Widder-Arendt-Kiszyński theorem more conditions related to fractional homomorphisms and α -times integrated semigroups.

Theorem 4.2. *Let \mathcal{A} be a Banach algebra, let ω be a non-negative number, let $(r_\lambda)_{\lambda \in (\omega, +\infty)}$ be a pseudo-resolvent in \mathcal{A} and let*

$$M = \sup\{(\lambda - \omega)^n \|r_\lambda^n\| : n \in \mathbb{N}, \lambda \in (\omega, +\infty)\}.$$

Then the following conditions are equivalent:

- (i) $M < +\infty$.
- (ii) *There exists a continuous homomorphism $T : L_\omega^1(\mathbb{R}^+) \rightarrow \mathcal{A}$ such that $T(\epsilon_{-\lambda}) = r_\lambda$ for each $\lambda \in (\omega, +\infty)$.*
- (iii) *For any $\alpha > 0$, there exists an α -times integrated semigroup $(s_\alpha(t))_{t \geq 0} \subset \mathcal{A}$ such that $\|s_\alpha(t)\| \leq \frac{C}{\Gamma(\alpha + 1)} t^\alpha e^{\omega t}$ with $C, t \geq 0$ and $r_\lambda = \lambda^\alpha \int_0^\infty e^{-\lambda t} s_\alpha(t) dt$, with $\lambda > \omega$.*
- (iv) *For any $\alpha > 0$, there exists a continuous homomorphism $T_\alpha : AC_\omega^{(\alpha)}(\mathbb{R}^+) \rightarrow \mathcal{A}$ such that $T_\alpha(\epsilon_{-\lambda}) = r_\lambda$ for each $\lambda \in (\omega, +\infty)$ and $\sup_{\alpha > 0} \|T_\alpha\| < +\infty$.*

Furthermore, if a continuous homomorphism $T : L_\omega^1(\mathbb{R}^+) \rightarrow \mathcal{A}$ such that $T(\epsilon_{-\lambda}) = r_\lambda$ for each $\lambda \in (\omega, +\infty)$ exists, then it is unique, $T(f) = T_\alpha(f)$ for $f \in AC_\omega^{(\alpha)}(\mathbb{R}^+)$ and

$$M = \|T\| = \sup_{\alpha > 0} \|T_\alpha\| = \inf\{C : \|s_\alpha(t)\| \leq C \frac{t^\alpha e^{\omega t}}{\Gamma(\alpha + 1)}, t \geq 0\}.$$

Proof. (ii) \Rightarrow (iii) Take $(R_t^\alpha)_{t > 0}$ to be the family of Bochner-Riesz functions of order $\alpha > 0$. Then $(R_t^\alpha)_{t \geq 0} \subset L_\omega^1(\mathbb{R}^+)$ and we define $s_\alpha(t) := T(R_t^\alpha)$, for $t > 0$ and $s_\alpha(0) := 0$. Since $(R_t^\alpha)_{t > 0}$ is an α -times integrated semigroup in $L_\omega^1(\mathbb{R}^+)$, $(s_\alpha(t))_{t \geq 0}$ is an α -times integrated semigroup in \mathcal{A} and

$$\|s_\alpha(t)\| \leq \|T\| \|R_t^\alpha\|_\omega \leq \|T\| \frac{t^\alpha}{\Gamma(\alpha + 1)} e^{\omega t}.$$

By continuity of T , we have for $\lambda > \omega$,

$$\lambda^\alpha \int_0^\infty e^{-\lambda t} s_\alpha(t) dt = \lambda^\alpha T \left(\int_0^\infty e^{-\lambda t} R_t^\alpha dt \right) = \lambda^\alpha T(W^{-\alpha} \epsilon_\lambda) = T(\epsilon_\lambda) = r_\lambda.$$

(iii) \Rightarrow (iv) We define $T_\alpha : AC_\omega^{(\alpha)}(\mathbb{R}^+) \rightarrow \mathcal{A}$ by

$$T_\alpha(f) := \int_0^\infty W^\alpha f(t) s_\alpha(t) dt, \quad f \in \mathcal{D}_+.$$

Following the same arguments as in [10, Theorem 3.1], we prove that T_α are continuous homomorphisms, $\|T_\alpha\| \leq C$ for any $\alpha > 0$ and $\sup_{\alpha > 0} \|T_\alpha\| \leq C < +\infty$. Since $\epsilon_\lambda \in AC_\omega^{(\alpha)}(\mathbb{R}^+)$ for $\lambda > \omega$ and $W^\alpha(\epsilon_\lambda) = \lambda^\alpha \epsilon_\lambda$, we have $T_\alpha(\epsilon_\lambda) = r_\lambda$ for $\lambda > \omega$.

(iv) \Rightarrow (ii) It is proved by Theorem 3.1(ii).

If a continuous homomorphism $T : L_\omega^1(\mathbb{R}^+) \rightarrow \mathcal{A}$ satisfying $T(\epsilon_{-\lambda}) = r_\lambda$ for each $\lambda \in (\omega, +\infty)$ exists, then $AC_\omega^{(\alpha)}(\mathbb{R}^+) \hookrightarrow L_\omega^1(\mathbb{R}^+)$ and $T(f) = T_\alpha(f)$ for $f \in AC_\omega^{(\alpha)}(\mathbb{R}^+)$. Moreover, if we collect in the proof the following inequalities,

$$\|T\| = M = \sup_{\alpha > 0} \|T_\alpha\| \leq \inf\{C : \|s_\alpha(t)\| \leq C \frac{t^\alpha e^{\omega t}}{\Gamma(\alpha + 1)}, t \geq 0\} \leq \|T\|,$$

we finally get the equality. \square

Remark 4.3. If $0 < \alpha \leq 1$, we use $s_\alpha(t) = T(R_t^\alpha)$ ($t > 0$) to obtain

$$\|s_\alpha(t+h) - s_\alpha(t)\| \leq 2\|T\| \frac{h^\alpha}{\Gamma(\alpha + 1)} e^{\omega(t+h)},$$

with $t, h > 0$; see also [7, Corollary 3.3].

ACKNOWLEDGMENT

I thank the referee for careful reading of this paper and valuable suggestions and references that lead to its improvement.

REFERENCES

1. W. Arendt, *Vector-Valued Laplace Transforms and Cauchy Problems*, Israel J. Math. **59**(3) (1987), 327–352. MR0920499 (89a:47064)
2. A. Bobrowski, *On the Yosida Approximation and the Widder-Arendt Representation Theorem*, Studia Math. **124** (1997), 281–290. MR1456426 (99b:47058)
3. W. Chojnacki, *On the Equivalence of a Theorem of Kiszyński and the Hille-Yosida Generation Theorem*, Studia Math. **126**(2) (1998), 491–497. MR1415577 (98d:46050)
4. W. Chojnacki, *Multiplier Algebras, Banach Bundles, and One-Parameter Semigroups*, Ann. Scuola. Norm. Sup. Pisa Cl. Sci. (4) **28** (1999), 287–322. MR1736230 (2002a:46066)
5. W. Chojnacki, *A Generalization of the Widder-Arendt Theorem*, Proc. Edinburgh. Math. Soc. **45** (2002), 161–179. MR1884610 (2002k:44002)
6. B. Hennig and F. Neubrander, *On Representations, Inversions, and Approximations of Laplace Transforms in Banach Spaces*, Appl. Anal. **49** (1993), 151–170. MR1289741 (95m:46053)
7. M. Hieber, *Laplace Transforms and α -Times Integrated Semigroups*, Forum Math **3** (1991), 595–612. MR1130001 (92k:47075)
8. J. Kiszyński, *The Widder Spaces, Representations of the Convolution Algebra $L^1(\mathbb{R}^+)$ and one Parameter Semigroups of Operators*, Preprint no. 558, Institute of Mathematics, Polish Academy of Sciences, Warsaw, June 1998.
9. J. Kiszyński, *Around Widder's Characterization of the Laplace Transform of an Element of $L^\infty(\mathbb{R}^+)$* , Ann. Polon. Math. **74** (2000), 161–200. MR1808994 (2002j:44001)
10. P.J. Miana, *α -Times Integrated Semigroups and Fractional Derivation*, Forum Math. **14** (2002), 23–46. MR1880193 (2003k:47059)

11. K.S. Miller and B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, Wiley, New York, 1993. MR1219954 (94e:26013)
12. D. V. Widder, *The Laplace Transform*, Princeton Univ. Press, 1946. MR0005923 (3:232d)

DEPARTAMENTO DE MATEMÁTICAS, FACULTAD DE CIENCIAS, UNIVERSIDAD DE ZARAGOZA, 50009
ZARAGOZA, SPAIN

E-mail address: `pjmiana@unizar.es`