

TYCHONOFF'S THEOREM FOR LOCALLY COMPACT SPACES AND AN ELEMENTARY APPROACH TO THE TOPOLOGY OF PATH SPACES

ALAN L. T. PATERSON AND AMY E. WELCH

(Communicated by David R. Larson)

ABSTRACT. The path spaces of a directed graph play an important role in the study of graph C^* -algebras. These are topological spaces that were originally constructed using groupoid and inverse semigroup techniques. In this paper, we develop a simple, purely topological, approach to this construction, based on Tychonoff's theorem. In fact, the approach is shown to work even for higher dimensional graphs satisfying the finitely aligned condition, and we construct the groupoid of the graph. Motivated by these path space results, we prove a Tychonoff theorem for an infinite, countable product of locally compact spaces. The main idea is to include certain finite products of the spaces along with the infinite product. We show that the topology is, in a reasonable sense, a pointwise topology.

1. INTRODUCTION

Many of the most important C^* -algebras that have been studied in recent years (e.g. the Cuntz-Krieger algebras) arise in the following context. We are given a directed graph \mathcal{E} with sets of vertices and edges denoted by V, E respectively. Associated with the graph is a universal C^* -algebra $C^*(E)$ and also a reduced C^* -algebra $C_{red}^*(E)$. There is now a large literature studying these graph C^* -algebras and the C^* -algebras of higher dimensional graphs. A sample of papers in this area are: [1, 4, 5, 6, 7, 10, 13, 11, 12, 15, 16, 17, 18, 19, 20, 21, 22, 23]. One of the major approaches to graph C^* -algebras, developed in a number of the papers above, is through the construction of a locally compact groupoid associated with the graph called the *graph groupoid*. (Renault in his book [21] first developed the groupoid approach for the Cuntz graph (below) \mathcal{E}_n .)

The graph groupoid is built up out of paths in \mathcal{E} and indeed in all cases, the unit space of the groupoid is a space of paths. In the row-finite case (i.e. when each vertex emits only finitely many edges) this space is the set Z of all infinite paths in \mathcal{E} . The space Z is a zero-dimensional locally compact space. While Tychonoff's theorem can, in the row-finite case, be easily applied to topologize Z , the case of non-row-finite graphs posed problems in the development of the theory. Indeed, in

Received by the editors January 25, 2004.

2000 *Mathematics Subject Classification*. Primary 54B10, 46L05; Secondary 22A22, 46L85, 54B15.

Key words and phrases. Directed graphs, graph C^* -algebras, path spaces, Tychonoff's theorem.

Cuntz's fundamental paper ([4]), the study of the C^* -algebra O_∞ , corresponding to the graph \mathcal{E}_∞ —an infinite bouquet of circles, not row-finite—requires a somewhat different treatment from that of the C^* -algebra O_n (n finite) whose corresponding graph \mathcal{E}_n is a bouquet of n circles and is row-finite.

Simple examples, such as \mathcal{E}_∞ , illustrate that in the non-row-finite case, Z is not usually locally compact in the pointwise topology. The problem of the existence of a locally compact groupoid associated with a non-row-finite graph was solved by Paterson in [17] using the universal groupoid G associated with the graph inverse semigroup S . (See [16, 4.3] and [9] for the construction of the universal groupoid of an inverse semigroup.) A key step in this procedure is the construction of the unit space X of G . The important part of this unit space can be identified with $W = Y \cup Z$, where Y is the set of finite paths of \mathcal{E} . The space W is then a locally compact metric space, and is used to construct the graph groupoid almost exactly as Z was used in the row-finite construction.

It is natural to ask if it is possible to construct the topology on W using purely topological means, with no recourse to inverse semigroups and their universal groupoids. We will show that this can be done quite simply. Indeed, more generally, this purely topological construction applies, not just to 1-graphs, but even to k -graphs, and this is the setting of the second section. We describe the groupoid for a general, finitely aligned, k -graph¹ and, in particular, the pointwise topology of the unit space W . A related construction, in the context of partial group actions, has been given for Exel-Laca algebras ([6]).

Motivated by the second section, we obtain, in the third section of the paper, an extension of the classical Tychonoff theorem to the case of locally compact, Hausdorff, second countable spaces. It is elementary, of course, that a product of locally compact spaces X_i is locally compact if and only if all but finitely many of the spaces are compact. Our result (Corollary 3.3) requires no such restriction. The crucial point is that we have to include, along with the infinite product $X_1 \times X_2 \times \dots$, certain *finite* products of the spaces. Precisely, we show that there is a natural locally compact metric topology on

$$X = (X_1 \times X_2 \times \dots) \cup \left[\bigcup_{n=1}^{\infty} (X_1 \times X_2 \times \dots \times X_n) \right].$$

The proof uses the one-point compactifications of the X_i 's and quotient topology results.

We have been unable to find this result in topology texts. Something similar, though, appears in the theory of Stochastic processes, where adjunction of the one point at infinity (e.g. [8, p. 324]) gives an absorbing state, so that when a path enters into that state, it is “dead” and therefore can be regarded as a *finite* path. However, we have been unable to find in that literature a detailed account of our result. We also show that the topology on X can be regarded as a “pointwise” topology as in the classical Tychonoff theorem. It is, the authors think, intriguing that the study of groupoids and C^* -algebras, often regarded as *non-commutative* topology, suggests, in a natural way, a result in (commutative) topology itself!

¹In a lecture at the 2003 *Groupoidfest* meeting at the University of Colorado, Paul Muhly described joint work with C. M. Farthing and T. Yeend in which an inverse semigroup approach constructs the groupoid for finitely aligned, higher rank graphs, generalizing the 1-graph case treated in [17]. The first author is grateful to Dr. Muhly for information about this work.

For simplicity, all topological spaces in this paper are locally compact, Hausdorff, second countable (and hence metrizable). All graphs are directed and have countably many vertices and edges. The first author is grateful to Marcelo Laca and Alex Kumjian for helpful discussions concerning the results of this paper.

2. PATH SPACES AND THEIR TOPOLOGIES

In this section, we will construct a locally compact metric topology on the path space of a finitely aligned k -graph Λ . This topology is that of pointwise convergence on the space Y of finite paths in Λ , and is remarkably simple to describe. Local compactness follows from Tychonoff's theorem and we will describe the r -discrete groupoid G_Λ of Λ at the end of the section. (The C^* -algebras of this groupoid are the C^* -algebras of the k -graph.)

We first discuss higher rank graphs (k -graphs). Such graphs Λ were introduced and studied by A. Kumjian and D. Pask in [13]. (Graphs of rank 1 are just directed graphs.) They defined, in terms of generators and relations, the C^* -algebra $C^*(\Lambda)$ of Λ . Further, they showed, under the assumptions of row finiteness and the absence of sources, that the infinite path space Λ^∞ of Λ is a locally compact Hausdorff space. Using shift maps σ^p , they obtained an r -discrete groupoid \mathcal{G}_Λ for Λ , analogous to that for 1-graphs, and showed (among other results) that $C^*(\mathcal{G}_\Lambda) = C^*(\Lambda)$. We will show that their approach to the groupoid \mathcal{G}_Λ can be made to work when Λ is *finitely aligned*. (So Λ need not be row finite and there may be some sources.)

We first recall some definitions ([13]). Let $\mathbb{N} = \{0, 1, 2, \dots\}$ and $k \geq 1$. Then \mathbb{N}^k is ordered by: $m \leq n \iff m_i \leq n_i$ for $1 \leq i \leq k$. Also, if $m, n \in \mathbb{N}^k$, then $m \vee n \in \mathbb{N}^k$ is given by $(m \vee n)_i = \max\{m_i, n_i\}$. Further, regard \mathbb{N}^k as a small category with one object, 0 as identity and addition as composition of morphisms. A k -graph is a countable small category Λ together with a functor $d : \Lambda \rightarrow \mathbb{N}^k$, the *degree map*, with unique factorization, i.e. if $\lambda \in \Lambda$ and $m, n \in \mathbb{N}^k$ is such that $d(\lambda) = m + n$, then there are unique elements $\mu, \nu \in \Lambda$ such that $\lambda = \mu\nu$ with $d(\mu) = m, d(\nu) = n$. We write $\mu \leq \lambda$. For $\lambda \in \Lambda$, let $\Lambda^n = d^{-1}(n)$; then $\Lambda = \bigcup_{n \in \mathbb{N}^k} \Lambda^n$. Further, Λ^0 can be identified with the set of objects of Λ . Note that a 1-graph Λ determines a directed graph with a set of vertices Λ^0 and a set of edges Λ^1 , and Λ^n is just the set of paths of length n . (The reader is referred to [13, 18] for more details.)

In [13], paths in Λ are defined in terms of k -graph morphisms from the category $\Omega_k = \{(m, n) \in \mathbb{N}^k \times \mathbb{N}^k : m \leq n\}$ into Λ . For the purposes of the present paper, it is convenient to reformulate this in terms of sequences in Λ indexed by the elements of certain subsets of \mathbb{N}^k . (This reformulation is effectively contained in the account of [13].) An advantage of this is that we can think of a path in Λ rather as we regard a path in an ordinary directed graph in terms of the sequence of its initial segments increasing with segment length.

Let $N \in (\mathbb{N} \cup \infty)^k$. Then an N -path in Λ is a sequence $\{\lambda_m : m \in \mathbb{N}^k, m \leq N\}$ such that $d(\lambda_m) = m$ and $\lambda_m \leq \lambda_n$ whenever $m \leq n$. The range map on such a path is given by $r(\{\lambda_m\}) = r(\lambda_0)$. The N -paths are identifiable with the k -graph morphisms $x : \Omega(N) \rightarrow \Lambda$, where $\Omega(N)$ is the small category whose set of objects is $O(N) = \{p \in \mathbb{N}^k : p \leq N\}$ and whose morphisms are pairs of the form $(m, n) \in \mathbb{N}^k \times \mathbb{N}^k$, where $m \leq n \leq N$, $r(m, n) = m, s(m, n) = n$ and $(m, n)(n, p) = (m, p)$. Indeed, given an N -path $\{\lambda_m\}$, the morphism x is obtained by factorizing $\lambda_n = \lambda_m x(m, n)$ where $d(\lambda_m) = m, d(x(m, n)) = n - m$, while conversely, given

such a morphism x , then we just take $\lambda_m = x(0, m)$. Let W^N be the set of N -paths in Λ and let

$$W = \bigcup_{N \in (\mathbb{N} \cup \infty)^k} W^N$$

be the *path space* of Λ . If $N \in \mathbb{N}^k$, and $\{\lambda_m\}$ is an N -path, then by the uniqueness of factorization, we can identify the path with the single element $\lambda_N \in \Lambda^N$. So the set Y of such paths can be called the *finite path space*, and can be identified with $\bigcup_{N \in \mathbb{N}^k} \Lambda^N$. The set

$$\bigcup_{N \in (\mathbb{N} \cup \infty)^k \setminus \mathbb{N}^k} W^N$$

is called the *infinite path space* of Λ , and is denoted by Z . (Note that if $N \in (\mathbb{N} \cup \infty)^k \setminus \mathbb{N}^k$, then $N_i = \infty$ for some i , and so $\{\lambda_m\} \in W^N$ has infinitely many components.) So $W = Y \cup Z$. Next, we need to be able to form products of paths yw for $y \in Y, w \in W$ and $s(y) = r(w)$. To define this let $y \in W^N, w \in W^{N'}$ and $n \in \mathbb{N}^k, n \leq N + N'$. Choose $p \in \mathbb{N}^k$ such that $p \leq N', n \leq N + p$. Then $(yw)_n$ is defined by $yw_p = (yw)_n \mu$ where $d((yw)_n) = n, d(\mu) = N + p - n$.

Let $v \in \Lambda^0$. Define

$$W_v = \{\{\lambda_n\} \in W : \lambda_0 = v\}.$$

Then $W = \bigcup_{v \in \Lambda^0} W_v$. We will show that, with a natural topology of “pointwise convergence”, each W_v is a compact Hausdorff space. (So W is a locally compact Hausdorff space with the disjoint union topology.)

We say that Λ is *finitely aligned* ([18, Definition 5.3]) if given $\lambda, \mu \in Y = \Lambda$, the set

$$E_{\lambda, \mu} = \{\nu \in Y : \lambda \leq \nu, \mu \leq \nu, d(\nu) = d(\lambda) \vee d(\mu)\}$$

is finite. As observed in [19], finite alignment is the condition required for the new Cuntz-Krieger relation of [19] to be C^* -algebraic rather than spatial. In the present paper, the condition has a very simple interpretation: it is exactly what is required to make the path space locally compact. (For the 1-graph case, finite alignment is trivially satisfied.)

We now define a map $\alpha : W \rightarrow 2^Y = \{0, 1\}^Y$ by setting $\alpha(w)(y) = 1$ if and only if $y \leq w$. For each $a \in 2^Y$, let $A_a = \{y \in Y : a(y) = 1\}$. It is obvious that α is one-to-one, since if $\alpha(w) = a$, then for each n , w_n is the unique element of A_a in W^n . The next result identifies the image of α in 2^Y .

Lemma 2.1. *Let $a \in 2^Y$. Then $a \in \alpha(W)$ if and only if A_a satisfies the following conditions:*

- (i) $A_a \neq \emptyset$;
- (ii) if $y_1 \leq y_2$ in Y and $y_2 \in A_a$, then $y_1 \in A_a$;
- (iii) A_a is directed upwards, i.e. if $y_1, y_2 \in A_a$, then there exists $y \in A_a$ such that $y_1 \leq y, y_2 \leq y$.

Proof. Suppose that $a \in \alpha(W)$. Then $a = \alpha(w)$ for some $w \in W$. Then $A_a \neq \emptyset$ since it contains w_0 . (ii) follows since if $y_1 \leq y_2$ and $y_2 \leq w$, then $y_1 \leq w$. For (iii), suppose that $y_1, y_2 \in A_a$. Then $y_1 \leq w, y_2 \leq w$. Let $y_1 \in W^{N_1}, y_2 \in W^{N_2}, w \in W^{N_3}$. Then $N_1 \leq N_3, N_2 \leq N_3$. (Note that the entries of N_1, N_2 are finite, but those of N_3 may not be.) Let $N_4 = N_1 \vee N_2 \leq N_3$ and $y = w_{N_4}$. Then $y_1 \leq y, y_2 \leq y$ and $y \in A_a$.

Conversely, suppose that $a \in 2^Y$ satisfies (i), (ii) and (iii). For each $y \in A_a$, let N_y be such that $y \in W^{N_y}$. Define N to be the smallest element of $(\mathbb{N} \cup \infty)^k$ such that $N_y \leq N$ for all $y \in A_a$. (So for each i , $1 \leq i \leq k$, $N_i = \sup_{y \in A_a} (N_y)_i$.) Let $n \in \mathbb{N}^k$, $n \leq N$. Then for each i , there exists $y(i) \in A_a$ such that $n_i \leq (N_{y(i)})_i \leq N_i$. By (iii), there exists $y \in A_a$ such that $y(i) \leq y$ for $1 \leq i \leq k$. Then $n \leq N_y$. Define $w_n = y_n \in \Lambda$. We claim that w is well defined, that $w \in W^N$ and that $\alpha(w) = a$.

To this end, suppose that $n \leq N' \in \mathbb{N}^k$, $N' \leq N$, and that $y' \in W^{N'} \cap A_a$. Then by (iii), there exists $z \in A_a$ with $y \leq z, y' \leq z$. Then $y_n = z_n = y'_n$ so that w_n is well defined. It also follows that $w \in W^N$. We now show that $\alpha(w) = a$. Suppose first that $a(y) = 1$. Then $N_y \leq N$ and by the above, $y \leq w$ and $\alpha(w)(y) = 1$. Conversely, suppose that $\alpha(w)(y) = 1$. Then $y \leq w$. By the above argument, there exists $y' \in A_a$ such that $N_{y'} \geq N_y$. Then $y = w_{N_y} = y'_{N_y} \in A_a$ by (ii). Hence $\alpha(w) = a$. □

Theorem 2.2. *Assume that Λ is finitely aligned. Then for each $v \in \Lambda^0$, the set $\alpha(W_v)$ is a closed subset of 2^Y . So identifying W_v with $\alpha(W_v)$ under the map α , W_v is a compact metric space under the topology of pointwise convergence on Y .*

Proof. Let $\{w^n\}$ be a sequence in W_v such that $\alpha(w^n) \rightarrow a$ in 2^Y . It is sufficient to show that A_a satisfies (i), (ii) and (iii) of Lemma 2.1.

(i) follows since $a(v) = \lim \alpha(w^n)(v) = 1$ since $v \leq w^n$ for all n . For (ii), suppose that $y_1 \leq y_2$ in Y and $y_2 \in A_a$. Since $w^n \rightarrow a$ and $a(y_2) = 1$, it follows that $\alpha(w^n)(y_2) = 1$ eventually. By Lemma 2.1, $\alpha(w^n)(y_1) = 1$ eventually, so that $a(y_1) = 1$, and (ii) follows. Next, for (iii), suppose that $y_1, y_2 \in A_a$. We have to show that there exists $y \in A_a$ such that $y_1 \leq y, y_2 \leq y$. To this end, again since $\alpha(w^n) \rightarrow a$, we have eventually that $\alpha(w^n)(y_1) = 1 = \alpha(w^n)(y_2)$. We can assume that this is true for all n . Let $N = d(y_1) \vee d(y_2)$ and let N_n be such that $w^n \in W^{N_n}$. Then $N_n \geq N$ for all n . Let $y^n = w^n_N$. Then $y^n \in E_{y_1, y_2}$, and since E_{y_1, y_2} is, by finite alignment, finite, we can, by going to a subsequence, assume that $y^n = y$ is constant for all n . Since $1 = \alpha(w^n)(y) \rightarrow a(y)$, we have $a(y) = 1$ and $y \in A_a$. So (iii) is satisfied and $a \in \alpha(W)$. □

Corollary 2.3. *Let Λ be finitely aligned. Then the path space W is a locally compact metric space under the topology of pointwise convergence on Y . This topology is the disjoint union topology of the compact spaces W_v .*

Corollary 2.4. *Let Λ be finitely aligned. Then the family of sets of the form $\lambda W_{s(\lambda)}$ forms a basis of compact open subsets for the topology of W .*

Proof. This follows by intersecting W by basic open subsets of 2^Y . □

Remarks for 1-graphs. Let \mathcal{E} be a directed graph and E, V the sets of edges and vertices respectively. Then Y is the set of finite paths in \mathcal{E} and Z the set of infinite paths in \mathcal{E} . Then $W = Y \cup Z$. The graph \mathcal{E} is said to be *row-finite* if for each $v \in V$, there are only finitely many edges e starting at v . For each $v \in V$, let Z_v, Y_v be the sets of infinite, finite paths $z = z_1 z_2 \dots$ of edges z_i in E starting at v . Then $W_v = Y_v \cup Z_v$ and is a compact open subset of W . If \mathcal{E} is row-finite, then Z_v is a closed subset of W_v , and for every $y \in W$, the singleton set $\{y\}$ is open in W .

Let $\ell(w)$ be the length of the path w ($0 \leq \ell(w) \leq \infty$). Abusing our earlier notation, let w_i be the i th component of a path w . Then a sequence $w^n \rightarrow w$ in W_v if and only if: either (a) $\ell(w) = \infty$ in which case $\ell(w^n) \rightarrow \infty$ and $w_i^n \rightarrow w_i$

for all i , or (b) $\ell(w) < \infty$, in which case eventually, $\ell(w^n) \geq \ell(w)$, $w_i^n \rightarrow w_i$ for $1 \leq i \leq \ell(w)$ and $w_{\ell(w)+1}^n \rightarrow \infty$ (where $w_{\ell(w)+1}^n$ is taken to be ∞ if $\ell(w^n) = \ell(w)$). A similar result concerning convergent sequences can be formulated for finitely aligned k -graphs in general. We will also see this kind of convergence again in the next section within a much more general, topological context.

A good illustration of the preceding comments (and indeed one of the main motivations for the theory developed in this paper) is provided by the Cuntz graph \mathcal{E}_∞ , which has one vertex v and a countably infinite set of loops beginning and ending at v . These loops will be labelled by the set \mathbb{P} of positive integers $1, 2, 3, \dots$. Then

$$(2.1) \quad W = \{v\} \cup \left(\bigcup_{k=1}^\infty \mathbb{P}^k\right) \cup \prod_{n=1}^\infty \mathbb{P} = W_v$$

is a compact metric space. (For an alternative account of the topology on W in this case, see [21, pp. 139-140].) Examples of convergent sequences in W are $123 \dots n \rightarrow 1234 \dots$, $12n111 \dots \rightarrow 12$, and $nnn \dots \rightarrow v$.

Let Λ be a general, finitely aligned k -graph as above. The path groupoid of Λ is then constructed in a manner analogous to that of [13] but taking into account the presence of finite paths. Let G_Λ be the set of triples of the form $(\lambda w, d(\lambda) - d(\mu), \mu w)$ for $\lambda, \mu \in \Lambda$ and $w \in W$ where $r(w) = s(\lambda) = s(\mu)$. Multiplication and inversion on G_Λ are defined in the usual way making it into a groupoid: $(x, n, y)(y, \ell, z) = (x, n + \ell, z)$ and $(x, n, y)^{-1} = (y, -n, x)$. The family of sets

$$Z(\lambda, \mu) = \{(\lambda w, d(\lambda) - d(\mu), \mu w) : w \in W, r(w) = s(\lambda) = s(\mu)\}$$

is a basis for a topology on G_Λ . Indeed, if $(x, n, y) \in Z(\lambda, \mu) \cap Z(\lambda', \mu')$, then $(x, n, y) \in Z(\lambda \vee \lambda', \mu \vee \mu') \subset Z(\lambda, \mu) \cap Z(\lambda', \mu')$. Further, G_Λ is a second countable, ample ([16, p. 48]), locally compact metric groupoid with counting measure as left Haar system (cf. [13, Proposition 2.8]). For example, $Z(\lambda, \mu)$ is homeomorphic to the basic compact, open subsets (Corollary 2.4) $\lambda W_{s(\lambda)}, \mu W_{s(\mu)}$ of the unit space W of G_Λ under the range and source maps: $(\lambda w, d(\lambda) - d(\mu), \mu w) \rightarrow \lambda w, (\lambda w, d(\lambda) - d(\mu), \mu w) \rightarrow \mu w$. It seems very likely (cf. [13]) that the groupoid approach to the C^* -algebras of a 1-graph ([12, 17, 21]) can be extended to the case of finitely aligned k -graphs using the groupoid G_Λ .

3. TYCHONOFF'S THEOREM FOR LOCALLY COMPACT SPACES

The natural locally compact metric topology existing on W_v for \mathcal{E}_∞ ((2.1)) motivates the search for a *locally compact* version of Tychonoff's theorem. We will show in this section that the general version of this, for a product of locally compact spaces, holds. The topological techniques used to prove this generalization are quite different from those of the preceding section.

So let X_n be a second countable, locally compact metric space ($n = 1, 2, 3, \dots$). Let $Y^k = X_1 \times \dots \times X_k, Y_0 = \bigcup_{k=0}^\infty Y^k, Z = \prod_{n=1}^\infty X_n$ and

$$W_0 = Y_0 \cup Z.$$

In the preceding sentence, Y^0 is understood to be $\{0\}$. (0 plays the role of the vertex v in the \mathcal{E}_∞ case.) Let $Y = \bigcup_{k=1}^\infty Y^k$ and $W = Y \cup Z$. So W is obtained from W_0 by removing 0. (Our notation W differs here from that of the preceding section (where vertices were contained in W) but we really want W in the present

context to involve only the X_i 's with no reference to 0.) It will be shown that there is a natural pointwise topology on W_0 which makes W_0 into a compact metric space. Removing the point 0 from W_0 then makes W into a locally compact metric space.

For each n , adjoin a point ∞_n to X_n to give the one-point compactification X_n^∞ of X_n . Then X_n^∞ is a compact metric space ([14, p. 146, Exercise 4]). Let $A = \prod_{n=1}^\infty X_n^\infty$. Then A is a compact metric space by Tychonoff's theorem. We now want to relate A to W_0 in the hope that we can use the compactness of A to obtain a "pointwise" compact topology for W_0 . Unfortunately, there are elements of A that contain ∞_n 's as components, and yet the elements of W_0 do not contain any ∞_n 's. The map $Q : A \rightarrow W_0$, that we now introduce, makes identifications in A that solves this difficulty.

Definition 3.1. The map $Q : A \rightarrow W_0$ is defined as follows:

$$Q(\{x_i\}) = \begin{cases} \{x_i\} & \text{if no } x_i \text{ is } \infty_i, \\ (x_1, x_2, \dots, x_n) & \text{if there is a smallest integer } n \geq 1 \\ & \text{such that } x_{n+1} = \infty_{n+1}, \\ 0 & \text{if } x_1 = \infty_1. \end{cases}$$

It is obvious that Q is onto W_0 and is 1-1 on Z . We give W_0 the quotient topology. For $x = \{x_i\} \in A$ let $N(x)$ be, for the 3 cases of the definition respectively, ∞ , n and 0. Then if $Q(x) = Q(y)$, it follows that $N(x) = N(y)$. The map Q is not 1-1 on A .

Theorem 3.2. W_0 is a compact metric space.

Proof. Let R be the equivalence relation on A determined by Q : so $\{x_i\} \sim \{x'_i\}$ if and only if $Q(\{x_i\}) = Q(\{x'_i\})$. Then $A/R = W_0$, and by [2, p. 105] and [3, p. 159], we have to show that R (identified with its graph) is closed in $A \times A$. To this end let $(x^n, y^n) \rightarrow (x, y) \in A \times A$, where for all n , $(x^n, y^n) \in R$. Write $x^n = \{x_i^n\}$, $y^n = \{y_i^n\}$, $x = \{x_i\}$, $y = \{y_i\}$. Then for all i , $\{x_i^n\} \rightarrow \{x_i\}$, $\{y_i^n\} \rightarrow \{y_i\}$. Also, set $N(x^n) = N(y^n) = m^n$ for all n . We have to show that $(x, y) \in R$. There are two cases to be considered.

Suppose first that the sequence $\{m^n\}$ is bounded. We can, by going to a subsequence, suppose that $m^n = m \geq 0$ is constant. Then $x_{m+1}^n = \infty_{m+1} = y_{m+1}^n$ for all n , and so taking limits, $x_{m+1} = \infty_{m+1} = y_{m+1}$. Next, for $1 \leq j \leq m$, $x_j^n = y_j^n$ for all n by the definition of Q , and taking limits, we get $x_j = y_j$ for $1 \leq j \leq m$. It follows that the first occurrence of an ∞_k in x and y is at exactly the same place, the j th place say, and $1 \leq j \leq m + 1$. So $x_i = y_i$ for $1 \leq i \leq j - 1$, and $(x, y) \in R$.

Next suppose that the sequence $\{m^n\}$ is unbounded. Taking a subsequence, we can, without loss of generality, suppose that $m^n \rightarrow \infty$. Let $i \in \mathbb{P}$. Since $m^n > i$ eventually, we have $x_i^n = y_i^n \in X_i$ for large enough n , and taking limits gives $x_i = y_i \in X_i^\infty$. So $x = y$ and $(x, y) \in R$. \square

Corollary 3.3. The space $W = W_0 \setminus \{0\}$ is a locally compact metric space in the relative quotient topology of W_0 .

For $w \in W_0$ let $\ell(w)$ be the length of w . (We take $\ell(0) = 0$.) Note that for all $x \in A$, $N(x) = \ell(Q(x))$. We now want to show that the topology on W is, in a natural sense, a "pointwise" topology. To this end, we need only check that the convergent sequences in W are of the appropriate form. The form for W_0 (and hence for W) is given in the following definition.

Definition 3.4. Let $\{x^n\}$ ($x^n = \{x_i^n\}$) be a sequence in W_0 and $x \in W_0$. We say that $x^n \rightarrow x$ *pointwise* if the following hold:

- (i) if $x \in Z$, then $\ell(x^n) \rightarrow \infty$ and $x_i^n \rightarrow x_i$ for all i ;
- (ii) if $x \in Y_0$, then $\ell(x^n) \geq \ell(x)$ eventually, $x_i^n \rightarrow x_i$ for $1 \leq i \leq \ell(x)$ and $x_{\ell(x)+1}^n \rightarrow \infty_{\ell(x)+1}$.

(Earlier (§2), we met this kind of convergence for sequences in the path spaces of 1-graphs.) We now want to show that the convergent sequences for W_0 are exactly the pointwise convergent sequences, so that we can regard the topology of W as a natural “pointwise” topology. This will now be proved. For notational ease, we write, for $x \in W_0$, $\bar{x} = Q^{-1}(\{x\}) \subset A$. This is an R -equivalence class in A .

Proposition 3.5. $x^n \rightarrow x$ in $W_0 \iff$ every cluster point of every sequence $\{y^n\}$, where $y^n \in \bar{x}^n$, is in \bar{x} .

Proof. \Rightarrow Suppose that for some sequence $\{y^n\}$, where $y^n \in \bar{x}^n$, there is a subsequence $y^{n_k} \rightarrow y \in A$. Then $x^{n_k} = Q(y^{n_k}) \rightarrow Q(y) = x$ so that $y \in \bar{x}$.

\Leftarrow Suppose that $\{x^n\}$ does not converge to x . By going to a subsequence, we can suppose that there is an open subset U of W_0 with $x \in U$ and $x^n \in W_0 \setminus U$ for all n . Let $\{y^n\}$ be a sequence in A with $y^n \in \bar{x}^n$. By compactness, there exists a cluster point y of $\{y^n\}$. Then for some subsequence $\{y^{n_k}\}$ of $\{y^n\}$, we have $y^{n_k} \rightarrow y$. Then $x^{n_k} = Q(y^{n_k}) \rightarrow Q(y)$. Since x^{n_k} is in the closed set $W_0 \setminus U$, it follows that $Q(y) \in W_0 \setminus U$ and $Q(y) \neq x$. This contradicts $y \in \bar{x}$. \square

Theorem 3.6. $x^n \rightarrow x$ in $W_0 \iff x^n \rightarrow x$ *pointwise*.

Proof. There are two cases to be considered depending on whether $x \in Y_0$ or $x \in Z$. We will give the proof for the first case, the proof for the second case being similar and easier.

Suppose then that $x \in Y_0$ and $x^n \rightarrow x$ in W_0 . Let $y^n \in \bar{x}^n$. Then by Proposition 3.5, every cluster point of $\{y^n\}$ is in \bar{x} . Suppose that it is not true that $\ell(x^n) \geq \ell(x)$ eventually. Then there is a subsequence $\{x^{n_k}\}$ such that $\ell(x^{n_k}) = b < \ell(x)$. So the $(b + 1)$ th entry of each y^{n_k} is ∞_{b+1} . We can suppose by compactness that $y^{n_k} \rightarrow y$ for some $y \in A$. Since this convergence is pointwise in A , it follows that $y_{b+1} = \infty_{b+1}$ so that $y \notin \bar{x}$. This is a contradiction. So $\ell(x^n) \geq \ell(x)$ eventually. Now let $1 \leq i \leq \ell(x)$. We claim that $x_i^n \rightarrow x_i$. For suppose not. Then there exists a subsequence $\{x^{n_k}\}$ and a neighborhood V_i of x_i in X_i such that for all k , $x_i^{n_k} \notin V_i$. Let $y^{n_k} \in \bar{x}^{n_k}$. By the above, without loss of generality, we can suppose that $N(y^{n_k}) \geq \ell(x)$ for all k and $y^{n_k} \rightarrow y$ for some y in A . Then $y_i = \lim x_i^{n_k} \in X^\infty \setminus V_i$. But then $y \notin \bar{x}$ since the i th entry of y does not equal that of x and $i \leq \ell(x)$. This is a contradiction. So $x_i^n \rightarrow x_i$ for $1 \leq i \leq \ell(x)$. A similar argument shows that $x_{\ell(x)+1}^n \rightarrow \infty_{\ell(x)+1}$ so that by (ii) of Definition 3.4, $x^n \rightarrow x$ pointwise.

Conversely, suppose that $x^n \rightarrow x$ pointwise. Let $y^n \in \bar{x}^n$ and $\{y^{n_k}\}$ be a subsequence such that $y^{n_k} \rightarrow y \in A$. By Proposition 3.5, we have to show that $y \in \bar{x}$. To this end, we have that $y_i^{n_k} \rightarrow y_i$ in X_i^∞ for all i . We note that eventually, $N(y^{n_k}) = \ell(x^{n_k}) \geq \ell(x)$ and that for all $1 \leq i \leq \ell(x)$, $y_i^{n_k} = x_i^{n_k} \rightarrow x_i$ so that $y_i = x_i$ for $1 \leq i \leq \ell(x)$. It remains to show that $y_{\ell(x)+1} = \infty_{\ell(x)+1}$, for this would give that $y \in \bar{x}$.

There are two cases to be considered. Suppose first that $y_{\ell(x)+1}^{n_k} = \infty_{\ell(x)+1}$ for infinitely many k . Then by pointwise convergence in A , $y_{\ell(x)+1} = \infty_{\ell(x)+1}$. For the second case, we can suppose, without loss of generality, that $y_{\ell(x)+1}^{n_k} \in X_{\ell(x)+1}$

for all k . Then $N(y^{n_k}) = \ell(x^{n_k}) \geq \ell(x) + 1$. So $x_{\ell(x)+1}^{n_k} = y_{\ell(x)+1}^{n_k} \in X_{\ell(x)+1}$, and $y_{\ell(x)+1} = \lim_{k \rightarrow \infty} y_{\ell(x)+1}^{n_k} = \lim_{k \rightarrow \infty} x_{\ell(x)+1}^{n_k} = \infty_{\ell(x)+1}$. \square

Notes. In the \mathcal{E}^∞ case, i.e. where $X_i = \mathbb{P}$ for all i , there are two ways of defining a compact metric topology on $W_0 (= W_v ((2.1)))$. The first of these, which we will call (W_0, rel) is the relative topology obtained by regarding W_0 as a subset of 2^{Y_0} (using α), discussed in the previous section. The second is $(W_0, quot)$ obtained by giving W_0 the quotient topology coming from the map $Q : A \rightarrow W_0$. As one might expect, these are the same. To see this, let $\beta : A \rightarrow 2^{Y_0}$ be given by $\beta(x)(y) = 1$ if x begins with y and is 0 otherwise. It is easy to check that β is continuous and that $\beta = \alpha \circ Q$. It follows by the definition of the quotient topology that the identity map from $(W_0, quot) \rightarrow (W_0, rel)$ is continuous. An elementary compactness result then gives that the two topologies coincide on W_0 .

In the case of Tychonoff's theorem for locally compact metric spaces (Theorem 3.2), when the X_i 's are all the same space X , it is natural to ask if W is the unit space of some locally compact groupoid. (When $X = \mathbb{P}$, W is the unit space of the Cuntz groupoid in the \mathcal{E}_∞ case.) There is a natural groupoid \mathcal{G}_X whose elements are triples of the form $(yw, \ell(y) - \ell(y'), y'w)$ just as in the graph case. However, we have been unable to topologize this groupoid in any reasonable way.

REFERENCES

- [1] T. Bates, D. Pask, I. Raeburn and W. Szymanski, *The C^* -algebras of row-finite graphs*, New York J. Math. 6(2000), 307-324. MR1777234 (2001k:46084)
- [2] N. Bourbaki, *Elements of Mathematics, General Topology, Chapters 1-4*, Springer-Verlag, New York, 1989. MR0979294 (90a:54001a)
- [3] N. Bourbaki, *Elements of Mathematics, General Topology, Chapters 5-10*, Springer-Verlag, New York, 1989. MR0979295 (90a:54001b)
- [4] J. Cuntz, *Simple C^* -algebras generated by isometries*, Comm. Math. Phys. 57(1977), 173-185. MR0467330 (57:7189)
- [5] J. Cuntz and W. Krieger, *A class of C^* -algebras and topological Markov chains*, Invent. Math. 56(1980), 251-268. MR0561974 (82f:46073a)
- [6] R. Exel and M. Laca, *Cuntz-Krieger algebras for infinite matrices*, J. Reine Angew. Math. 512 (1999), 119-172. MR1703078 (2000i:46064)
- [7] A. an Huef and I. Raeburn, *The ideal structure of Cuntz-Krieger algebras*, Ergodic Theory and Dynamical Systems 17(1997), 611-624. MR1452183 (98k:46098)
- [8] O. Kallenberg, *Foundations of Modern Probability*, Springer, New York, 1997. MR1464694 (99e:60001)
- [9] M. Khoshkam and G. Skandalis, *Regular Representation of Groupoids and Applications to Inverse Semigroups*, J. Reine Angew. Math. 546(2002), 47-72. MR1900993 (2003f:46084)
- [10] A. Kumjian, *Notes on C^* -algebras of graphs*, Contemporary Mathematics 228(1998), 189-200. MR1667662 (99m:46137)
- [11] A. Kumjian, D. Pask and I. Raeburn, *Cuntz-Krieger algebras of directed graphs*, Pacific J. Math. 184(1998), 161-174. MR1626528 (99i:46049)
- [12] A. Kumjian, D. Pask, I. Raeburn and J. Renault, *Graphs, Groupoids and Cuntz-Krieger algebras*, J. Funct. Anal. 144(1997), 505-541. MR1432596 (98g:46083)
- [13] A. Kumjian and D. Pask, *Higher Rank Graph C^* -algebras*, New York Journal of Mathematics 6(2000), 1-20. MR1745529 (2001b:46102)
- [14] J. Nagata, *Modern General Topology*, North Holland, Amsterdam, 1968. MR0264579 (41:9171)
- [15] D.A. Pask and C.E. Sutherland, *Filtered inclusions of path algebras: a combinatorial approach to Doplicher-Roberts duality*, J. Operator Theory 31 (1994) 99-121. MR1316986 (96i:46073)
- [16] A.L.T. Paterson *Groupoids, inverse semigroups and their operator algebras*, Progress in Mathematics, v. 170, Birkäuser Boston, New York, 1999. MR1724106 (2001a:22003)

- [17] A.L.T. Paterson *Graph inverse semigroups, groupoids, and their C^* -algebras*, J. Operator Theory 48(2002), 645-662. MR1962477 (2004h:46066)
- [18] I. Raeburn and A. Sims, *Product systems of graphs and the Toeplitz algebras of higher-rank graphs*, preprint, 2001.
- [19] I. Raeburn, A. Sims and T. Yeend, *The C^* -algebras of finitely aligned higher-rank graphs*, J. Funct. Anal. 213(2004), 206-240. MR2069786
- [20] I. Raeburn, A. Sims and T. Yeend, *Higher-rank graphs and their C^* -algebras*, Proc. Edinburgh Math. Soc. 46(2003), 99-115. MR1961175 (2004f:46068)
- [21] J. N. Renault, *A groupoid approach to C^* -algebras*, Lecture Notes in Mathematics, Vol. 793, Springer-Verlag, New York, 1980. MR0584266 (82h:46075)
- [22] W. Szymanski, *Simplicity of Cuntz-Krieger algebras of infinite matrices*, Pacific J. Math. 199(2001) 249-256. MR1847155 (2002i:46056)
- [23] W. Szymanski, *Bimodules for Cuntz-Krieger algebras of infinite matrices*, Bull. Austral. Math. Soc. 62(2000), 87-94. MR1775890 (2001g:46151)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSISSIPPI, UNIVERSITY, MISSISSIPPI 38677
E-mail address: mmap@olemiss.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSISSIPPI, UNIVERSITY, MISSISSIPPI 38677
E-mail address: amy3welch@yahoo.com