

ON THE STRUCTURE OF QUANTUM PERMUTATION GROUPS

TEODOR BANICA AND SERGIU MOROIANU

(Communicated by Martin Lorenz)

ABSTRACT. The quantum permutation group of the set $X_n = \{1, \dots, n\}$ corresponds to the Hopf algebra $A_{aut}(X_n)$. This is an algebra constructed with generators and relations, known to be isomorphic to $\mathbb{C}(S_n)$ for $n \leq 3$, and to be infinite dimensional for $n \geq 4$. In this paper we find an explicit representation of the algebra $A_{aut}(X_n)$, related to Clifford algebras. For $n = 4$ the representation is faithful in the discrete quantum group sense.

INTRODUCTION

A general theory of unital Hopf \mathbb{C}^* -algebras was developed by Woronowicz in [11], [12], [13]. The main results are the existence of the Haar functional, an analogue of Peter-Weyl theory and of Tannaka-Krein duality, and explicit formulae for the square of the antipode. As for examples, these include algebras of continuous functions on compact groups, q -deformations of them with $q > 0$, and \mathbb{C}^* -algebras of discrete groups.

Of particular interest is the algebra $A_{aut}(X_n)$ constructed by Wang in [9]. This is the universal Hopf \mathbb{C}^* -algebra coacting on the set $X_n = \{1, \dots, n\}$. In other words, the compact quantum group associated to it is a kind of analogue of the symmetric group S_n .

The algebra $A_{aut}(X_n)$ is constructed with generators and relations. There are n^2 generators, labeled u_{ij} with $i, j = 1, \dots, n$. The relations are those making u a magic biunitary matrix. This means that all coefficients u_{ij} are projections, and on each row and each column of u these projections are mutually orthogonal, and sum up to 1. The comultiplication is given by $\Delta(u_{ij}) = \sum u_{ik} \otimes u_{kj}$, and the fundamental coaction is given by $\alpha(\delta_i) = \sum u_{ji} \otimes \delta_j$.

For $n = 1, 2, 3$ the canonical quotient map $A_{aut}(X_n) \rightarrow \mathbb{C}(S_n)$ is an isomorphism. For $n \geq 4$ the algebra $A_{aut}(X_n)$ is infinite dimensional, and just a few things are known about it. Its irreducible corepresentations are classified in [3], with the conclusion that their fusion rules coincide with those for irreducible representations of $SO(3)$, independently of $n \geq 4$. In [10] Wang proves that the compact quantum

Received by the editors November 24, 2004 and, in revised form, July 28, 2005.

2000 *Mathematics Subject Classification*. Primary 16W30; Secondary 81R50.

Key words and phrases. Hopf algebra, magic biunitary matrix, inner faithful representation.

The second author was partially supported by the Marie Curie grant MERG-CT-2004-006375 funded by the European Commission, and by a CERES contract (2004).

©2006 American Mathematical Society
Reverts to public domain 28 years from publication

group associated to $A_{aut}(X_n)$ with $n \geq 4$ is simple. In [3] it is shown that the discrete quantum group associated to $A_{aut}(X_n)$ with $n \geq 5$ is not amenable. Various quotients of $A_{aut}(X_n)$, corresponding to quantum symmetry groups of polyhedra, colored graphs etc., are studied in [4] by using planar algebra techniques.

These results certainly bring some light on the structure of $A_{aut}(X_n)$. However, for $n \geq 4$ this remains an abstract algebra, constructed with generators and relations.

In this paper we find an explicit representation of $A_{aut}(X_n)$. The construction works when n is a power of 2, and uses a magic biunitary matrix related to Clifford algebras. For $n = 4$ the representation is inner faithful, in the sense that the corresponding unitary representation of the discrete quantum group associated to $A_{aut}(X_4)$ is faithful.

As a conclusion, there might be a geometric interpretation of Hopf algebras of type $A_{aut}(X_n)$. We should mention here that for the algebra $A_{aut}(X)$ with an X finite graph, such an interpretation would be of real help, for instance in computing fusion rules.

1. MAGIC BIUNITARY MATRICES

Let A be a unital \mathbb{C}^* -algebra. That is, we have a unital algebra A over the field of complex numbers \mathbb{C} , with an antilinear antimultiplicative map $a \rightarrow a^*$ satisfying $a^{**} = a$, and with a Banach space norm satisfying $\|a^*a\| = \|a\|^2$.

A projection is an element $p \in A$ satisfying $p^2 = p^* = p$. Two projections p, q are said to be orthogonal when $pq = 0$. A partition of unity in A is a finite set of projections, which are mutually orthogonal, and sum up to 1.

Definition 1.1. A matrix $v \in M_n(A)$ is called magic biunitary if all its rows and columns are partitions of the unity of A .

A magic biunitary is indeed a biunitary, in the sense that both v and its transpose v^t are unitaries. The other word – magic – comes from a vague similarity with magic squares.

The basic example comes from the symmetric group S_n . Consider the sets of permutations $\{\sigma \in S_n \mid \sigma(j) = i\}$. When i is fixed and j varies, or vice versa, these sets form partitions of S_n . Thus their characteristic functions $v_{ij} \in \mathbb{C}(S_n)$ form a magic biunitary.

Of particular interest is the “universal” magic biunitary matrix. This has coefficients in the universal algebra $A_{aut}(X_n)$ constructed by Wang in [9].

Definition 1.2. $A_{aut}(X_n)$ is the universal \mathbb{C}^* -algebra generated by n^2 elements u_{ij} , subject to the magic biunitarity condition.

In other words, we have the following universal property. For any magic biunitary matrix $v \in M_n(A)$ there is a morphism of \mathbb{C}^* -algebras $A_{aut}(X_n) \rightarrow A$ mapping $u_{ij} \rightarrow v_{ij}$.

A more elaborate version of this property, to be discussed now, states that $A_{aut}(X_n)$ is a Hopf \mathbb{C}^* -algebra, whose underlying quantum group is a kind of analogue of S_n .

The following definition is due to Woronowicz [13].

Definition 1.3. A unital Hopf \mathbb{C}^* -algebra is a unital \mathbb{C}^* -algebra A , together with a morphism of \mathbb{C}^* -algebras $\Delta : A \rightarrow A \otimes A$, subject to the following conditions.

- (i) Coassociativity condition: $(\Delta \otimes id)\Delta = (id \otimes \Delta)\Delta$.
- (ii) Cocancellation condition: $\Delta(A)(1 \otimes A)$ and $\Delta(A)(A \otimes 1)$ are dense in $A \otimes A$.

The basic example is the algebra $\mathbb{C}(G)$ of continuous functions on a compact group G , with $\Delta(\varphi) : (g, h) \rightarrow \varphi(gh)$. Here coassociativity of Δ follows from associativity of the multiplication of G , and cocancellation in $\mathbb{C}(G)$ follows from cancellation in G .

Another example is the group algebra $\mathbb{C}^*(\Gamma)$ of a discrete group Γ . This is obtained from the usual group algebra $\mathbb{C}[\Gamma]$ by a standard completion procedure. The comultiplication is defined on generators $g \in \Gamma$ by the formula $\Delta(g) = g \otimes g$.

In general, associated to a Hopf \mathbb{C}^* -algebra A are a compact quantum group G and a discrete quantum group Γ , according to the heuristic formula $A = \mathbb{C}(G) = \mathbb{C}^*(\Gamma)$.

Definition 1.4. A coaction of A on a finite set X is a morphism of \mathbb{C}^* -algebras $\alpha : \mathbb{C}(X) \rightarrow \mathbb{C}(X) \otimes A$, subject to the following conditions:

- (i) Coassociativity condition: $(\alpha \otimes id)\alpha = (id \otimes \Delta)\alpha$.
- (ii) Natural condition: $(\Sigma \otimes id)v = \Sigma(\cdot)1$, where $\Sigma(\varphi)$ is the sum of values of φ .

The basic example is with a group G of permutations of X . Consider the action map $a : X \times G \rightarrow X$, given by $a(i, \sigma) = \sigma(i)$. The formula $\alpha\varphi = \varphi a$ defines a morphism of \mathbb{C}^* -algebras $\alpha : \mathbb{C}(X) \rightarrow \mathbb{C}(X \times G)$. This can be regarded as a coaction of $\mathbb{C}(G)$ on X .

In general, coactions of A can be thought of as coming from actions of the underlying compact quantum group G . With this interpretation, the natural condition says that the action of G must preserve the counting measure on X . This assumption cannot be dropped.

The following fundamental result is due to Wang [9].

- Theorem 1.1.** (i) $A_{aut}(X_n)$ is a Hopf \mathbb{C}^* -algebra, with comultiplication $\Delta(u_{ij}) = \sum u_{ik} \otimes u_{kj}$.
- (ii) The linear map $\alpha(\delta_j) = \sum \delta_i \otimes u_{ji}$ is a coaction of $A_{aut}(X_n)$ on $X_n = \{1, \dots, n\}$.
- (iii) $A_{aut}(X_n)$ is the universal Hopf \mathbb{C}^* -algebra coacting on X_n .

The idea for proving (i) is that we can define Δ by using the universal property of $A_{aut}(X_n)$. Coassociativity is clear, and cocancellation follows from a result of Woronowicz in [13], stating that this is automatic whenever there is a counit and an antipode. But these can be defined by $\varepsilon(u_{ij}) = \delta_{ij}$ and $S(u_{ij}) = u_{ji}$, once again by using universality of $A_{aut}(X_n)$.

We know that the compact quantum group G_n associated to $A_{aut}(X_n)$ is a kind of quantum analogue of the symmetric group S_n . In particular there should be an inclusion $S_n \subset G_n$. Here is the exact formulation of this observation; see Wang [9] for details.

Proposition 1.1. There is a Hopf \mathbb{C}^* -algebra morphism $\pi_n : A_{aut}(X_n) \rightarrow \mathbb{C}(S_n)$, mapping the generators u_{ij} to the characteristic functions of the sets $\{\sigma \in S_n \mid \sigma(j) = i\}$.

The question is now whether π_n is an isomorphism or not. For instance a 2×2 magic biunitary must be of the following special form, where p is a projection:

$$\begin{pmatrix} p & 1-p \\ 1-p & p \end{pmatrix}.$$

The algebra generated by p is canonically isomorphic to \mathbb{C}^2 if $p \neq 0, 1$, and to \mathbb{C} if not. Thus the universal algebra $A_{aut}(X_2)$ is isomorphic to \mathbb{C}^2 , and π_2 is an isomorphism.

The map π_3 is an isomorphism as well; see [4] for a proof.

At $n = 4$ we have the following example of a magic biunitary matrix:

$$\begin{pmatrix} p & 1-p & 0 & 0 \\ 1-p & p & 0 & 0 \\ 0 & 0 & q & 1-q \\ 0 & 0 & 1-q & q \end{pmatrix}.$$

We can choose the projections p, q such that the algebra $\langle p, q \rangle$ they generate is infinite dimensional and not commutative. It follows that $A_{aut}(X_4)$ is infinite dimensional and not commutative as well, so π_4 cannot be an isomorphism.

Proposition 1.2. *For $n \geq 4$ the algebra $A_{aut}(X_n)$ is infinite dimensional and not commutative. In particular π_n is not an isomorphism.*

This follows by gluing an identity matrix of size $n - 4$ to the above 4×4 matrix.

There is a quantum group interpretation here. Consider the compact and discrete quantum groups defined by the formula $A_{aut}(X_4) = \mathbb{C}(G_4) = \mathbb{C}^*(\Gamma_4)$. When p, q are free the surjective morphism of \mathbb{C}^* -algebras $A_{aut}(X_4) \rightarrow \langle p, q \rangle$ can be thought of as coming from a surjective morphism of discrete quantum groups $\Gamma_4 \rightarrow \mathbb{Z}_2 * \mathbb{Z}_2$. This makes it clear that Γ_4 is infinite. Now G_4 being the Pontrjagin dual of Γ_4 , it must be infinite as well.

See Bichon [5], Wang [9], [10] and Banica [4] for further speculations on this subject.

2. INNER FAITHFUL REPRESENTATIONS

We would like to find an explicit representation of $A_{aut}(X_n)$. As with any Hopf \mathbb{C}^* -algebra, there is a problem here, because there are two notions of faithfulness.

Consider for instance a discrete subgroup Γ of the unitary group $U(n)$. The inclusion $\Gamma \subset U(n)$ can be regarded as a unitary group representation $\Gamma \rightarrow U(n)$, and we get a \mathbb{C}^* -algebra representation $\mathbb{C}^*(\Gamma) \rightarrow M_n(\mathbb{C})$. This latter representation is far from being faithful: for instance its kernel is infinite dimensional, hence non-empty, when Γ is an infinite group. However, the representation $\mathbb{C}^*(\Gamma) \rightarrow M_n(\mathbb{C})$ must be “inner faithful” in some Hopf \mathbb{C}^* -algebra sense, because the representation $\Gamma \rightarrow U(n)$ it comes from is faithful.

So, we are led to the following question. Let H be a unital Hopf \mathbb{C}^* -algebra, and let $\pi : H \rightarrow A$ be a morphism of \mathbb{C}^* -algebras. If Γ is the discrete quantum group associated to H we know that π corresponds to a unitary representation $\pi_i : \Gamma \rightarrow U(A)$. The question is: when is π inner faithful, meaning that π_i is faithful?

A simple answer is obtained by using the formalism of Kustermans and Vaes [6]. Associated to H is a von Neumann algebra H_{vN} , obtained by a certain completion procedure. Now coefficients of π belong to the dual algebra \widehat{H}_{vN} , and we can say that π is inner faithful if these coefficients generate \widehat{H}_{vN} . This notion is used by Vaes in [7], and a version of it is used by Wang in [9].

In this paper we use an equivalent definition from [2].

Definition 2.1. Let H be a unital Hopf \mathbb{C}^* -algebra. A \mathbb{C}^* -algebra representation $\pi : H \rightarrow A$ is called inner faithful if the $*$ -algebra generated by its coefficients is dense in H_{alg}^* .

Here H_{alg} is the dense $*$ -subalgebra of H consisting of “representative functions” on the underlying compact quantum group, constructed by Woronowicz in [13]. This is a Hopf $*$ -algebra in the usual sense. Its dual complex vector space H_{alg}^* is a $*$ -algebra, with multiplication Δ^* and involution $**$. Finally, coefficients of π are the linear forms $\varphi\pi$ with $\varphi \in A^*$, and the density assumption is with respect to the weak topology on H_{alg}^* . See e.g. the book of Abe [1] for Hopf algebras and [2] for details regarding this definition.

The main example is with a discrete group Γ . As expected, a representation $\mathbb{C}^*(\Gamma) \rightarrow A$ is inner faithful if and only if the corresponding unitary group representation $\Gamma \rightarrow U(A)$ is faithful. Some other examples are discussed in [2].

For how to use inner faithfulness see Vaes [7].

Definition 2.2. The character of a magic biunitary matrix $v \in M_n(A)$ is the sum of its diagonal entries $\chi(v) = v_{11} + v_{22} + \dots + v_{nn}$.

The terminology comes from the case where $v = u$ is the universal magic biunitary matrix, with coefficients in $A = A_{aut}(X_n)$. Indeed, the matrix u is a corepresentation of $A_{aut}(X_n)$ in the sense of Woronowicz [11], and the element $\chi(u)$ is its character.

Lemma 2.1. Let $v \in M_n(A)$ be a magic biunitary matrix, with $n \geq 4$. Assume that there is a unital linear form $\varphi : A \rightarrow \mathbb{C}$ such that

$$\varphi(\chi(v)^k) = \frac{1}{k+1} \binom{2k}{k}$$

for any k . Then the representation $\pi : A_{aut}(X_n) \rightarrow A$ defined by $u_{ij} \rightarrow v_{ij}$ is inner faithful.

Proof. The numbers in the statement are the Catalan numbers, appearing as multiplicities in the representation theory of $SO(3)$. The result will follow from the following fact from [3]. The finite-dimensional irreducible corepresentations of $A_{aut}(X_n)$ can be arranged in a sequence $\{r_k\}$, such that their fusion rules are the same as those for representations of $SO(3)$:

$$r_k \otimes r_s = r_{|k-s|} + r_{|k-s|+1} + \dots + r_{k+s}.$$

Let $h : A_{aut}(X_n) \rightarrow \mathbb{C}$ be the Haar functional constructed by Woronowicz in [11]. Also consider the character of the fundamental corepresentation of $A_{aut}(X_n)$:

$$\chi(u) = u_{11} + u_{22} + \dots + u_{nn}.$$

The Poincaré series of $A_{aut}(X_n)$ is defined by the following formula:

$$f(z) = \sum_{k=0}^{\infty} h(\chi(u)^k) z^k.$$

By the above result, this is equal to the Poincaré series for $SO(3)$:

$$f(z) = \sum_{k=0}^{\infty} \frac{1}{k+1} \binom{2k}{k} z^k.$$

The assumption of the lemma says that the equality $\varphi\pi = h$ holds on all powers of $\chi(u)$. By linearity, this equality must hold on the algebra $\langle\chi(u)\rangle$ generated by $\chi(u)$. Now by positivity of h it follows that the restriction of π to this algebra $\langle\chi(u)\rangle$ is injective.

On the other hand, once again from fusion rules, we see that $\chi(u)$ generates the algebra of characters $A_{aut}(X_n)_{central}$ constructed by Woronowicz in [11].

Summing up, we know that π is faithful on $A_{aut}(X_n)_{central}$.

Now consider the “minimal model” construction in [2]. This is the factorisation of π into a Hopf \mathbb{C}^* -algebra morphism $A_{aut}(X_n) \rightarrow H$, and an inner faithful representation $H \rightarrow A$:

$$A_{aut}(X_n) \rightarrow H \rightarrow A.$$

Since π is faithful on $A_{aut}(X_n)_{central}$, so is the map on the left. By Woronowicz’s analogue of the Peter-Weyl theory in [11], it follows that the map on the left is an isomorphism. Thus π coincides with the map on the right, which is by definition inner faithful. \square

It is possible to reformulate this result by using notions from Voiculescu’s free probability theory [8]. A non-commutative \mathbb{C}^* -probability space is a pair (A, φ) consisting of a unital \mathbb{C}^* -algebra A together with a positive unital linear form $\varphi : A \rightarrow \mathbb{C}$.

Associated to a self-adjoint element $x \in A$ is its spectral measure μ_x . This is a probability measure on the spectrum of x , defined by the formula

$$\varphi(f(x)) = \int_{\mathbb{R}} f(t) d\mu_x(t).$$

This equality must hold for any continuous function f on the spectrum of x . By density we can restrict our attention to polynomials $f \in \mathbb{C}[X]$, then by linearity it is enough to have this equality for monomials $f(t) = t^k$. We say that μ_x is uniquely determined by its moments,

$$\varphi(x^k) = \int_{\mathbb{R}} t^k d\mu_x(t).$$

The following notion plays a central role in free probability. See [8], page 26.

Definition 2.3. An element x in a non-commutative \mathbb{C}^* -probability space is called semicircular if its spectral measure is $d\mu_x(t) = (2\pi)^{-1}\sqrt{4-t^2} dt$ on $[-2, 2]$, and 0 elsewhere.

In terms of moments, we must have the following equalities, for any k :

$$\varphi(x^k) = \frac{1}{2\pi} \int_{-2}^2 t^k \sqrt{4-t^2} dt.$$

The integral is 0 when k is odd, and equal to a Catalan number when k is even,

$$\varphi(x^{2k}) = \frac{1}{k+1} \binom{2k}{k}.$$

We get in this way a reformulation of the above lemma.

Theorem 2.1. *A magic biunitary matrix whose character has the same spectral measure as the square of a semicircular element produces an inner faithful representation of Wang’s algebra.*

The assumption $n \geq 4$ was removed, because it is superfluous. Indeed, for $n = 1, 2, 3$ finite dimensionality of $A_{aut}(X_n)$ implies that the spectrum of any $\chi(v)$ is discrete.

3. GEOMETRIC CONSTRUCTIONS

Consider the Pauli matrices,

$$1 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad i = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad j = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad k = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}.$$

These satisfy the relations for quaternions $i^2 = j^2 = k^2 = -1$, $ij = ji = -k$, etc.

To any $x \in SU(2)$ we associate the following matrix:

$$\begin{pmatrix} 1 \\ i \\ j \\ k \end{pmatrix} x \begin{pmatrix} 1 & i & j & k \end{pmatrix} = \begin{pmatrix} x & xi & xj & xk \\ ix & ixi & ixj & ixc \\ jx & jxi & jxj & jxc \\ kx & kxi & kxj & kxc \end{pmatrix}.$$

Each row and each column of this matrix is an orthogonal basis of $M_2(\mathbb{C}) \simeq \mathbb{C}^4$ with respect to the inner product

$$\langle x, y \rangle = \frac{1}{2} \text{Tr}(xy^*),$$

since i, j, k are skew-adjoints. Thus the matrix of corresponding orthogonal projections is a magic biunitary.

Theorem 3.1. *There is an inner faithful representation*

$$\pi : A_{aut}(X_4) \rightarrow \mathbb{C}(SU(2), M_4(\mathbb{C}))$$

mapping the universal 4×4 magic biunitary matrix to the 4×4 matrix

$$v(x) = \begin{pmatrix} P_x & P_{xi} & P_{xj} & P_{xk} \\ P_{ix} & P_{ixi} & P_{ixj} & P_{ixk} \\ P_{jx} & P_{jxi} & P_{jxj} & P_{jxk} \\ P_{kx} & P_{kxi} & P_{kxj} & P_{kxk} \end{pmatrix},$$

where for $y \in SU(2)$ we denote by P_y the orthogonal projection onto the space $\mathbb{C}y \subset M_2(\mathbb{C})$, and we regard it as a continuous function of y , with values in $M_2(M_2(\mathbb{C})) \simeq M_4(\mathbb{C})$.

Proof. We have to compute the character of $v = v(x)$:

$$\chi(v) = P_x + P_{ixi} + P_{jxj} + P_{kxk}.$$

We make the convention that Greek letters designate quaternions in $\{1, i, j, k\}$. We decompose x as a sum with real coefficients $x = \sum x_\alpha \alpha$. We have the following formula for $\chi(v)$:

$$\chi(v) = \sum_{\alpha} P_{\alpha x \alpha}.$$

With the notations $\alpha\beta = (-1)^{N(\alpha, \beta)}\beta\alpha$ and $\alpha^2 = (-1)^{N(\alpha)}$ we can compute $\alpha x \alpha$:

$$\alpha x \alpha = \sum_{\beta} (-1)^{N(\alpha, \beta) + N(\alpha)} x_{\beta} \beta.$$

Now using the above-mentioned canonical scalar product on $M_2(\mathbb{C})$, this gives the following formula for $P_{\alpha x \alpha}$, after cancelling the $(-1)^{2N(\alpha)} = 1$ term:

$$\langle P_{\alpha x \alpha} \beta, \gamma \rangle = (-1)^{N(\alpha, \beta) + N(\alpha, \gamma)} x_\beta x_\gamma.$$

Now summing over α gives the formula of the character $\chi(v)$:

$$\langle \chi(v) \beta, \gamma \rangle = \sum_{\alpha} (-1)^{N(\alpha, \beta) + N(\alpha, \gamma)} x_\beta x_\gamma.$$

The coefficient of $x_\beta x_\gamma$ can be computed by using the multiplication table of quaternions,

$$\sum_{\alpha} (-1)^{N(\alpha, \beta) + N(\alpha, \gamma)} = 4\delta_{\beta, \gamma}.$$

Thus $\chi(v)$ is a diagonal matrix, having the numbers $4x_\beta^2$ on the diagonal:

$$\chi(v) = \text{diag}(4x_\beta^2).$$

Consider the linear form $\varphi = \int \otimes tr$, where the integral is with respect to the Haar measure of $SU(2)$, and tr is the normalised trace of 4×4 matrices, meaning $1/4$ times the usual trace. The moments of $\chi(v)$ with respect to φ are computed as follows:

$$\int tr(\chi(v)^k) dx = 4^{k-1} \sum_{\beta} \int x_\beta^{2k} dx.$$

By symmetry reasons the four integrals are all equal, say to the first one,

$$\int tr(\chi(v)^k) dx = 4^k \int x_1^{2k} dx.$$

It follows that $\chi(v)$ has the same spectral measure as $4x_1^2$,

$$\mu_{\chi(v)} = \mu_{4x_1^2}.$$

But the variable $2x_1$ is semicircular. This can be seen in many ways, for instance by direct computation, after identifying $SU(2)$ with the real sphere S^3 , or by using the fact that $2x_1 = \text{Tr}(x)$ is the character of the fundamental representation of $SU(2)$, whose moments are computed using Clebsch-Gordon rules. The result now follows by applying Theorem 2.1. \square

The construction of π has the following generalisation. Consider the Clifford algebra $Cl(\mathbb{R}^s)$. This is a finite-dimensional algebra, having a basis formed by products $e_{i_1} \dots e_{i_k}$ with $1 \leq i_1 < \dots < i_k \leq s$, with multiplicative structure given by $e_i^2 = -1$ and $e_i e_j = -e_j e_i$ for $i \neq j$.

It is convenient to use the notation $e_I = e_{i_1} \dots e_{i_k}$ with $I = (i_1, \dots, i_k)$.

As an example, the Clifford algebra $Cl(\mathbb{R}^2)$ is spanned by the elements $e_\emptyset = 1$, e_1 , e_2 and $e_{12} = e_1 e_2$. The generators e_1, e_2 are subject to the relations $e_1^2 = e_2^2 = -1$ and $e_1 e_2 = -e_2 e_1$. Now these relations are satisfied by the Pauli matrices i, j , and the corresponding representation of $Cl(\mathbb{R}^2)$ turns out to be faithful. That is, we have the following identifications:

$$e_\emptyset = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad e_1 = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad e_{12} = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}.$$

We can label as well indices of 4×4 matrices by elements of the set $\{\emptyset, 1, 2, 12\}$. With these notations, the representation in Theorem 3.1 is given by $\pi(u_{IJ}) = P_{e_I x e_J}$.

The same formula works for an arbitrary number s .

Theorem 3.2. *There is a representation $\pi_n : A_{aut}(X_n) \rightarrow \mathbb{C}(G_n, M_n(\mathbb{C}))$ mapping the universal $n \times n$ magic biunitary matrix to the $n \times n$ matrix*

$$v = (P_{e_I x e_J})_{IJ}$$

where $n = 2^s$, the unitary group of the Clifford algebra $Cl(\mathbb{R}^s)$ is denoted G_n , and the algebra of endomorphisms of $Cl(\mathbb{R}^s)$ is identified with $M_n(\mathbb{C})$.

The first part of proof of Theorem 3.1 extends to this general situation. We get that $\chi(v)$ is diagonal, with eigenvalues $\{nx_I^2\}$. This does not seem to be related to semicircular elements when $s \geq 3$. The representation π_n probably comes from an inner faithful representation of a quotient of $A_{aut}(X_n)$, corresponding to a “subgroup” of the quantum permutation group.

ACKNOWLEDGMENTS

We are deeply grateful to Georges Skandalis for all the discussions and his essential contribution to the results of this paper.

REFERENCES

- [1] E. Abe, Hopf algebras, Cambridge Univ. Press (1977). MR0594432 (83a:16010)
- [2] T. Banica, Hopf algebras and subfactors associated to vertex models, *J. Funct. Anal.* **159** (1998), 243–266. MR1654119 (2001a:46069)
- [3] T. Banica, Symmetries of a generic coaction, *Math. Ann.* **314** (1999), 763–780. MR1709109 (2001g:46146)
- [4] T. Banica, Quantum automorphism groups of homogeneous graphs, *J. Funct. Anal.* **224** (2005), 243–280. MR2146039
- [5] J. Bichon, Quantum automorphism groups of finite graphs, *Proc. Amer. Math. Soc.* **131** (2003), 665–673. MR1937403 (2003j:16049)
- [6] J. Kustermans and S. Vaes, Locally compact quantum groups in the von Neumann algebraic setting, *Math. Scand.* **92** (2003), 68–92. MR1951446 (2003k:46081)
- [7] S. Vaes, Strictly outer actions of groups and quantum groups, *J. Reine Angew. Math.* **578** (2005), 147–184. MR2113893 (2005k:46167)
- [8] D. Voiculescu, K. Dykema and A. Nica, Free random variables, CRM Monograph Series 1, AMS (1993). MR1217253 (94c:46133)
- [9] S. Wang, Quantum symmetry groups of finite spaces, *Comm. Math. Phys.* **195** (1998), 195–211. MR1637425 (99h:58014)
- [10] S. Wang, Simple compact quantum groups I, preprint.
- [11] S.L. Woronowicz, Compact matrix pseudogroups, *Comm. Math. Phys.* **111** (1987), 613–665. MR0901157 (88m:46079)
- [12] S.L. Woronowicz, Tannaka-Krein duality for compact matrix pseudogroups. Twisted $SU(N)$ groups, *Invent. Math.* **93** (1988), 35–76. MR0943923 (90e:22033)
- [13] S.L. Woronowicz, Compact quantum groups, in *Symétries Quantiques – Les Houches 1995*, North-Holland (1998), 845–884. MR1616348 (99m:46164)

LABORATOIRE EMILE PICARD, UNIVERSITÉ PAUL SABATIER, 118 ROUTE DE NARBONNE, 31062 TOULOUSE, FRANCE

E-mail address: banica@picard.ups-tlse.fr

INSTITUTUL DE MATEMATICĂ AL ACADEMIEI ROMÂNE, P.O. BOX 1-764, RO-014700 BUCUREȘTI, ROMANIA

E-mail address: moroianu@alum.mit.edu