

LOW FOR RANDOM REALS AND POSITIVE-MEASURE DOMINATION

BJØRN KJOS-HANSEN

(Communicated by Julia Knight)

ABSTRACT. The low for random reals are characterized topologically, as well as in terms of domination of Turing functionals on a set of positive measure.

1. INTRODUCTION

A function $f : \omega \rightarrow \omega$ is *uniformly almost everywhere (a.e.) dominating* if for measure-one many X , and all g computable from X , f dominates g . Such functions were first studied by Kurtz [8] who showed that uniformly a.e. dominating functions exist and that in fact $0'$, the Turing degree of the halting problem, computes one of them. If we replace measure by category, there are no such functions, as is not hard to see. A few decades later Dobrinen and Simpson [5] made use of a.e. domination in Reverse Mathematics. They made a couple of fundamental conjectures that were promptly refuted in [2] and [4]. In this article we strengthen the results of [2] to provide a characterization of a related concept, positive-measure domination, in terms of lowness for randomness. Conversely, we characterize low for random reals in terms of such domination. The following characterizations are already known. (We assume the reader is familiar with the definition of Martin-Löf random reals and of prefix-free Kolmogorov complexity K .)

Theorem 1.1 (Nies, Hirschfeldt, Stephan, Terwijn [6], [10], [11]). *The following are equivalent for $A \in 2^\omega$:*

- *A is low for random: each Martin-Löf random real is Martin-Löf random relative to A .*
- *A is K -trivial: $\exists c \forall n K(A \upharpoonright n) \leq K(\emptyset \upharpoonright n) + c$.*
- *A is low for K : $\exists c \forall n K(n) \leq K^A(n) + c$.*
- *$\exists Z \geq_T A$, Z is ML-random relative to A .*
- *$A \leq_T 0'$ and Ω is ML-random relative to A .*

The low for random reals induce a Σ_3^0 nonprincipal ideal in the Turing degrees bounded above by a low₂ Δ_2^0 degree [10], and have already found application to long-standing open problems in computability theory. Our characterizations in

Received by the editors November 29, 2005 and, in revised form, January 21, 2006.

2000 *Mathematics Subject Classification*. Primary 03D28, 68Q30.

The author thanks the Institute for Mathematical Sciences of the National University of Singapore for support during the preparation of this manuscript at the *Computational Prospects of Infinity* conference in Summer 2005. The author also thanks Denis R. Hirschfeldt for proving upon request a lemma used in an earlier proof of the case $B \leq_T 0'$ of Theorem 3.2.

©2007 American Mathematical Society
Reverts to public domain 28 years from publication

this paper are distinguished by not being couched in the language of randomness and Kolmogorov complexity. They do however refer to measure; it remains open whether a characterization purely in terms of domination or traces can be given such as that found for low for Schnorr random reals [1], [13].

The first main result of Section 2 is Theorem 2.10, which is a characterization of the low for random reals in terms of containment of effectively closed sets of positive measure. Building on this result, Theorem 2.12 is a characterization of low for random reals in terms of positive-measure domination. Section 3 contains, first, a characterization of the Turing degrees relative to which $0'$ is low for random, in terms of positive-measure domination. Finally, with a view toward future research, we include a proof that there is a Turing functional that is universal for this kind of domination.

2. LOW FOR RANDOM REALS

To obtain our topological characterization, we will pass first from a certain universal Martin-Löf test (given in terms of K) to an arbitrary Martin-Löf test, and then to an arbitrary open set of measure < 1 .

Theorem 2.1 (Kraft-Chaitin Theorem [3]). *Suppose $\langle n_k, \sigma_k \rangle$, $k \in \omega$, is a recursive sequence, with $\sum_k 2^{-n_k} \leq 1$. Then there exists a prefix-free machine M and a collection of strings τ_k with $|\tau_k| = n_k$ and $M(\tau_k) = \sigma_k$.*

Definition 2.2 (Chaitin). Let $A \in 2^\omega$. An *information content measure relative to A* is a partial function $\hat{K} : 2^{<\omega} \rightarrow \omega$ such that

$$\sum_{\sigma \in 2^{<\omega}} 2^{-\hat{K}(\sigma)} \leq 1$$

and $\{\langle \sigma, k \rangle : \hat{K}(\sigma) \leq k\}$ is r.e. in A .

Proof. Let σ_k be the k th string to enter V_n^A , and let $n_k = |\sigma_k| - n$. Note that $\sum_k 2^{-n_k} = 2^n \sum_k 2^{-|\sigma_k|} \leq 1$, so we can let M be as in Theorem 2.1. In other words, if σ enters V_n^A , then produce an M -description of σ of length $\leq |\sigma| - n$. Let $\hat{K}(\sigma)$ be the length of the shortest description of σ so produced. \square

Lemma 2.3 (Chaitin). *If \hat{K} is an information content measure relative to a real A , then for all n , $K^A(n) \leq \hat{K}(n) + \mathcal{O}(1)$.*

Definition 2.4. For any real X , let $S^A = \{S_n^A\}_{n \in \omega}$, where $S_n^A = \{X : \exists m K^A(X \upharpoonright m) \leq m - n\}$.

Lemma 2.5. *If V^A is a Martin-Löf test relative to A , then for each n there exists p such that $V_p^A \subseteq S_n^A$.*

Proof. For each m , write $V_{2m}^A = \bigcup \{[\sigma_{m,k}] : k \in \omega\}$ where the function f given by $f(m, k) = \sigma_{m,k}$ is computable and the sets $\{\sigma_{m,k} : k \geq 1\}$ are prefix-free. Define numbers $n_{m,k} = |\sigma_{m,k}| - m + 1$ for $m, k \in \omega$. We have

$$\sum_{m,k} 2^{-|n_{m,k}|} = \sum_m 2^{m-1} \sum_k 2^{-|\sigma_{m,k}|} = \sum_m 2^{m,1} \mu V_{2m} \leq \sum_m 2^{m-1} 2^{-2m} = 1.$$

Hence by Theorem 2.1, we have a partial A -recursive prefix-free machine M and strings $\tau_{m,k}$ with $|\tau_{m,k}| = n_{m,k}$ and $M(\tau_{m,k}) = \sigma_{m,k}$. Thus K_M , complexity based on the machine M , satisfies $K_M(\sigma_{m,k}) \leq n_{m,k}$, and so by Lemma 2.3, there is a

constant c such that $K^A(\sigma_{m,k}) \leq n_{m,k} + c = |\sigma_{m,k}| - m + 1 + c$. This means that $V_{2m}^A \subseteq S_{m-c-1}^A$ for each m .

Thus, given n , let $m = n + c + 1$ and $p = 2m$. Then $V_p^A = V_{2m}^A \subseteq S_{m-c-1}^A = S_n^A$, as desired. \square

Schnorr [12] showed that S^A is a universal Martin-Löf test relative to A .

Proof. In Lemma 2.5 we can conclude that $\bigcap_n V_n^A \subseteq \bigcap_n S_n^A$. Moreover $\mu S_n^A = \sum \{2^{-|\sigma|} : K(\sigma) \leq |\sigma| - n\} < 2^{-n}$. \square

Definition 2.6. Let $n \geq 1$. Let Σ_n^μ denote the collection of all Σ_n^0 classes of measure < 1 . The complement of a Σ_n^μ class is a Π_n^μ class. The complement of U is denoted \bar{U} . The clopen subset of 2^ω generated by $\sigma \in 2^{<\omega}$ is denoted $[\sigma]$, and concatenation of strings is denoted by juxtaposition.

If U, V are open subsets of 2^ω given by $U = \bigcup\{[\sigma] : \sigma \in \hat{U}\}$ and $V = \bigcup\{[\sigma] : \sigma \in \hat{V}\}$, where U and V are prefix-free sets of strings, then we define

$$UV = \bigcup\{[\sigma\tau] : \sigma \in \hat{U}, \tau \in \hat{V}\}.$$

This product depends on \hat{U} and \hat{V} , not just on U and V . So when considering a Σ_1^0 class U , we implicitly fix a suitable recursively enumerable set \hat{U} for U .

We define $U^n = U^{n-1}U$ where $U^1 = U$. We can also think of this exponentiation as acting on a closed set Q , defining Q^n via the equation $\overline{Q^n} = \overline{Q^n}$. It will be clear whether we are considering a set as open or closed.

Lemma 2.7 (Kučera [7]). *For each $\Pi_1^\mu(A)$ class Q there is a computable function f such that $\{\overline{Q^{f(n)}}\}_{n \in \omega}$, is a Martin-Löf test relative to A .*

Proof. Let $q > 0$ be a rational number such that $\mu Q \geq q$. Let $P = \overline{Q}$. Then $\mu P^n = (\mu P)^n \leq (1 - q)^n$. Let f be a computable function such that for all $k \in \omega$, $\mu P^{f(k)} \leq 2^{-k}$. Let $V_k^A = P^{f(k)}$. Then V^A is a Martin-Löf test relative to A . \square

Lemma 2.8. *If P is an open set such that P^n is contained in a Σ_1^μ class for some $n \geq 2$, then P itself is contained in a Σ_1^μ class.*

Proof. We write $U|\sigma = \bigcup\{[\tau] : [\sigma\tau] \subseteq U\}$. Note that if P is open, then so is P^2 . Hence by iteration, it suffices to consider the case $n = 2$. So suppose $(\exists U) P^2 \subseteq U \in \Sigma_1^\mu$. Case 1: $\exists \sigma, \mu(U|\sigma) < 1, \sigma \in \hat{P}$. Then $P^2 \cap [\sigma] = [\sigma]P$, the product of $[\sigma]$ and P . Then $P = ([\sigma]P)|\sigma = (P^2 \cap [\sigma])|\sigma = P^2|\sigma \subseteq U|\sigma \in \Sigma_1^\mu$. Case 2: Otherwise; so $\hat{P} \subseteq \{\sigma : \mu(U|\sigma) = 1\}$. Fix $\epsilon > 0$ such that $\mu U < 1 - \epsilon$, and let $V = \bigcup\{[\sigma] : \mu(U|\sigma) \geq 1 - \epsilon\}$. Note that V is Σ_1^0 , contains P , and $\mu V < 1$ because $(1 - \epsilon)\mu V \leq \mu U < 1 - \epsilon$. \square

As usual, an A -random is a real that is Martin-Löf random relative to A . If $A, B \in 2^\omega$, then A is a tail of B if there exists n such that $A(k) = B(n + k)$ for all $k \in \omega$.

Lemma 2.9 (Kučera [7]). *For each $A \in 2^\omega$, each $\Pi_1^\mu(A)$ class contains a tail of each A -random real.*

Proof. Let Q be a $\Pi_1^\mu(A)$ class and suppose X is A -random. Then by Lemma 2.7, there is an m such that $X \in Q^m$. If $m = 2$, then clearly, as Q is closed, some tail of X is an element of Q . If $m > 2$, the result follows by iteration since each Q^m is closed. \square

Theorem 2.10. *Let $A \in 2^\omega$. The following are equivalent:*

- (1) *Each 1-random real is A -random (A is low for random [10]).*
- (2) *For each Π_1^μ class Q consisting entirely of 1-random reals, there exist σ, n such that $Q \cap [\sigma] \neq \emptyset$ but $Q \cap S_n^A \cap [\sigma] = \emptyset$.*
- (3) *For some n , $\overline{S_n^A}$ has a Π_1^μ subclass.*
- (4) *For each A -Martin-Löf test V_n^A , there exists an n such that $\overline{V_n^A}$ has a Π_1^μ subclass.*
- (5) *For each $\Pi_1^\mu(A)$ class Q there exists an n such that Q^n has a Π_1^μ subclass.*
- (6) *Each $\Pi_1^\mu(A)$ class has a Π_1^μ subclass.*
- (7) *Some $\Pi_1^\mu(A)$ class consisting entirely of A -random reals has a Π_1^μ subclass.*
- (8) *The class of A -random reals has a Π_1^μ subclass.*

Proof. (1) \Rightarrow (2): For this implication we use an argument of Nies and Stephan [9]. Suppose A is low for random but (2) fails. So there is a Π_1^μ class Q consisting entirely of 1-random reals, such that for all σ, n , if $Q \cap [\sigma] \neq \emptyset$, then $Q \cap S_n^A \cap [\sigma] \neq \emptyset$. Let $\sigma_0 = \lambda$, and $\sigma_{n+1} \succeq \sigma_n$, with $[\sigma_{n+1}] \subseteq S_n^A$ but $[\sigma_{n+1}] \cap Q \neq \emptyset$. Then $Y = \bigcup_{n \in \omega} \sigma_n$ is not A -random, but is 1-random, since $Y \in Q$.

(2) \Rightarrow (3): Let Q be as in (2), and let n, σ be as guaranteed by (2) for Q . Then $Q \cap [\sigma]$ is the desired subclass. It has positive measure because no 1-random belongs to a Π_1^0 class of measure zero.

(3) \Rightarrow (4): Lemma 2.5.

(4) \Rightarrow (5): Let Q be a $\Pi_1^\mu(A)$ class. By Lemma 2.7, $V_k^A = \overline{Q^{f(k)}}$ is a Martin-Löf test relative to A for some computable f . By (4), $Q^{f(m)} = \overline{V_m^A} \supseteq F$ for some $F \in \Pi_1^\mu$ and m ; let $n = f(m)$.

(5) \Rightarrow (6): Lemma 2.8.

(6) \Rightarrow (7): If U^A is a universal Martin-Löf test for A -randomness, then we can let $Q = \overline{U_1}$.

(7) \Rightarrow (8): Since any class consisting entirely of A -randoms is contained in the class of all A -randoms.

(8) \Rightarrow (1): Suppose X is 1-random; we need to show X is A -random. Let F be a Π_1^μ subclass of the class of A -randoms. By Lemma 2.9, some tail of X is an element of F . Hence a tail of X is A -random, and thus X itself is A -random. \square

To characterize the low for random reals in terms of domination we first introduce some notation. We write

$$\text{Tot}(\Phi) = \{X : \Phi^X \text{ is total}\} \quad \text{and} \quad \varphi^X(n) = (\mu s)(\forall m < n)(\Phi_s^X(m) \downarrow \leq s).$$

Note that $\text{Tot}(\Phi)$ is a Π_2^0 class for each Φ , and $\text{Tot}(\Phi) = \text{Tot}(\varphi)$. The function φ is the running time of Φ , explicitly satisfying $\Phi^X(n) \leq \varphi^X(n)$ for all n . Let Φ be a Turing functional and $B \in 2^\omega$. If there exists $f \leq_T B$ such that for positive-measure many X , Φ^X is dominated by f , then we write $\Phi < B$. By σ -additivity this is equivalent to the statement that there exists $f \leq_T B$ such that for positive-measure many X , Φ^X is *majorized* by f . We also write $\Phi < B$ in the case that $\text{Tot}(\Phi)$ has measure zero.

Lemma 2.11 (implicit in [5]). *Let $B \in 2^\omega$ and let Φ be a Turing functional. Then $\varphi < B$ iff $\text{Tot}(\Phi)$ has a $\Pi_1^\mu(B)$ subclass.*

Proof. First suppose $\varphi < B$, as witnessed by f . Then $\{X : \forall n \Phi_{f(n)}^X(n) \downarrow\}$ is a $\Pi_1^\mu(B)$ subclass of $\text{Tot}(\Phi)$. Conversely, let F be a $\Pi_1^\mu(B)$ subclass of $\text{Tot}(\Phi)$. By compactness, $\{\varphi^X(n) : X \in F\}$ is finite for each n , and $\{\langle n, m \rangle : \forall X (X \in F \rightarrow \varphi^X(n) < m)\}$ is a $\Sigma_1^0(B)$ class. Hence by $\Sigma_1^0(B)$ uniformization there is a function $f \leq_T B$ such that $\forall n \forall X (X \in F \rightarrow \varphi^X(n) < f(n))$; i.e., f witnesses that $\varphi < B$. \square

Theorem 2.12. *Let $A \in 2^\omega$. The following are equivalent:*

- (1) A is low for random.
- (2) Each $\Pi_1^\mu(A)$ class has a Π_1^μ subclass.
- (3) (i) $A \leq_T 0'$, and (ii) for each Φ , if $\text{Tot}(\Phi)$ has a $\Pi_1^\mu(A)$ subclass, then $\varphi < 0$.
- (4) (i) $A \leq_T 0'$, and (ii) for each Φ , if $\varphi < A$, then $\varphi < 0$.

Proof. (1) \Leftrightarrow (2) was shown in Theorem 2.10.

(2) \Rightarrow (3): Nies [10] shows that if A is low for random, then $A \leq_T 0'$. Suppose $\text{Tot}(\Phi)$ has a $\Pi_1^\mu(A)$ subclass Q . By (2), $\text{Tot}(\Phi)$ has a Π_1^μ subclass F . By Lemma 2.11, we are done.

(3) \Rightarrow (2): Suppose (3) holds and suppose Q is a $\Pi_1^\mu(A)$ class. Pick Ψ such that $Q = \{X : \Psi^{X \oplus A}(0) \uparrow\}$. Since $A \leq_T 0'$, $A = \lim_s A_s$, the limit of a computable approximation. Let $\Phi^X(s) = \mu t > s(\Psi_t^{X \oplus A_t}(0) \uparrow)$. Then $Q = \text{Tot}(\Phi)$. Applying (3) to this Φ , we have $\varphi < 0$, and so by Lemma 2.11 we are done.

(3) \Leftrightarrow (4) is immediate from Lemma 2.11. \square

3. POSITIVE-MEASURE DOMINATION

In [5] it was asked whether the Turing degrees A of uniformly a.e. dominating functions are characterized by either of the inequalities $A \geq 0'$ and $A' \geq_T 0''$. The case $A \geq 0'$ was refuted by a direct construction in [4]. The case $A' \geq_T 0''$ was refuted in [2] using precursors to the results presented here. Namely, the dual of property 4(ii) above is $\forall \varphi (\varphi < A)$ or equivalently $\forall \varphi (\varphi < 0' \rightarrow \varphi < A)$. Relativizing our proofs gives that this is equivalent to: $0'$ is low for random relative to A . If we restrict ourselves to $A \leq_T 0'$, then by [10] this implies $A' \geq_{tt} 0''$, which is strictly stronger than $A' \geq_T 0''$. We do not know whether the assumption $A \leq_T 0'$ is necessary for either of the conclusions $A' \geq_{tt} 0''$, $A' \geq_T 0''$.

We say that A is *positive-measure dominating* if for each Φ , $\Phi < A$. If each B -random real is A -random, then we write $A \leq_{LR} B$ (A is low for random relative to B) following [10]. We write Φ^A for the functional $X \mapsto \Phi^{A \oplus X}$.

Lemma 3.1. *Let $A \in 2^\omega$ and $\mathcal{C} \subseteq 2^\omega$. Then \mathcal{C} is a $\Pi_2^0(A)$ class iff \mathcal{C} is $\text{Tot}(\Phi^A)$ for some Turing functional Φ .*

Proof. Suppose \mathcal{C} is a $\Pi_2^0(A)$ class, i.e. $\mathcal{C} = \{X : \forall y \exists s R(y, s, A, X)\}$, where R is a formula in the language of second-order arithmetic all of whose quantifiers are first-order and bounded. Then we can let $\Phi^{A \oplus X}(y) = \mu s (R(y, s, A, X))$. Conversely, $\text{Tot}(\Phi^A) = \{X : \forall y \exists s (\Phi_s^{A \oplus X}(y) \downarrow)\}$. \square

Theorem 3.2. *Let $B \in 2^\omega$. Then $0'$ is low for random relative to B iff B is positive-measure dominating.*

Proof. This is the special case $A = 0$ of the fact that for each $A, B \in 2^\omega$, the following are equivalent:

- (1) $A' \leq_{LR} A \oplus B$.
- (2) Each $\Pi_1^\mu(A')$ class has a $\Pi_1^\mu(A \oplus B)$ subclass.
- (3) Each $\Pi_2^\mu(A)$ class has a $\Pi_1^\mu(A \oplus B)$ subclass.
- (4) $\forall \Phi$, if $\text{Tot}(\Phi^A)$ has positive measure, then it has a $\Pi_1^\mu(A \oplus B)$ subclass.
- (5) $\forall \Phi(\varphi^A < A \oplus B)$.

The equivalences are proved as follows. (1) \Leftrightarrow (2): Relativization of Theorem 2.10 gives: $A \leq_{LR} B$ iff each $\Pi_1^\mu(A)$ class has a $\Pi_1^\mu(B)$ subclass. (3) \Leftrightarrow (4): Lemma 3.1. (4) \Leftrightarrow (5): Relativization of Lemma 2.11. (2) \Leftrightarrow (3): Let $A \in 2^\omega$. A' is uniformly a.e. dominating relative to A , hence A' is positive-measure dominating relative to A . Hence by putting $B = A'$ in (3) \Leftrightarrow (5), each $\Pi_2^\mu(A)$ class has a $\Pi_1^\mu(A')$ subclass. \square

Universal functionals. Suppose Φ_i , $i \in \omega$, are all the Turing functionals. As observed in [4], the functional Ψ given by $\Psi^{0^i 1^X} = \Phi_i^X$ is *universal* for uniform a.e. domination, in the sense that any function that dominates Ψ on almost every X is a uniformly a.e. dominating function. As $\Psi < 0$, Ψ is not universal for positive-measure domination; however, the following functional is universal.

Fix $c \in \omega$. Let $\Xi_c^X(s)$ be the least stage $t > s$ at which X looks like it is 2-random, with constant c . That is,

$$\Xi_c^X(s) = (\mu t > s)(\forall n \leq t K_t^{0^i}(X \upharpoonright n) \geq n - c).$$

Here 0_t^i is the approximation to 0^i at stage t , and K_t^A the approximation to K^A for any $A \in 2^\omega$. Considering S_c^X (Definition 2.4), it is clear that Ξ_c is total for positive-measure many X , all of which are 2-randoms. The running time ξ_c of Ξ_c is universal for positive-measure domination in the following sense.

Theorem 3.3. *The class $\{A : A \text{ is positive-measure dominating}\}$ is Σ_3^0 . In fact, for each $A \in 2^\omega$ and $c \in \omega$, A is positive-measure dominating iff $\xi_c < A$.*

Proof. The complement of $\text{Tot}(\Xi_c)$ is $\{X : \exists n K^{0^i}(X \upharpoonright n) < n - c\}$, which is open. Hence $\text{Tot}(\Xi_c)$ is closed and is in fact a $\Pi_1^\mu(0^i)$ class. Suppose $\xi_c < A$. By Lemma 2.11, $\text{Tot}(\Xi_c)$ has a $\Pi_1^\mu(A)$ subclass. Thus: Some $\Pi_1^\mu(0^i)$ class consisting entirely of 0^i -randoms has a $\Pi_1^\mu(A)$ subclass. By Theorem 2.10(7) relativized, $0^i \leq_{LR} A$, and so by Theorem 3.2, A is positive-measure dominating. \square

REFERENCES

1. A. Nies, B. Kjos-Hanssen and F. Stephan, *Lowness for the class of Schnorr random reals*, SIAM J. Computing **35** (2006), no. 3, 647–657. MR2201451
2. S. Binns, B. Kjos-Hanssen, M. Lerman, and D.R. Solomon, *On a question of Dobrinen and Simpson concerning almost everywhere domination*, J. Symbolic Logic **71** (2006), no. 1, 119–136.
3. G.J. Chaitin, *A theory of program size formally identical to information theory*, J. Assoc. Comput. Mach. **22** (1975), 329–340. MR0411829 (53:15557)
4. P. Cholak, N. Greenberg, and J.S. Miller, *Uniform almost everywhere domination*, J. Symbolic Logic **71** (2006), no. 3, 1057–1072. MR2251556
5. N.L. Dobrinen and S.G. Simpson, *Almost everywhere domination*, J. Symbolic Logic **69** (2004), 914–922. MR2078930 (2005d:03079)
6. D.R. Hirschfeldt, A. Nies, and F. Stephan, *Using random sets as oracles*, to appear.

7. Antonín Kučera, *Measure, Π_1^0 -classes and complete extensions of PA*, Recursion theory week (Oberwolfach, 1984), Lecture Notes in Math., vol. 1141, Springer, Berlin, 1985, pp. 245–259. MR0820784 (87e:03102)
8. S.A. Kurtz, *Randomness and genericity in the degrees of unsolvability*, Ph.D. thesis, University of Illinois at Urbana-Champaign, 1981, pp. VII+131 pages.
9. A. Nies, *Low for random reals: the story*, unpublished.
10. ———, *Lowness properties and randomness*, Adv. Math. **197** (2005), 274–305. MR2166184
11. A. Nies, F. Stephan, and S.A. Terwijn, *Randomness, relativization and Turing degrees*, J. Symbolic Logic **70** (2005), no. 2, 515–535. MR2140044
12. C.P. Schnorr, *A unified approach to the definition of a random sequence*, Mathematical Systems Theory **5** (1971), 246–258. MR0354328 (50:6808)
13. S.A. Terwijn and D. Zambella, *Computational randomness and lowness*, J. Symbolic Logic **66** (2001), 1199–1205. MR1856736 (2002j:03044)

DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, ITHACA, NEW YORK 14853
E-mail address: bjorn@math.cornell.edu