

AN ISOPERIMETRIC INEQUALITY  
FOR THE SECOND EIGENVALUE OF THE LAPLACIAN  
WITH ROBIN BOUNDARY CONDITIONS

JAMES KENNEDY

(Communicated by Matthew J. Gursky)

ABSTRACT. We prove that the second eigenvalue of the Laplacian with Robin boundary conditions is minimized among all bounded Lipschitz domains of fixed volume by the domain consisting of the disjoint union of two balls of equal volume.

1. INTRODUCTION

Let  $\Omega \subset \mathbb{R}^N$  be a bounded Lipschitz domain (not necessarily connected) and consider the eigenvalue problem for the Laplacian with Robin boundary condition

$$(1.1) \quad \begin{aligned} -\Delta u &= \lambda u && \text{in } \Omega, \\ \frac{\partial u}{\partial \nu} + \beta u &= 0 && \text{on } \partial\Omega, \end{aligned}$$

where  $\beta > 0$  is a constant and  $\nu$  is the outer unit normal to  $\Omega$ . This problem is sometimes referred to as the third boundary value problem, or as the problem of the elastically supported membrane. It is well known that, as in the case of Dirichlet boundary conditions, the associated operator on  $L^2(\Omega)$  has compact resolvent, with the eigenvalues forming a sequence  $0 < \lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \dots \rightarrow \infty$ .

It has been shown in [1, 5] that the first eigenvalue  $\lambda_1 = \lambda_1(\Omega)$  satisfies the isoperimetric, or Faber-Krahn, inequality  $\lambda_1(\Omega) \geq \lambda_1(B)$ , where  $B$  is a ball having the same volume as  $\Omega$ . Our current goal is to prove a similar inequality for  $\lambda_2(\Omega)$ .

**Theorem 1.1.** *The second eigenvalue  $\lambda_2(\Omega)$  of (1.1) on a bounded Lipschitz domain  $\Omega \subset \mathbb{R}^N$  satisfies  $\lambda_2(\Omega) \geq \lambda_2(D)$ , where  $D$  is a domain of the same volume as  $\Omega$  consisting of the disjoint union of two balls of equal volume.*

We defer the proof of Theorem 1.1 to Section 3 and first discuss some background issues and consequences.

---

Received by the editors January 30, 2008.

2000 *Mathematics Subject Classification.* Primary 35P15, 35J25.

*Key words and phrases.* Isoperimetric inequality, Laplacian, Robin boundary conditions, elastically supported membrane.

©2008 American Mathematical Society  
Reverts to public domain 28 years from publication

## 2. OBSERVATIONS AND REMARKS

We first wish to consider the Laplacian with Dirichlet boundary conditions. Not only is the minimizing domain  $D$  the same for both Dirichlet and Robin boundary conditions, but our proof uses ideas from the Dirichlet case. For this reason we will give a brief sketch of the proof of the latter here. A more complete proof, together with further references, can be found in [10, Sec. 4].

Let  $\varphi$  denote an eigenfunction of the second Dirichlet eigenvalue, which we will call  $\mu_2(\Omega)$ . The idea is to consider the nodal domains  $\Omega^+ := \{x \in \Omega : \varphi(x) > 0\}$  and  $\Omega^- := \{x \in \Omega : \varphi(x) < 0\}$ . Then  $\varphi$  is an eigenfunction of the Dirichlet Laplacian that does not change sign in  $\Omega^+$ , so that  $\mu_2(\Omega) = \mu_1(\Omega^+)$  ( $\mu_1$  being the first Dirichlet eigenvalue). If we denote by  $B^+$  a ball of the same volume as  $\Omega^+$ , then the usual Faber-Krahn inequality applied to  $\Omega^+$  gives  $\mu_2(\Omega) \geq \mu_1(B^+)$ . Similarly, if  $B^-$  is a ball of the same volume as  $\Omega^-$ , then  $\mu_2(\Omega) \geq \mu_1(B^-)$ .

Hence  $\mu_2(\Omega) \geq \max\{\mu_1(B^+), \mu_1(B^-)\}$ . The latter is minimized if  $B^+ = B^- =: B$  has half the volume of  $\Omega$ . But  $D$  (defined in Theorem 1.1) can be written as the disjoint union of two copies of  $B$ , so that  $\mu_2(D) = \mu_1(D) = \mu_1(B) \leq \mu_2(\Omega)$ .

We would like to use a similar idea in the Robin case. Denoting an eigenfunction of  $\lambda_2(\Omega)$  by  $\psi$ , we wish to describe  $\lambda_2(\Omega)$  as the first eigenvalue of a problem on  $\Omega^+$  (and  $\Omega^-$ ) with mixed Robin-Dirichlet boundary conditions  $\frac{\partial\psi}{\partial\nu} + \beta\psi = 0$  on  $\partial\Omega^+ \cap \partial\Omega$  and  $\psi = 0$  on  $\partial\Omega^+ \cap \Omega$ . This should be greater than the first eigenvalue  $\lambda_1(\Omega^+)$  of the pure Robin problem on  $\Omega^+$ . By the Faber-Krahn inequality for Robin problems [1, 5],  $\lambda_1(\Omega^+) \geq \lambda_1(B^+)$ , and we proceed as before.

However, there is a major complication. We cannot directly apply the inequality from [1, 5] to  $\Omega^+$ ,  $\Omega^-$  since that result is only valid for Lipschitz domains and in general  $\Omega^+$ ,  $\Omega^-$  may not be that smooth. The problem is twofold.

First, we have no control over the behaviour of  $\partial\Omega^+$ ,  $\partial\Omega^-$  near where the nodal surface  $\{x \in \Omega : \psi(x) = 0\}$  meets  $\partial\Omega$ . At such points  $x$ , supposing the boundary condition holds pointwise we have  $\frac{\partial\psi}{\partial\nu}(x) = 0$ . This is in general consistent with the possibility that  $\nabla\psi(x) = 0$ .

Second, even though the eigenfunction  $\psi$  will be  $C^\infty$  in  $\Omega$ , this is not enough to guarantee that the nodal surface is a smooth manifold in the interior. Sard's Lemma (see [11, Ch. 3, Theorem 1.3]) does not suffice, since 0 may be in the null set of nonregular values of  $\psi$ .

We overcome these problems by constructing suitable approximations to the nodal domains. Note that we do not use Sard's Lemma or even the Courant-Hilbert Theorem [3, Ch. VI, Sec. 6].

*Remark 2.1.* (i) We emphasize that we do not require our domains to be connected. Although connectedness of  $\Omega$  was implicitly assumed in the proof sketched above, and is explicitly assumed in Section 3, there is a standard and easy way to remove this assumption. Suppose that Theorem 1.1 holds for connected domains and that  $\Omega$  is not connected. Then either  $\lambda_2(\Omega) = \lambda_2(\Omega_0)$  for some connected component  $\Omega_0$  of  $\Omega$  or there exist disjoint connected components  $\Omega_1, \Omega_2$  of  $\Omega$  such that  $\lambda_1(\Omega) = \lambda_1(\Omega_1)$ ,  $\lambda_2(\Omega) = \lambda_1(\Omega_2)$ . In the former case Theorem 1.1 applied to  $\Omega_0$ , together with the monotonicity of  $\lambda_1(D) = \lambda_2(D)$  with respect to the volume of  $D$ , yields the result. In the latter case if we replace  $\Omega^+$  by  $\Omega_1$  and  $\Omega^-$  by  $\Omega_2$ , we may repeat verbatim the argument used when  $\Omega$  is connected. This argument works equally well for Dirichlet and Robin boundary conditions.

(ii) We leave as an open problem the sharpness of the inequality. That is, is the domain  $D$  the *unique* minimizer of  $\lambda_2$ ? It is for the Dirichlet Laplacian, at least up to sets of capacity zero and rigid transformations such as translations and rotations. Moreover, the inequality for the first eigenvalue of the Robin problem (1.1) is sharp, at least for  $C^2$  domains (see [7, Theorem 1.1]). Our method is unlikely to yield a sharpness result, as it uses approximation arguments.

(iii) We also note as an open problem the case when  $\beta < 0$ . In this case the eigenvalues still form a sequence  $\lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \dots \rightarrow \infty$ , but now  $\lambda_1(\Omega) < 0$  for every  $\Omega$  (see for example [12, Sec. 2]). Moreover, it is easy to see that we cannot minimize  $\lambda_m(\Omega)$  with respect to  $\Omega$  for any  $m$ . Indeed, let  $\Omega_n$  be a sequence of smooth domains of fixed volume such that the surface measure  $\sigma(\partial\Omega_n) \rightarrow \infty$ . Using any constant as a test function in the Rayleigh quotient for  $\lambda_1(\Omega_n)$  (see [12, Eq. 2.1]), we see that  $\lambda_1(\Omega_n) \leq \beta\sigma(\partial\Omega_n)/|\Omega_n| \rightarrow -\infty$  as  $n \rightarrow \infty$ . Letting  $\tilde{\Omega}_n$  be the disjoint union of  $m$  copies of  $\Omega_n$ , we have  $\lambda_m(\tilde{\Omega}_n) = \lambda_1(\Omega_n) \rightarrow -\infty$  as  $n \rightarrow \infty$ . Instead we seek a maximizer for  $\lambda_m$ . It is conjectured that for  $\lambda_1(\Omega)$  this is the ball [13]. For  $\lambda_2$ ,  $D$  cannot be the maximizer for all values of  $\beta < 0$  since  $\lambda_2(D) < 0$  for all  $\beta$ , but there exist  $\Omega$  and  $\beta$  for which  $\lambda_2(\Omega) > 0$ . (For example if  $\Omega = B_1$ , the ball of radius 1, then by [12, Sec. 2.4]  $B_1$  has  $1 + \lfloor |\beta| \rfloor$  negative eigenvalues. So  $\lambda_2(B_1) > 0$  if  $\beta > -1$ .) In fact we suggest  $B$  is the best candidate for the maximizer, since  $\lambda_2(B)$  is maximal for  $\beta = 0$  (Neumann boundary conditions) and  $\lambda_2$  should depend continuously on  $\beta \in \mathbb{R}$  for fixed  $\Omega$ .

(iv) Finally, we might ask if there is a minimizer of  $\lambda_2$  (for  $\beta > 0$ ) among all *connected* domains. In the Dirichlet case there is none: we can find a sequence of connected domains  $\Omega_n$  with  $\mu_2(\Omega_n) \rightarrow \mu_2(D)$ , with  $D$  being the unique overall minimizer of  $\mu_2$  (see [10, Sec. 4]). A similar construction works in the Robin case (see Example 2.2), but we cannot yet finish the argument as we do not yet know if  $D$  is the unique minimizer of  $\lambda_2$ .

**Example 2.2.** We construct a sequence of connected domains  $\Omega_n$  of fixed volume such that  $\lambda_2(\Omega_n) \rightarrow \lambda_2(D)$  (see Figure 1). Our domains are almost identical to the “dumbbells” used in [10]. Start with  $D = B_1 \cup B_2$  and join  $B_1$  to  $B_2$  with a cylinder  $C_n$  of total volume  $\frac{1}{n}$ . To keep the volume of  $\Omega_n$  constant, remove part of  $B_1$  and  $B_2$  in a small neighbourhood  $U_n$  near where they meet  $C_n$  (as in Figure 1) in such a way that the resulting boundary is still smooth. It now follows from [4, Corollary 3.7] that  $\lambda_2(\Omega_n) \rightarrow \lambda_2(D)$ , since the  $U_n$  can be chosen in such a way that Assumption 3.2 of [4] is satisfied.




FIGURE 1. The domain  $\Omega_n$

3. PROOF OF THEOREM 1.1

First we fix our notation. Let  $\Omega \subset \mathbb{R}^N$  be a bounded Lipschitz domain. As noted in Remark 2.1(i) we may assume without loss of generality that  $\Omega$  is connected. Its second eigenvalue  $\lambda_2(\Omega)$  has an eigenfunction  $\psi \in H^1(\Omega) \cap C(\overline{\Omega}) \cap C^\infty(\Omega)$  (interior regularity is standard and continuity up to the boundary comes from combining [6, Corollary 5.5] with [14, Corollary 2.9]). Since  $\Omega$  is connected,  $\psi$  changes sign in  $\Omega$ , so that the open subsets  $\Omega^+ = \{x \in \Omega : \psi(x) > 0\}$  and  $\Omega^- = \{x \in \Omega : \psi(x) < 0\}$  are both nonempty. Set  $\psi^+ := \max\{\psi, 0\}, \psi^- := \max\{-\psi, 0\}$ . Then  $\psi^+, \psi^- \in H^1(\Omega) \cap C(\overline{\Omega})$  and  $\nabla\psi^+ \neq 0$  only on  $\Omega^+$ , with a similar statement for  $\nabla\psi^-$  (see [9, Lemma 7.6]). Henceforth  $\lambda_1(V)$  will denote the first eigenvalue of the Robin problem (1.1) on the domain  $V$ . We will denote  $N$ -dimensional Lebesgue measure by  $|\cdot|$  and  $(N - 1)$ -dimensional surface (Hausdorff) measure by  $\sigma$ .

Let  $B^+, B^-$  be balls having the same volume as  $\Omega^+, \Omega^-$  respectively. As sketched at the beginning of Section 2, to prove Theorem 1.1 it suffices to show  $\lambda_2(\Omega) \geq \max\{\lambda_1(B^+), \lambda_1(B^-)\}$ . Without loss of generality we only consider  $\Omega^+$  and prove  $\lambda_2(\Omega) \geq \lambda_1(B^+)$ .

The key idea is to attach a thin strip near  $\partial\Omega$  to  $\Omega^+$  to avoid any problems when  $\{x \in \Omega : \psi(x) = 0\}$  meets  $\partial\Omega$ . See Figure 2. Fix  $\varepsilon > 0$  and set  $S_\varepsilon := \{x \in \Omega : \text{dist}(x, \partial\Omega) < \delta\}$ , where  $\delta = \delta(\varepsilon) > 0$  is chosen such that  $|S_\varepsilon| < \varepsilon$ . Set  $U_\varepsilon := \Omega^+ \cup S_\varepsilon$ . Then  $\partial\Omega \subset \partial U_\varepsilon$ . Denote the rest of  $\partial U_\varepsilon$  by  $\Gamma_\varepsilon$ . Then  $\Gamma_\varepsilon$  is compactly contained in  $\Omega$ , with  $\text{dist}(\partial\Omega, \Gamma_\varepsilon) \geq \delta$ . Moreover,  $|U_\varepsilon \setminus \Omega^+| \leq |S_\varepsilon| < \varepsilon$ . Note however that  $\Gamma_\varepsilon$  may not be Lipschitz.




FIGURE 2. The approximating domain  $U_\varepsilon = \Omega^+ \cup S_\varepsilon$

We consider the mixed problem on  $U_\varepsilon$ :

$$\begin{aligned}
 (3.1) \quad & -\Delta u = \lambda u && \text{in } U_\varepsilon, \\
 & \frac{\partial u}{\partial \nu} + \beta u = 0 && \text{on } \partial\Omega, \\
 & u = 0 && \text{on } \Gamma_\varepsilon.
 \end{aligned}$$

Denote by  $H_{U_\varepsilon}$  the space of weak solutions to (3.1). This is given by the closure in  $H^1(U_\varepsilon)$  (equivalently, in  $H^1(\Omega)$ ) of  $C_c^\infty(U_\varepsilon \cup \partial\Omega)$ , the space of all  $C^\infty(\overline{\Omega})$  functions with support compactly contained in  $U_\varepsilon \cup \partial\Omega$ . (Any element of  $H_{U_\varepsilon}$  may be considered an element of  $H^1(\Omega)$  by extending it by zero outside  $U_\varepsilon$ .) The problem (3.1) has a first eigenvalue, call it  $\Lambda_1(U_\varepsilon)$ , given by the variational formula

$$(3.2) \quad \Lambda_1(U_\varepsilon) = \inf_{\varphi \in H_{U_\varepsilon}} Q(\varphi, U_\varepsilon) := \inf_{\varphi \in H_{U_\varepsilon}} \frac{\int_{U_\varepsilon} |\nabla\varphi|^2 dx + \int_{\partial\Omega} \beta\varphi^2 d\sigma}{\int_{U_\varepsilon} \varphi^2 dx}.$$

In fact for arbitrary open  $V \subset \Omega$  with  $\partial\Omega \subset \partial V$  and  $\text{dist}(\partial\Omega, \partial V \setminus \partial\Omega) > 0$ , we may consider the problem (3.1) on  $V$ , with  $\Lambda_1(V)$  and  $H_V$  given by the obvious analogues of  $\Lambda_1(U_\varepsilon)$  and  $H_{U_\varepsilon}$ . We denote by  $Q(\varphi, V)$  the Rayleigh quotient of (3.2) on  $V$  for a given function  $\varphi \in H_V$ . We have the following important estimate for  $\lambda_2(\Omega)$ .

**Lemma 3.1.** *For any  $\varepsilon > 0$ ,  $\lambda_2(\Omega) \geq \Lambda_1(U_\varepsilon)$ .*

The proof of Lemma 3.1 is based on the following characterization of  $\lambda_2(\Omega)$ , combined with (3.2).

**Lemma 3.2.** *For any  $\varepsilon > 0$ , we have*

$$(3.3) \quad \lambda_2(\Omega) = \frac{\int_{U_\varepsilon} |\nabla\psi^+|^2 dx + \int_{\partial\Omega} \beta(\psi^+)^2 d\sigma}{\int_{U_\varepsilon} (\psi^+)^2 dx}.$$

*Proof.*  $\lambda_2(\Omega)$  satisfies

$$\int_{\Omega} \nabla\psi \cdot \nabla\varphi dx + \int_{\partial\Omega} \beta\psi\varphi d\sigma = \lambda_2(\Omega) \int_{\Omega} \psi\varphi dx$$

for all  $\varphi \in H^1(\Omega)$ . Choosing  $\psi^+ \in H^1(\Omega) \cap C(\overline{\Omega})$  as a test function, we have  $\nabla\psi \cdot \nabla\psi^+ = |\nabla\psi^+|^2$  in  $\Omega$  (see for example [9, Lemma 7.6]) and  $\psi\psi^+ = (\psi^+)^2$  pointwise in  $\overline{\Omega}$ . Since  $\|\psi^+\|_2 \neq 0$ ,

$$\lambda_2(\Omega) = \frac{\int_{\Omega} |\nabla\psi^+|^2 dx + \int_{\partial\Omega} \beta(\psi^+)^2 d\sigma}{\int_{\Omega} (\psi^+)^2 dx}.$$

But the integrands in the volume integrals are nonzero only if  $x \in \Omega^+ \subset U_\varepsilon$ . Hence (3.3) follows.  $\square$

*Proof of Lemma 3.1.* By Lemma 3.2 and (3.2), we only have to prove that  $\psi^+ \in H_{U_\varepsilon}$ . Since  $H_{U_\varepsilon}$  is closed in the  $H^1$  norm, it suffices to prove that there exist  $u_n \in H_{U_\varepsilon}$  such that  $u_n \rightarrow \psi^+$  in  $H^1(U_\varepsilon)$ .

Noting that  $\psi^+ \in H^1(U_\varepsilon) \cap C(\overline{U_\varepsilon})$  and  $\psi^+ = 0$  on  $\Gamma_\varepsilon$ , our claim follows from a trivial modification of the proof of [2, Théorème IX.17] (see also Remarque 20 there). The only difference is that the approximating functions  $u_n$  there will have support compactly contained in  $U_\varepsilon \cup \partial\Omega$  rather than  $U_\varepsilon$  (so that our limit function will lie in  $H_{U_\varepsilon}$  rather than  $H_0^1(U_\varepsilon)$ ).  $\square$

Next, since  $\Gamma_\varepsilon$  may not be smooth, we approximate  $U_\varepsilon$  by a suitable sequence of smooth domains  $U_n \subset U_\varepsilon$  such that  $\Lambda_1(U_n) \rightarrow \Lambda_1(U_\varepsilon)$ .

**Lemma 3.3.** *There exists a sequence of Lipschitz domains  $U_n \subset U_\varepsilon$  such that  $\partial\Omega \subset \partial U_n$  and  $\text{dist}(\partial\Omega, \partial U_n \setminus \partial\Omega) > 0$  for all  $n$ ,  $|U_\varepsilon \setminus U_n| \rightarrow 0$  and  $\Lambda_1(U_n) \rightarrow \Lambda_1(U_\varepsilon)$  as  $n \rightarrow \infty$ .*

*Proof.* The existence of the  $U_n$  converging in volume is a standard result (see for example [8, Ch. V, Theorem 4.20]); note that since  $\partial\Omega$  and  $\Gamma_\varepsilon$  are separated, this is equivalent to having a sequence of the form  $\mathbb{R}^N \setminus (\Omega \setminus U_n)$  converge to  $\mathbb{R}^N \setminus (\Omega \setminus U_\varepsilon)$ . Moreover, the  $U_n$  can be chosen such that  $\{x \in U_\varepsilon : \text{dist}(x, \Gamma_\varepsilon) > \frac{1}{n}\} \subset U_n$ .

Comparing the characterization (3.2) for  $U_n$  and  $U_\varepsilon$ , and since  $H_{U_n} \subset H_{U_\varepsilon}$ , we have  $\Lambda_1(U_n) \geq \Lambda_1(U_\varepsilon)$ . So it remains to prove

$$\limsup_{n \rightarrow \infty} \Lambda_1(U_n) \leq \Lambda_1(U_\varepsilon).$$

Fix  $\delta > 0$  and choose  $\varphi \in H_{U_\varepsilon}$  such that the Rayleigh quotient  $Q(\varphi, U_\varepsilon) < \Lambda_1(U_\varepsilon) + \delta$ . By density, we may assume  $\varphi \in C_c^\infty(U_\varepsilon \cup \partial\Omega)$ .

Then  $\text{supp } \varphi$  (the support of  $\varphi$ ) and  $\Gamma_\varepsilon$  are compact and  $\text{dist}(\text{supp } \varphi, \Gamma_\varepsilon) > 0$ . In particular,  $\text{dist}(\text{supp } \varphi, \Gamma_\varepsilon) > \frac{1}{n}$  for all  $n \in \mathbb{N}$  large enough. By choice of the  $U_n$  it follows that  $\text{supp } \varphi$  is compactly contained in  $U_n \cup \partial\Omega$  for  $n$  large enough.

In particular,  $\varphi \in C_c^\infty(U_n \cup \partial\Omega)$ ; whence  $Q(\varphi, U_\varepsilon) = Q(\varphi, U_n) \geq \Lambda_1(U_n)$  for all  $n$  large enough (where we have used  $\partial\Omega \subset \partial U_n$  to get  $Q(\varphi, U_\varepsilon) = Q(\varphi, U_n)$ ). That is, for any  $\delta > 0$ ,  $\Lambda_1(U_n) < \Lambda_1(U_\varepsilon) + \delta$  for all  $n$  large enough.  $\square$

Choose a sequence of  $U_n$  as in Lemma 3.3 and consider the Robin problem (1.1) on  $U_n$ . Since  $\partial U_n$  is Lipschitz for each  $n$ , we may use the Faber-Krahn inequality for Robin problems and then pass to the limit. So let  $B_n$  be a ball of the same volume as  $U_n$ . Then  $\lambda_1(U_n) \geq \lambda_1(B_n)$  for all  $n$ , by [5, Theorem 1.1]. Moreover, by a direct comparison of variational formulae and since  $H_{U_n} \subset H^1(U_n)$ ,  $\Lambda_1(U_n) \geq \lambda_1(U_n)$ .

Now let  $B_\varepsilon$  be a ball having the same volume as  $U_\varepsilon$ . As  $n \rightarrow \infty$  and  $|U_n| \rightarrow |U_\varepsilon|$ , we have  $|B_n| \rightarrow |B_\varepsilon|$ . Since the first eigenvalue of (1.1) on the ball depends continuously on the size of the ball,  $\lambda_1(B_n) \rightarrow \lambda_1(B_\varepsilon)$ . By Lemma 3.3,  $\Lambda_1(U_n) \rightarrow \Lambda_1(U_\varepsilon)$ . Putting this all together we have

$$\Lambda_1(U_\varepsilon) \leftarrow \Lambda_1(U_n) \geq \lambda_1(U_n) \geq \lambda_1(B_n) \rightarrow \lambda_1(B_\varepsilon),$$

that is,  $\Lambda_1(U_\varepsilon) \geq \lambda_1(B_\varepsilon)$ .

Finally, let  $\varepsilon \rightarrow 0$ . Since  $|U_\varepsilon| \rightarrow |\Omega^+|$ , we have  $|B_\varepsilon| \rightarrow |B^+|$  and so  $\lambda_1(B_\varepsilon) \rightarrow \lambda_1(B^+)$ . Also, since  $\lambda_2(\Omega) \geq \Lambda_1(U_\varepsilon)$  by Lemma 3.1, we conclude  $\lambda_2(\Omega) \geq \lambda_1(B^+)$ . In light of our earlier comments, this completes the proof of Theorem 1.1.

#### ACKNOWLEDGEMENTS

The author would like to thank his thesis supervisor, Daniel Daners, for helpful discussions and advice, as well as the referee for helping to improve this paper.

#### REFERENCES

1. M.-H. Bossel, *Membranes élastiquement liées: Extension du théorème de Rayleigh-Faber-Krahn et de l'inégalité de Cheeger*, C. R. Acad. Sci. Paris Sér. I Math. **302** (1986), 47–50. MR0827106 (87f:35186)
2. H. Brezis, *Analyse fonctionnelle. Théorie et applications*, Masson, Paris, 1983. MR0697382 (85a:46001)
3. R. Courant and D. Hilbert, *Methods of Mathematical Physics*, vol. 1, Interscience Publishers, New York, 1953. MR0065391 (16:426a)
4. E. N. Dancer and D. Daners, *Domain perturbation for elliptic equations subject to Robin boundary conditions*, J. Differential Equations **138** (1997), 86–132. MR1458457 (98e:35017)
5. D. Daners, *A Faber-Krahn inequality for Robin problems in any space dimension*, Math. Ann. **335** (2006), 767–785. MR2232016 (2007b:35242)
6. D. Daners, *Robin boundary value problems on arbitrary domains*, Trans. Amer. Math. Soc. **352** (2000), 4207–4236. MR1650081 (2000m:35048)
7. D. Daners and J. Kennedy, *Uniqueness in the Faber-Krahn inequality for Robin problems*, SIAM J. Math. Anal. **39** (2007/08), 1191–1207. MR2368899
8. D. E. Edmunds and W. D. Evans, *Spectral theory and differential operators*. Oxford Mathematical Monographs. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1987. MR0929030 (89b:47001)
9. D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*, 2nd ed., Grundlehren der Mathematischen Wissenschaften, vol. 224, Springer-Verlag, Berlin, 1983. MR0737190 (86c:35035)
10. A. Henrot, *Minimization problems for eigenvalues of the Laplacian*, J. Evol. Equ. **3** (2003), 443–461. MR2019029 (2005a:49078)

11. M. W. Hirsch, *Differential topology*. Graduate Texts in Mathematics, No. 33, Springer-Verlag, New York-Heidelberg, 1976. MR0448362 (56:6669)
12. A. A. Lacey, J. R. Ockendon and J. Sabina, *Multidimensional reaction diffusion equations with nonlinear boundary conditions*, SIAM J. Appl. Math. **58** (1998), 1622–1647. MR1637882 (99g:35066)
13. Shape Analysis for Eigenvalues. Abstracts from the workshop held April 8–14, 2007. Organised by D. Bucur, G. Buttazzo and A. Henrot, Oberwolfach Reports, vol. 4, no. 2 (2007), 995–1026.
14. M. Warma, *The Robin and Wentzell-Robin Laplacians on Lipschitz domains*, Semigroup Forum **73** (2006), 10–30. MR2277314 (2007g:35032)

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SYDNEY, NEW SOUTH WALES 2006,  
AUSTRALIA

*E-mail address:* J.Kennedy@maths.usyd.edu.au