

A SHORT PROOF OF PITT'S COMPACTNESS THEOREM

SYLVAIN DELPECH

(Communicated by Nigel J. Kalton)

ABSTRACT. We give a short proof of Pitt's theorem that every bounded linear operator from ℓ_p or c_0 into ℓ_q is compact whenever $1 \leq q < p < \infty$.

A bounded linear operator between two Banach spaces X and Y is said to be *compact* if it maps the closed unit ball of X into a relatively compact subset of Y .

Theorem (Pitt; see for example [1], p. 175). *Let $1 \leq q < p \leq +\infty$, and put $X_p = \ell_p$ if $p < +\infty$ and $X_\infty = c_0$. Then every bounded linear operator from X_p into ℓ_q is compact.*

Proof. Let $T : X_p \rightarrow \ell_q$ be a norm-one operator. As $1 < p$, the dual of X_p is separable. Hence every bounded sequence in X_p has a weakly Cauchy subsequence. Thus, for proving the compactness of T , it is enough to show that T is weak-to-norm continuous. So, let us consider a weakly null sequence (h_n) in X_p . We have to show that $\lim_{n \rightarrow \infty} \|T(h_n)\| = 0$. We claim that

- (1) for every $x \in c_0$ and for every weakly null sequence (w_n) in c_0 ,

$$\limsup_{n \rightarrow \infty} \|x + w_n\| = \max(\|x\|, \limsup_{n \rightarrow \infty} \|w_n\|),$$

- (2) for every $x \in \ell_r$, $1 \leq r < \infty$, and for every weakly null sequence (w_n) in ℓ_r ,

$$\limsup_{n \rightarrow \infty} \|x + w_n\|^r = \|x\|^r + \limsup_{n \rightarrow \infty} \|w_n\|^r.$$

Indeed this is obvious when x is finitely supported, because the coordinates of (w_n) along the support of x tend to 0 in norm. The general case is true by the density of finitely supported elements in X_p and since the norm is a Lipschitzian function.

Fix $0 < \varepsilon < 1$. By definition of the norm of T , there exists $x_\varepsilon \in X_p$ such that $\|x_\varepsilon\| = 1$ and $1 - \varepsilon \leq \|T(x_\varepsilon)\| \leq 1$. Moreover, for all $n \in \mathbb{N}$ and for all $t > 0$

$$(0) \quad \|T(x_\varepsilon) + T(th_n)\| \leq \|x_\varepsilon + th_n\|.$$

In the left-hand side of (0), we apply claim (2) in ℓ_q , with $x = T(x_\varepsilon)$ and the weakly null sequence $(T(th_n))$.

First, assume $p < +\infty$. We apply claim (2) to the right-hand side of (0) with $r = p$, $x = x_\varepsilon$ and the weakly null sequence (th_n) to obtain

$$\left[\|T(x_\varepsilon)\|^q + t^q \limsup_{n \rightarrow \infty} \|T(h_n)\|^q \right]^{\frac{1}{q}} \leq \left[\|x_\varepsilon\|^p + t^p \limsup_{n \rightarrow \infty} \|h_n\|^p \right]^{\frac{1}{p}}.$$

Received by the editors February 6, 2008, and, in revised form, April 16, 2008.

2000 *Mathematics Subject Classification*. Primary 46B25.

Key words and phrases. ℓ_p space, c_0 space, compact operator.

©2008 American Mathematical Society
 Reverts to public domain 28 years from publication

Recall that $\|x_\varepsilon\| = 1$, $1 - \varepsilon \leq \|T(x_\varepsilon)\| \leq 1$ and that (h_n) is weakly convergent, thus bounded by some $M > 0$. This gives

$$\limsup_{n \rightarrow \infty} \|T(h_n)\|^q \leq \frac{1}{t^q} \left[(1 + t^p M^p)^{q/p} - (1 - \varepsilon)^q \right].$$

Taking $t = \varepsilon^{\frac{1}{p}}$ here, we get

$$\limsup_{n \rightarrow \infty} \|T(h_n)\|^q \leq \frac{1}{\varepsilon^{q/p}} \left[1 + \frac{q}{p} M^p \varepsilon - (1 - q\varepsilon) + o(\varepsilon) \right].$$

Now, letting $\varepsilon \rightarrow 0$ here, we get that $\limsup_{n \rightarrow \infty} \|T(h_n)\|^q \leq 0$, and therefore the sequence $(T(h_n))$ norm-converges to 0.

Second, assume $p = +\infty$. We apply claim (1) to the right-hand side of (0) to obtain

$$\limsup_{n \rightarrow \infty} \|T(h_n)\|^q \leq \frac{1}{t^q} [\max(1, t^q M^q) - (1 - \varepsilon)^q].$$

Considering here any $0 < \varepsilon < M^{-2q}$ and then taking $t = \varepsilon^{\frac{1}{2q}}$, we get that

$$\limsup_{n \rightarrow \infty} \|T(h_n)\|^q \leq \frac{1}{\varepsilon^{1/2}} [1 - (1 - \varepsilon)^q].$$

Now, letting $\varepsilon \rightarrow 0$ here, we get as before that the sequence $(T(h_n))$ norm-converges to 0. \square

The framework of this paper was inspired by [2]. The proof given in [2], devoted to the case $p < +\infty$, uses Stegall's variational principle.

REFERENCES

- [1] M. Fabian, P. Habala, P. Hájek, V. Montesinos Santalucía, J. Pelant and V. Zizler, *Functional analysis and infinite-dimensional geometry*, CMS Books in Mathematics, Springer-Verlag, New York, 2001. MR1831176 (2002f:46001)
- [2] M. Fabian and V. Zizler, *A "nonlinear" proof of Pitt's compactness theorem*, Proc. Amer. Math. Soc. **131** (2003), 3693–3694. MR1998188 (2004g:46026)

INSTITUT DE MATHÉMATIQUES DE BORDEAUX, UMR 5251, UNIVERSITÉ BORDEAUX I, 351,
COURS DE LA LIBÉRATION, 33405 TALENCE CEDEX, FRANCE
E-mail address: sylvain.delpech@gmail.com