

CONGRUENCE PROPERTIES OF HERMITIAN MODULAR FORMS

TOSHIYUKI KIKUTA AND SHOYU NAGAOKA

(Communicated by Wen-Ching Winnie Li)

In celebration of Tomoyoshi Ibukiyama's 60th birthday

ABSTRACT. We study the existence of a modular form satisfying a certain congruence relation. The existence of such modular forms plays an important role in the determination of the structure of a ring of modular forms modulo p . We give a criterion for the existence of such a modular form in the case of Hermitian modular forms.

1. INTRODUCTION

In [7], H. P. F. Swinnerton-Dyer determined the structure of a ring of modular forms mod p in the elliptic modular case. In his argument, the existence of a certain modular form plays an important role. Namely, he used the fact that there exists a modular form f of weight $p - 1$ with p -integral Fourier coefficients such that

$$f \equiv 1 \pmod{p}.$$

(Also cf. Serre [6].) In the elliptic modular case, such a form can be constructed easily. In fact, we may take $f = E_{p-1}$ (the normalized Eisenstein series of weight $p - 1$). However, the problem of existence in the case of Siegel modular forms turns out to be difficult. For example, the Siegel–Eisenstein series $E_{p-1}^{(n)}$ of weight $p - 1$ is no longer a solution in general. In [2], S. Boecherer and the second author studied this problem and gave some criteria for the existence problem in the case of Siegel modular forms.

In this paper, we give a criterion of the existence problem in the case of Hermitian modular forms over the imaginary quadratic fields $\mathbb{Q}(\sqrt{-1})$ and $\mathbb{Q}(\sqrt{-3})$.

2. HERMITIAN MODULAR FORMS

We start by recalling the definition of Hermitian modular forms. For details, please refer to [3]. The Hermitian half-space \mathbb{H}_n of degree n is defined by

$$\mathbb{H}_n := \left\{ Z \in M_n(\mathbb{C}) \mid \frac{1}{2\sqrt{-1}}(Z - {}^t\bar{Z}) > 0 \right\}.$$

Received by the editors April 1, 2008.

2000 *Mathematics Subject Classification.* Primary 11F33; Secondary 11F55.

Key words and phrases. Congruences for modular and p -adic modular forms.

The second author was supported in part by Grant-in-Aid for Scientific Research 19540061.

©2008 American Mathematical Society
Reverts to public domain 28 years from publication

Let \mathbb{K} be an imaginary quadratic field with discriminant $d_{\mathbb{K}}$. We denote by $\mathcal{O} = \mathcal{O}_{\mathbb{K}}$ the ring of integers and by \mathcal{O}^{\times} the group of units in \mathcal{O} .

The Hermitian modular group of degree n over \mathbb{K} ,

$$U_n(\mathcal{O}) := \{M \in M_{2n}(\mathcal{O}) \mid {}^t\overline{M}J_nM = J_n\}, \quad J_n = \begin{pmatrix} 0 & 1_n \\ -1_n & 0 \end{pmatrix},$$

acts on \mathbb{H}_n by

$$Z \longmapsto M \langle Z \rangle := (AZ + B)(CZ + D)^{-1}$$

for all $Z \in \mathbb{H}_n$ and $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in U_n(\mathcal{O})$.

Let $\Gamma \subset U_n(\mathcal{O})$ be a subgroup of $U_n(\mathcal{O})$. A holomorphic function $F(Z)$ on \mathbb{H}_n is called a Hermitian modular form of weight k for Γ if it satisfies the functional equations:

$$F(M \langle Z \rangle) = \det(CZ + D)^k F(Z)$$

for all $Z \in \mathbb{H}_n$ and $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma$. (We assume the holomorphy at the cusps in the case $n = 1$.) We denote by $M_k(\Gamma)$ the space of Hermitian modular forms of weight k for Γ . Later we mainly deal with the case $\Gamma = U_n(\mathcal{O})$ or $SU_n(\mathcal{O}) := U_n(\mathcal{O}) \cap SL_{2n}(\mathcal{O})$. In both cases, $F \in M_k(\Gamma)$ has a Fourier expansion of the form

$$F(Z) = \sum_{0 \leq T \in \Lambda_n} a_F(T) \exp\{2\pi\sqrt{-1}\text{tr}(TZ)\},$$

where T runs over the lattice

$$\Lambda_n = \Lambda_n(\mathbb{K}) := \{T = (t_{ij}) \in \text{Her}_n(\mathbb{K}) \mid t_{ii} \in \mathbb{Z}, \sqrt{d_{\mathbb{K}}}t_{ij} \in \mathcal{O}\}$$

(cf. [3]).

3. MAIN RESULT

In this section, we state the main result of this paper, which gives a criterion on the existence of a modular form satisfying a certain congruence relation.

Let p be a rational prime and $\mathbb{Z}_{(p)} = \{\frac{a}{b} \in \mathbb{Q} \mid p \nmid b\}$ denote the localization of \mathbb{Z} at p . We denote by $M_k(\Gamma)_{\mathbb{Z}_{(p)}}$ the subset of $M_k(\Gamma)$ consisting of $F \in M_k(\Gamma)$ such that all of its Fourier coefficients belong to $\mathbb{Z}_{(p)}$.

Our main result can be stated as follows:

Theorem 3.1. (1) Assume that $\mathbb{K} = \mathbb{Q}(\sqrt{-1})$ or $\mathbb{K} = \mathbb{Q}(\sqrt{-3})$. There exists a Hermitian modular form $F_{p-1} \in M_{p-1}(SU_n(\mathcal{O}))_{\mathbb{Z}_{(p)}}$ such that

$$F_{p-1} \equiv 1 \pmod{p}$$

if

$$p \equiv 1 \pmod{4}.$$

(2) Assume that $\mathbb{K} = \mathbb{Q}(\sqrt{-1})$. There exists a Hermitian modular form $F_{p-1} \in M_{p-1}(U_n(\mathcal{O}))_{\mathbb{Z}_{(p)}}$ such that

$$F_{p-1} \equiv 1 \pmod{p}$$

if and only if

$$p \equiv 1 \pmod{4}.$$

4. p -SPECIAL HERMITIAN MATRIX

In order to prove our theorem, we need to consider the existence of a p -special Hermitian matrix.

A Hermitian matrix $H = {}^t\overline{H} \in Her_m(\mathbb{K})$ is called p -special if H satisfies the following four conditions:

- (i) H is positive definite,
- (ii) H is even integral, namely, $H \in 2\Lambda_m$,
- (iii) $\det H = \left(\frac{2}{\sqrt{|d_{\mathbb{K}}|}}\right)^m$,
- (iv) (the main condition) there exists a p -group C_p in the finite unitary group

$$U_m(H; \mathcal{O}) := \{U \in M_m(\mathcal{O}) \mid {}^t\overline{U}HU = H\}$$

such that the group C_p acts freely on $\mathcal{O}^m \setminus \{\mathbf{0}\}$.

Our proof of the main result mainly depends on the existence of a p -integral Hermitian matrix. To demonstrate the existence of such a matrix we use the result of Bayer-Fluckiger, which guarantees the existence of a suitable even unimodular lattice over \mathbb{Z} .

Theorem 4.1 (Bayer-Fluckiger [1]). *Let m be a positive integer such that m is not a power of 2. Then there exists a definite unimodular lattice having an automorphism with characteristic polynomial Φ_m if and only if m is mixed and $\varphi(m)$ is divisible by 8. Here Φ_m is the m -th cyclotomic polynomial.*

Our result in this section is as follows:

Proposition 4.2. *Assume that $\mathbb{K} = \mathbb{Q}(\sqrt{-1})$ or $\mathbb{Q}(\sqrt{-3})$. If p is a prime number such that $p \equiv 1 \pmod{4}$, then there exists a p -special Hermitian matrix H of rank $p - 1$.*

Proof. First we assume that $\mathbb{K} = \mathbb{Q}(\sqrt{-1})$ and $p \equiv 1 \pmod{4}$. If we put $m = 4p$, then m is not a prime power and $\varphi(m) = 2(p - 1)$ is divisible by 8. Hence, by Theorem 4.1, there exists an even unimodular positive definite lattice (L, S) having an automorphism with characteristic polynomial Φ_m (m -th cyclotomic polynomial), where S is the associated bilinear form. We denote by t such an automorphism. The order of the automorphism t^p is 4, and L becomes a $\mathbb{Z}[\sqrt{-1}]$ -module by identifying $\sqrt{-1}$ with t^p . Since $\mathbb{Z}[\sqrt{-1}]$ is principal, one may construct a $\mathbb{Z}[\sqrt{-1}]$ -basis of L . Hence L becomes a $\mathbb{Z}[\sqrt{-1}]$ -lattice of rank $p - 1$. (The \mathbb{Z} -rank of L is $\varphi(m) = 2(p - 1)$.) The corresponding Gram matrix H is the desired matrix. Indeed, one can confirm that the Hermitian matrix H satisfies the conditions (i)-(iv) of the p -special Hermitian matrix.

- (i) The positivity of H comes from that of L .
- (ii) The bilinear form $S : L \times L \rightarrow \mathbb{Z}$ satisfies

$$S(\mathbf{x}, \mathbf{y}) = \frac{1}{2}(h(\mathbf{x}, \mathbf{y}) + h(\mathbf{y}, \mathbf{x})),$$

where $h : L \times L \rightarrow \mathbb{Q}(\sqrt{-1})$ is the Hermitian form associated with H . Since $h(\mathbf{x}, \mathbf{x}) = S(\mathbf{x}, \mathbf{x}) \in 2\mathbb{Z}$, H is even integral.

(iii) If we denote by $B \in \text{Sym}_{2(p-1)}(\mathbb{Z})$ the Gram matrix associated with the bilinear form S , then we have

$$\det B = (\det H)^2 \cdot \left(\frac{\sqrt{|d_{\mathbb{K}}|}}{2} \right)^{2(p-1)}.$$

Since $\det B = 1$ and $|d_{\mathbb{K}}| = 4$, we have $\det H = 1$. This shows that the matrix H satisfies the condition (iii).

(iv) We recall the definition of the automorphism t mentioned above. In this case, the p -group $C_p := \langle t^4 \rangle$ acts freely on $\mathcal{O}^{p-1} \setminus \{\mathbf{0}\}$ because the characteristic polynomial of t is Φ_m .

Next we assume that $\mathbb{K} = \mathbb{Q}(\sqrt{-3})$ and $p \equiv 1 \pmod{4}$. If we put $m = 3p$, then, by a similar argument to that stated above, there exists an even unimodular positive definite lattice (L, S) having an automorphism with characteristic polynomial Φ_m . We denote by s such an automorphism. The order of the automorphism s^p is 3, and L becomes an $\mathcal{O}_{\mathbb{K}}$ -module by identifying $\omega = \frac{-1+\sqrt{-3}}{2}$ with s^p . Since $\mathcal{O}_{\mathbb{K}} = \mathbb{Z} + \omega\mathbb{Z}$ is principal, one may construct an $\mathcal{O}_{\mathbb{K}}$ -basis of L . One can prove that the corresponding Gram matrix satisfies the conditions of the p -special Hermitian matrix in a way similar to the case $\mathbb{K} = \mathbb{Q}(\sqrt{-1})$. This completes the proof of Proposition 4.2. \square

Example 4.3. We give examples of H in the case $p = 5$:

The case $\mathbb{K} = \mathbb{Q}(\sqrt{-1})$,
$$H = \begin{pmatrix} 2 & 0 & 1 + \sqrt{-1} & \sqrt{-1} \\ 0 & 2 & \sqrt{-1} & 1 - \sqrt{-1} \\ 1 - \sqrt{-1} & -\sqrt{-1} & 2 & 0 \\ -\sqrt{-1} & 1 + \sqrt{-1} & 0 & 2 \end{pmatrix}.$$

The case $\mathbb{K} = \mathbb{Q}(\sqrt{-3})$,
$$H = \begin{pmatrix} 2 & 0 & \frac{2}{\sqrt{-3}} & \frac{2}{\sqrt{-3}} \\ 0 & 2 & \frac{2}{\sqrt{-3}} & \frac{-2}{\sqrt{-3}} \\ \frac{-2}{\sqrt{-3}} & \frac{-2}{\sqrt{-3}} & 2 & 0 \\ \frac{-2}{\sqrt{-3}} & \frac{2}{\sqrt{-3}} & 0 & 2 \end{pmatrix}.$$

5. PROOF OF THE MAIN THEOREM

In this section, we prove our main theorem.

Proof. (1) We assume that $\mathbb{K} = \mathbb{Q}(\sqrt{-1})$ or $\mathbb{K} = \mathbb{Q}(\sqrt{-3})$ and that $p \equiv 1 \pmod{4}$. By Proposition 4.2, there exists a p -special Hermitian matrix H of rank $p - 1$. We denote by C_p the corresponding p -group (cf. section 4, the definition of a p -special Hermitian matrix (iv)). We associate the theta series

$$\vartheta_H(Z) := \sum_{X \in M_{p-1,n}(\mathcal{O})} \exp\{\pi\sqrt{-1} \text{tr}(H[X]Z)\}, \quad Z \in \mathbb{H}_n,$$

where $H[X] := {}^t\overline{X}HX$. The modularity of ϑ_H for $SU_n(\mathcal{O})$ comes from the conditions (i), (ii), and (iii) of the p -special matrix $H \in 2\Lambda_{p-1}$ (e.g. cf. Cohen and Resnikoff [4], p. 332); namely, we have $\vartheta_H(Z) \in M_{p-1}(SU_n(\mathcal{O}))$. In particular, we have $\vartheta_H(Z) \in M_{p-1}(U_n(\mathcal{O}))$ in the case that $\mathbb{K} = \mathbb{Q}(\sqrt{-1})$ because

$\sharp\mathcal{O}^\times = \sharp\mathbb{Z}[\sqrt{-1}]^\times = 4$ and the weight $p - 1$ is divisible by 4. The Fourier expansion is given as follows:

$$\vartheta_H(Z) = \sum_T A(H, T) \exp\{2\pi\sqrt{-1} \operatorname{tr}(TZ)\},$$

$$A(H, T) = \sharp\mathcal{A}(H, T), \quad \mathcal{A}(H, T) = \{X \in M_{p-1,n}(\mathcal{O}) \mid H[X] = 2T\}.$$

If $T \neq O_n$, then the p -group C_p acts freely on the set $\mathcal{A}(H, T)$. Therefore, the number $A(H, T)$ is divisible by p . Since $A(H, O_n) = 1$, we have

$$\vartheta_H(Z) \equiv 1 \pmod{p}.$$

This proves (1) of Theorem 3.1.

(2) Assume that $\mathbb{K} = \mathbb{Q}(\sqrt{-1})$ and that there exists a form

$$F_{p-1} \in M_{p-1}(U_n(\mathbb{Z}[\sqrt{-1}]))_{\mathbb{Z}(p)}$$

such that

$$F_{p-1} \equiv 1 \pmod{p}.$$

We recall the definition of the Φ -operator defined by

$$\Phi : M_k(U_n(\mathcal{O})) \longrightarrow M_k(U_{n-1}(\mathcal{O})), \quad \Phi(F)(Z) := \lim_{\lambda \rightarrow \infty} F \left(\begin{pmatrix} Z & 0 \\ 0 & i\lambda \end{pmatrix} \right), \quad Z \in \mathbb{H}_{n-1}.$$

If we apply the Φ -operator $n - 1$ times to F_{p-1} , then

$$\Phi^{(n-1)}(F_{p-1}) \in M_{p-1}(U_1(\mathbb{Z}[\sqrt{-1}]))_{\mathbb{Z}(p)}$$

still satisfies the congruence relation

$$\Phi^{(n-1)}(F_{p-1}) \equiv 1 \pmod{p}.$$

If $p \not\equiv 1 \pmod{4}$, this is impossible because

$$M_k(U_1(\mathbb{Z}[\sqrt{-1}])) = \begin{cases} M_k(SL_2(\mathbb{Z})) & \text{if } k \equiv 0 \pmod{4}, \\ 0 & \text{otherwise.} \end{cases}$$

(This comes from the fact that $U_1(\mathcal{O}) = \mathcal{O}^\times \cdot SL_2(\mathbb{Z})$.) We have proved the statement (2), thereby completing the proof of Theorem 3.1. \square

6. REMARK

In the case that $\mathbb{K} = \mathbb{Q}(\sqrt{-1})$ and $n = 2$, there is another construction of F_{p-1} , which is based on the theory of Hermitian Jacobi forms.

We assume that $\mathbb{K} = \mathbb{Q}(\sqrt{-1})$. Freitag [5] constructed a set of generators of the graded ring

$$M^{sym}(U_2(\mathbb{Z}[\sqrt{-1}])) = \bigoplus_k M_k^{sym}(U_2(\mathbb{Z}[\sqrt{-1}])),$$

where $M_k^{sym}(U_2(\mathbb{Z}[\sqrt{-1}]))$ is the subspace consisting of the symmetric Hermitian modular forms of weight k . (In general, $F \in M_k(U_n(\mathcal{O}))$ is called symmetric if $F({}^tZ) = F(Z)$.) We recall the weight 4 generator φ_4 of $M^{sym}(U_2(\mathbb{Z}[\sqrt{-1}]))$ (cf. [5]). It is known that all the Fourier coefficients of $\mathcal{E}_4 := \frac{1}{4}\varphi_4$ are integral and the constant term is equal to 1. We expand \mathcal{E}_4 as a Fourier-Jacobi series and take the index 1 Jacobi form $\Phi_{4,1}$. All of the Fourier coefficients of $\Phi_{4,1}$ are divisible by 240. We put $\phi_{4,1} := \frac{1}{240}\Phi_{4,1}$. Now we assume that $p \equiv 1 \pmod{4}$. Then

$$f_{p-1,1} := E_4^{\frac{p-5}{4}} \cdot \phi_{4,1}$$

becomes a Hermitian Jacobi form of weight $p - 1$ and index 1. Here $E_4 = 1 + 240 \sum_{n=1}^{\infty} \sigma_3(n)q^n$ is the ordinary Eisenstein series of weight 4 for $SL_2(\mathbb{Z})$. All of the Fourier coefficients of $f_{p-1,1}$ are integral and the constant term is equal to 1. We consider the Maass lift \mathcal{M}_k from the space of Hermitian Jacobi forms of weight k and index 1 to the space $M_k(U_2(\mathbb{Z}[\sqrt{-1}]))$. Then

$$F_{p-1} := -\frac{2(p-1)}{B_{p-1}} \mathcal{M}_{p-1}(f_{p-1,1})$$

is the desired form, namely, $F_{p-1} \in M_{p-1}(U_2(\mathbb{Z}[\sqrt{-1}]))_{\mathbb{Z}(p)}$ and

$$F_{p-1} \equiv 1 \pmod{p}.$$

(Since the Maass lift \mathcal{M}_k is defined only for k such that $k \equiv 0 \pmod{4}$ in this case, we need the assumption $p \equiv 1 \pmod{4}$.)

ACKNOWLEDGEMENT

The authors wish to thank Prof. G. Nebe for suggesting the proof of Proposition 4.2.

REFERENCES

1. E. Bayer-Fluckiger, *Definite unimodular lattices having an automorphism of given characteristic polynomial*. Comment. Math. Helv. **59** (1984), 509–538. MR780074 (86k:11032)
2. S. Boecherer, S. Nagaoka, *On mod p properties of Siegel modular forms*. Math. Ann. **338** (2007), 421–433. MR2302069 (2008d:11041)
3. H. Braun, *Hermitian modular functions*. Ann. of Math. (2) **50** (1949), 827–855. MR0032699 (11:333a)
4. D. M. Cohen, H. L. Resnikoff, *Hermitian quadratic forms and Hermitian modular forms*. Pac. J. Math. **76** (1978), 329–337. MR506135 (80b:10039)
5. E. Freitag, *Modulformen zweiten Grades zum rationalen und Gausschen Zahlkörper*. Sitzungsber. Heidelberger Akad. Wiss. Math.-Natur. Kl. (1967), 3–49. MR0214541 (35:5391)
6. J-P. Serre, *Formes modulaires et fonctions zêta p -adiques*. Modular Functions of One Variable III, Lecture Notes in Math. **350** (1973), 191–268, Springer. MR0404145 (53:7949a)
7. H. P. F. Swinnerton-Dyer, *On ℓ -adic representations and congruences for coefficients of modular forms*. Modular Functions of One Variable III, Lecture Notes in Math. **350** (1973), 1–55, Springer. MR0406931 (53:10717a)

DEPARTMENT OF MATHEMATICS, KINKI UNIVERSITY, HIGASHI-OSAKA, OSAKA 577-8502, JAPAN
E-mail address: kikuta@math.kindai.ac.jp

DEPARTMENT OF MATHEMATICS, KINKI UNIVERSITY, HIGASHI-OSAKA, OSAKA 577-8502, JAPAN
E-mail address: nagaoka@math.kindai.ac.jp