

ON GENERICITY AND WEIGHT IN THE FREE GROUP

ANAND PILLAY

(Communicated by Julia Knight)

ABSTRACT. We prove that the generic type of the (theory of the) free group F_n on $n \geq 2$ generators has infinite weight, strengthening the well-known result that these free groups are not superstable. A preliminary result, possibly of independent interest, is that the realizations in F_n of the generic type are precisely the primitives.

1. INTRODUCTION AND PRELIMINARIES

From the work by Sela [10] and Kharlampovich and Myasnikov [2], we know that all nonabelian free groups have the same elementary (or first order) theory, which we call T_{fg} . Sela [11] also recently proved that T_{fg} is stable. It has been known for a long time [9] that T_{fg} is not superstable. Rather recently, influenced by work on theories without the independence property, the notion of “strongly stable” has aroused interest. A theory T is said to be *strongly stable* if it is stable and every (finitary) type has finite weight (see below). Any superstable theory is strongly stable. It is natural to ask whether the free group is strongly stable. We show here that it is not. In fact we prove that in T_{fg} the *generic type* has infinite weight, strengthening our observation in [8] that the generic type has weight ≥ 2 . On the way to proving this we will show that the realizations in F_n of the generic type are precisely the primitives. Our proof makes use of recent work by Perin [6].

Many thanks to both Sasha Ivanov and Zlil Sela for helpful discussions and for pointing out relevant results in the literature. In particular, Ivanov explained to me how the special case “weight of generic type is ≥ 3 ” followed from our methods just by using a result of Nies [5] that F_2 is homogeneous. Thanks also to the referee and to Rizos Sklinos for helpful comments on a first version.

In the remainder of this section, I recall pertinent facts about free groups, stable groups, and weight. The main results and their proofs (quite easy) appear in section 2.

1.1. Free groups. Let F_n denote the free group on n generators, where usually $n \geq 2$. Insofar as we consider F_n as a structure in the sense of model theory, it will be in the language of groups $\{\cdot, {}^{-1}, e\}$. Sometimes we write ab in place of $a \cdot b$.

Received by the editors December 9, 2008, and, in revised form, March 3, 2009.

2000 *Mathematics Subject Classification.* Primary 03C45; Secondary 20F67.

This work was supported by Marie Curie Chair EXC 024052 as well as EPSRC grant EP/F009712/1.

©2009 American Mathematical Society
Reverts to public domain 28 years from publication

It is well-known that F_n is isomorphic to F_m iff $m = n$. If F_n is free on generators a_1, \dots, a_n , we call $\{a_1, \dots, a_n\}$ a basis of F_n , and an element of a basis is called a *primitive* of F_n . So the primitives of F_n form a single orbit under $\text{Aut}(F_n)$.

We will be making heavy use of the following fact, which follows easily from Whitehead's theorem, although we give an explanation.

Fact 1.1. Let $\{a_1, \dots, a_n\}$ be a basis of F_n . Let $m \leq n$ and let k_1, \dots, k_m be integers > 1 . Then $a_1^{k_1} \cdots a_m^{k_m}$ is not a primitive of F_n .

Explanation. Let $A = \{a_1, \dots, a_n\}$. A^{-1} denotes the set of inverses of A . An automorphism α of F_n is called a Whitehead automorphism if it is induced by either (i) a permutation of $A \cup A^{-1}$, or (ii) for some i , a map which fixes a_i , and for each $j \neq i$ takes a_j to a_j , $a_j a_i$, $a_i^{-1} a_j$, or $a_i^{-1} a_j a_i$. A word w (in the basis elements) is said to be cyclically reduced if it is reduced, and not of the form vcv^{-1} . A cyclic word is a cyclically reduced word, defined up to cyclic permutation. For w a reduced word, $l(w)$ denotes its length. Whitehead's theorem, which is Theorem 4 in section 10 of [1] or Proposition 4.17 of [3] says that if w, u are cyclic words which are in the same orbit of $\text{Aut}(F_n)$ and such that moreover $l(u)$ is minimal among lengths of words in this orbit, then there is a sequence τ_1, \dots, τ_s of Whitehead automorphisms of F_n such that for each $i = 1, \dots, s$, $l(w) \geq l(\tau_i \dots \tau_1 w)$. Moreover if $l(w) \neq l(u)$, then each of these inequalities is strict.

Now let w be our word $a_1^{k_1} \cdots a_m^{k_m}$. Clearly w is cyclically reduced and so is a cyclic word. We want to show that w is not primitive, namely is not in the same orbit under $\text{Aut}(F_n)$ as a_1 . Supposing otherwise, put $u = a_1$, also a cyclic word, clearly of minimal length in its orbit. So Whitehead's theorem applies to w and u . But by inspection, $l(\alpha w) \geq l(w)$ for any Whitehead automorphism of F_n , which gives a contradiction.

In [10], Sela proved:

Theorem 1.2. *If $m \geq n \geq 2$, then the natural embedding of F_n in F_m is an elementary embedding.*

Of course, this not only solves Tarski's problem on the elementary equivalence of finite rank nonabelian free groups, but also shows that the natural embeddings of F_κ into F_λ (for any finite or infinite cardinals $\kappa < \lambda$) are also elementary embeddings.

On the other hand, Chloé Perin [6] has recently proved a converse to Theorem 1.2:

Theorem 1.3. *Let $n \geq 2$ and let G be an elementary substructure of F_n . Then G is a free factor of F_n . Namely G is a free group of rank at most n , and some (any) basis of G extends to a basis of F_n .*

As mentioned in the introduction we let T_{fg} denote the common theory of non-abelian free groups of finite rank, which we know now to be complete. Sela [11] has also proved the striking result:

Theorem 1.4. *T_{fg} is stable.*

Some basic stability-theoretic properties of the free group will be discussed in the next section.

1.2. Stability. For basic model theory we refer the reader to [4]. In my recent paper [8] on stability-theoretic aspects of the free group, I gave a brief survey of stability and stable groups, directed towards nonexperts, so rather than repeat

myself I will direct readers to the introduction of that paper. Further facts about stability and stable groups can be found in [7].

Let us fix a complete countable stable theory T , and a model M . We let \bar{M} be a saturated elementary extension of M (so we allow $M = \bar{M}$). From stability theory we have the notion “ a is independent from b over C ” for a, b tuples from M (or even subsets of M) and C a subset of M . Technically the notion is “ $tp(a/C \cup \{b\})$ does not fork over C ”, where forking is as defined by Shelah. In any case “ a is independent from b over C ” is synonymous with “ a is independent from $C \cup \{b\}$ over C ”. We also may say “ a forks with b over C ” in place of “ a is not independent from b over C ”.

Among key properties of independence in a stable theory are

(a) (symmetry) a is independent from b over C iff b is independent from a over C .

(b) (transitivity) If $A \subseteq B \subseteq C$, then a is independent from C over A iff a is independent from C over B and a is independent from B over A .

(c) (local character) If a is a finite tuple and C any set, then there is a countable $C_0 \subseteq C$ such that a is independent from C over C_0 .

(d) (invariance) Whether or not a is independent from b over C depends on $tp(a, b, C)$ (so working in \bar{M} the notion is invariant under automorphism).

(e) (existence) Given a and C and $B \supseteq C$, there is $a' \in \bar{M}$ such that $tp(a'/C) = tp(a/C)$ and a' is independent from B over C .

(f) (uniqueness) If C is “algebraically closed in M^{eq} ” and a is a tuple, then $tp(a/C)$ is *stationary*, meaning that for any $B \supseteq C$, if a', a'' are such that $tp(a'/C) = tp(a''/C) = tp(a/C)$ and each of a', a'' is independent from B over C , then $tp(a'/B) = tp(a''/B)$.

Remark 1.5. It would be interesting to give a relatively explicit description of independence when M is a free group. For $C = \emptyset$, and a, b “generics” this will be done below.

Let $A \subseteq M$ and $\{b_i : i \in I\}$ be a set of tuples from M . We will say that $\{b_i : i \in I\}$ is A -independent, if for each $i \in I$, b_i is independent from $\{b_j : j \neq i\}$ over A . Let us note a couple of things. Firstly, if κ is a cardinal and $(b_i : i < \kappa)$ has the property that for each i , b_i is independent from $\{b_j : j < i\}$ over A , then in fact $\{b_i : i < \kappa\}$ is A -independent (by symmetry and transitivity of independence).

Secondly, if $p(x) \in S(A)$ is stationary, then any two A -independent sequences $(b_i : i < \kappa)$, $(c_i : i < \kappa)$ of realizations of p , have the same type over A . In particular $(b_i : i < \kappa)$ is totally indiscernible over A (every permutation is an elementary map over A in the sense of the ambient model M).

I will now give a definition of weight in a countable stable theory T . See section 4.4 of Chapter 1 of [7] for more details.

Definition 1.6. (i) Let a be a finite tuple from M and A a subset of M . The *preweight* of $tp(a/A)$ (written $pw(tp(a/A))$ or even $pw(a/A)$) is the supremum of the cardinals κ such that in \bar{M} there exists an A -independent set $\{b_i : i < \kappa\}$ such that a forks with each b_i over A .

(ii) For a, M, A as in (i), the *weight* of $tp(a/A)$, written $w(tp(a/A))$ or $w(a/A)$, is the supremum of the preweights of nonforking extensions of $tp(a/A)$, equivalently the supremum of the set of $w(a/B)$, for $B \subseteq \bar{M}$, $B \supseteq A$, and a independent from B over A .

Fact 1.7. For any a, A as in Definition 1.6, $w(a/A) \leq \omega$. Moreover, if $w(a/A) = \omega$, then for some $B \supseteq A$, possibly from \bar{M} , such that a is independent from B over A , the supremum is achieved for $tp(a/B)$; namely, there is a B -independent set $\{c_i : i < \omega\}$ such that a forks with each c_i over B .

In [7] we called a stable theory *thin* if every finitary type (namely type of some finite tuple over some set) has finite weight. Following work of Shelah [12], this is now called *strongly stable*. Any superstable theory is strongly stable. Considerable work has been done recently on theories without the independence property (called NIP theories, or dependent theories). Although there is no obvious notion of “superdependent”, Shelah gave a definition of “strongly dependent”, which for stable theories is precisely strongly stable. This explains some of the current interest in weight.

Example 1.8. (i) In a strongly minimal theory such as the theory of algebraically closed fields of a fixed characteristic, $w(b/A) = pw(b/A)$ and is the same as the Morley rank of $tp(b/A)$ (b a finite tuple, A a set of parameters).

(ii) If $p(x)$ is a complete stationary type over \emptyset of weight 1 in a stable theory T , then in any model M of T any two maximal independent sets of realizations of p have the same cardinality.

(iii) If p is a complete type of U -rank ω^α (in a stable theory), then $w(p) = 1$.

(iv) In the structure $(\mathbb{Z}/4\mathbb{Z}^\omega, +)$ the generic type (see below) has Morley rank 2 but weight 1.

(v) There are strongly stable but nonsuperstable groups, such as a vector space over \mathbb{Q} equipped with predicates for members of an infinite strictly descending chain of subspaces.

(vi) The generic type (see below) of a separably closed field F of infinite Ersov invariant has weight 1, although $Th(F)$ is nonsuperstable.

For our purposes a *stable group* is a definable group in a stable theory T . Namely, there are formulas $\phi(x)$ and $\psi(x, y, z)$ such that in some (any) model M of T , the set of solutions of ψ is the graph of a group operation on the set of solutions of ϕ . We will assume these formulas have no parameters, so the free group is a stable group, with $T = T_{fg}$. In a stable group, the theory of independence above has an equivariant variation, leading to the theory of generic types. Let us fix again a model M of T and let G be the interpretation of the relevant formulas in M . There are two equivalent definitions of a “generic type” or “generic element” of G : (i) Let $A \subseteq M$. Then $g \in G$ is a generic element of G over A , or $tp(g/A)$ is a generic of G , if for any A -definable subset X of G containing g , finitely many left translates of X cover G . (ii) Again for $A \subseteq M$, g is generic in G over A if (working possibly in \bar{M}) whenever $h \in G(\bar{M})$ is independent from g over A , then $h \cdot g$ is independent from h over A .

It is a fact that if $tp(g/A)$ is generic, then g is independent from A over \emptyset , and also that if $tp(g/A)$ is generic and g is independent from B over A , then $tp(g/B)$ is generic.

Let us remark on two additional properties of generics in stable groups: (iii) If $g \in G$ is generic over A , then g^{-1} is generic over A . (iv) If $g \in G$ is generic over A and $h \in G \cap A$, then each of gh and hg is generic over A .

A stable group is said to be *connected* if it has no proper definable subgroup of finite index. Again it is a basic fact that G is connected if and only if there

is a unique generic type of G : namely, working possibly in \bar{M} , for any set A of parameters there is a unique $tp(g/A)$ with g generic in G over A . For G connected, we denote by $p_0^G(x)$ the unique generic type of G over \emptyset . This type will be *stationary*, and for any set A of parameters, its unique nonforking extension over A will coincide with the unique generic type of G over A . In particular for any cardinal (maybe finite) κ , an independent set $\{a_i : i \in I\}$ of realizations of p_0^G is an indiscernible set. It is not hard to conclude, using also properties (iii) and (iv) above, that the Whitehead transformations applied to such an independent set are elementary maps. More precisely:

Fact 1.9. Suppose G is a connected stable group. Let $A = \{a_i : i \in I\}$ be an independent set of realizations of $p_0^G(x)$ in G . Let π be one of the following maps:

- (i) for some permutation σ of I , $\pi(a_i) = a_{\sigma(i)}$ or $a_{\sigma(i)}^{-1}$,
- (ii) for some $i \in I$, $\pi(a_i) = a_i$ and for every $j \neq i$, $\pi(a_j)$ is a_j , $a_j \cdot a_i$, $a_i^{-1} \cdot a_j$, or $a_i^{-1} \cdot a_j \cdot a_i$.

Then π is an elementary map in the sense of G . In particular $\{\pi(a_i) : i < \kappa\}$ is also an independent set of realizations of p_0^G .

The following was observed in [8], and moreover the results were shown to follow from Theorems 1.2 and 1.4, using elementary arguments due to Poizat [9].

Fact 1.10. (i) The free group is connected.

(ii) If F is a free group with basis $\{a_i : i \in I\}$, then $\{a_i : i \in I\}$ is an independent set of realizations of the unique generic type p_0 .

In fact it is worth remarking briefly that Poizat's main observation was that working in F_ω with free basis $\{a_i : i < \omega\}$, a definable subset X is generic (finitely many translates cover F_ω) if and only if X contains all but finitely many of the a_i . If we know in addition Theorems 1.2 and 1.4 of Sela, then Fact 1.10 follows quickly.

Bearing in mind that any two bases of a free group have the same cardinality, part (ii) above together with Example 1.7 (ii) might be considered evidence that the generic type of the free group has weight 1. In fact, in [8] we already pointed out that the generic type has weight at least two, simply because a generic in the free group is a product of two nongenerics. In the next section we will prove that in fact the generic type of the free group has infinite weight.

2. MAIN RESULTS

As above, T_{fg} is the theory of the noncommutative free group, and $p_0(x)$ is the generic type of T_{fg} over \emptyset . In fact we will be working entirely in "standard models" of T_{fg} , namely free groups of finite rank.

Our first result is a kind of converse to Fact 1.10 (ii).

Theorem 2.1. (i) Let F be a free group of finite rank (≥ 2). Then any realization of $p_0(x)$ in F is a primitive.

(ii) Any maximal independent set of realizations of $p_0(x)$ in F is a basis of F .

Proof. (i) Suppose F is free of rank n , with basis $\{a_1, \dots, a_n\}$, and consider F as a subgroup of F_{n+1} where the latter has basis $\{a_1, \dots, a_n, a_{n+1}\}$. By Theorem 1.2, F is an elementary substructure of F_{n+1} . Let b realize p_0 in F . So b also realizes p_0 in F_{n+1} .

Claim I. b, a_{n+1} are independent realizations of p_0 in F_{n+1} .

Proof. Work in F_{n+1} . By Fact 1.10(ii), a_{n+1} realizes p_0 , and is moreover independent from $\{a_1, \dots, a_n\}$ over \emptyset . But $b \in \text{dcl}(a_1, \dots, a_n)$, so a_{n+1} is independent from b over \emptyset . This suffices.

Let G be the subgroup of F_{n+1} generated by $\{b, a_{n+1}\}$.

Claim II. (b, a_{n+1}) has the same type in G as in F_{n+1} .

Proof. By stationarity of p_0 , the type of (b, a_{n+1}) in F_{n+1} is the same as the type of a basis of F_2 in F_2 . Hence G is free with basis (b, a_{n+1}) , and Claim II follows.

As $\{b, a_{n+1}\}$ generates G , it follows from Claim II that G is an elementary substructure of F_{n+1} . By Theorem 1.3, G is a free factor of F_{n+1} , whence $\{b, a_{n+1}\}$ extends to a basis $\{b, a_{n+1}, c_1, \dots, c_{n-1}\}$ of F_{n+1} . Let $\phi : F_{n+1} \rightarrow F$ be the surjective homomorphism defined by: ϕ is the identity on F and $\phi(a_{n+1}) = 1$. So $\phi(b) = b$, and $\{b, \phi(c_1), \dots, \phi(c_{n-1})\}$ generates F . As F is free of rank n , by Proposition 2.7 of [3], for example, $\{b, \phi(c_1), \dots, \phi(c_n)\}$ is a basis of F . So b is a primitive of F .

(ii) Let I be a maximal independent set of realizations of p_0 in F . By part (i) and Fact 1.10(ii), $|I| > 1$. As in the proof of part (i), the group G generated by I is an elementary substructure of F and is moreover free on I . By Theorem 1.3 again, G is a free factor of F whereby I extends to a basis of F . Again using Fact 1.10 and our maximality assumption on I , $G = F$, so I is a basis of F . The proof is complete.

Remark 2.2. By part (ii) of the proposition, any two maximal independent sets of realizations of $p_0(x)$ in a “free, finite rank” model of T_{fg} have the same cardinality, which is again a kind of “weight 1” behaviour.

Theorem 2.3. *The generic type $p_0(x)$ of T_{fg} has infinite weight.*

Proof. For each $n \geq 2$ we will find a realization g of p_0 in F_n and independent realizations b_1, \dots, b_n of p_0 in F_n such that g depends on (forks with) b_i for each $i = 1, \dots, n$. We will be systematically using Fact 1.9 to check that certain elements we construct are generic, or even independent sets of generics. In fact we could equally well observe that our “transformations” are taking bases to bases, and hence by Fact 1.10(ii) to independent sets of generics. In any case we will just say “by Fact 1.9”.

Let us fix a basis $\{a_1, \dots, a_n\}$ of F_n . Let $g = a_1 a_2^3 a_3^3 \dots a_i^3 \dots a_n^3$. Then

Claim 1. g realizes $p_0(x)$.

Proof. By Fact 1.9.

Now let $b_1 = a_1$, $b_2 = a_1 a_2$, and for $i = 3, \dots, n$, $b_i = a_1 a_2^3 \dots a_{i-1}^3 a_i$.

Claim 2. $\{b_1, \dots, b_n\}$ is an independent set of realizations of p_0 .

Proof. Again by Fact 1.9. In fact one sees directly that $\{b_1, \dots, b_n\}$ is a basis of F_n .

Claim 3. g forks with b_i for each $i = 1, \dots, n$.

Proof. Suppose for a contradiction that g is independent from b_i . By Fact 1.9, $b_i^{-1}g$ realizes p_0 , so in particular by Theorem 2.1(i) is a primitive element in F_n . Suppose first that $i = 1$. As $b_1 = a_1$, $b_1^{-1}g = a_2^3 \dots a_n^3$, but by Fact 1.1 the latter is

not primitive, a contradiction. Now suppose $i > 1$. Then $b_i^{-1}g = a_i^2 a_{i+1}^3 \dots a_n^3$ is also not primitive by Fact 1.1, again a contradiction.

Claims 1, 2 and 3 show that $pw(p_0) \geq n$. As n was arbitrary, $w(p_0) = \omega$, proving the theorem.

Note that the proof of Theorem 2.3 says that the generic type has arbitrarily large weight with respect to itself. In fact we can conclude from the proof a more precise version.

Corollary 2.4. *In some model G of T_{fg} , there is a realization g of p_0 and an independent set $\{b_i : i < \omega\}$ of realizations of p_0 such that g forks with each b_i over \emptyset .*

Proof. Let c_1, c_2, c_3, \dots be independent realizations of p_0 in a saturated model. Let $\Sigma(x, y_1, y_2, \dots)$ be a set of formulas expressing that x realizes p_0 , $\{y_1, y_2, \dots\}$ is an independent set of realizations of p_0 and that $x^{-1}y_n = c_1^3 \dots c_{n-1}^3 c_n^2$ for all $n = 1, 2, \dots$. The proof of Theorem 2.3 shows that Σ is consistent, so realized by g, b_1, b_2, \dots , and again using Fact 1.1, g forks with each b_i over \emptyset .

Remark 2.5. Can we find such a g and $(b_i)_i$ (as in the Corollary) in F_ω ? We conjecture not.

REFERENCES

1. D. E. Cohen, *Combinatorial Group Theory: A Topological Approach*, London Math. Soc. Student Texts 14, Cambridge University Press, 1989. MR1020297 (91d:20001)
2. O. Kharlampovich and A. Myasnikov, Elementary theory of free nonabelian groups, *J. Algebra* 302 (2006), 451-552. MR2293770 (2008e:20033)
3. R. C. Lyndon and P. E. Schupp, *Combinatorial Group Theory*, Springer-Verlag, 1977. MR0577064 (58:28182)
4. D. Marker, *Model Theory: An Introduction*, Springer, 2002. MR1924282 (2003e:03060)
5. A. Nies, Aspects of free groups, *Journal of Algebra* 263 (2003), 119-125. MR1974081 (2004g:20033)
6. C. Perin, Plongements élémentaires dans un groupe hyperbolique sans torsion, Ph.D. thesis, Caen, October 2008.
7. A. Pillay, *Geometric Stability Theory*, Oxford University Press, 1996. MR1429864 (98a:03049)
8. A. Pillay, Forking in the free group, *Journal of the Inst. of Math. Jussieu* 7 (2008), 375-389. MR2400726
9. B. Poizat, Groupes stables, avec types génériques réguliers, *Journal of Symbolic Logic* 48 (1983), 339-355. MR704088 (85e:03082)
10. Z. Sela, Diophantine geometry over groups, VI: The elementary theory of a free group, *Geom. Funct. Anal.* 16 (2006), 707-730. MR2238945 (2007j:20063)
11. Z. Sela, Diophantine geometry over groups, VIII: Stability (arXiv:math/0609096v1).
12. S. Shelah, Strongly dependent theories (arXiv:math/0504197v2), to appear.

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS, LS2 9JT, UNITED KINGDOM
E-mail address: pillay@maths.leeds.ac.uk