

STABILITY CRITERION FOR CONVOLUTION-DOMINATED INFINITE MATRICES

QIYU SUN

(Communicated by Marius Junge)

ABSTRACT. Let ℓ^p be the space of all p -summable sequences on \mathbb{Z} . An infinite matrix is said to have ℓ^p -stability if it is bounded and has bounded inverse on ℓ^p . In this paper, a practical criterion is established for the ℓ^p -stability of convolution-dominated infinite matrices.

1. INTRODUCTION

Let \mathcal{C} be the Gohberg-Baskakov-Sjöstrand class of infinite matrices $A := (a(j, j'))_{j, j' \in \mathbb{Z}}$ with

$$\|A\|_{\mathcal{C}} = \sum_{k \in \mathbb{Z}} \sup_{j - j' = k} |a(j, j')| < \infty.$$

Let $\ell^p := \ell^p(\mathbb{Z})$ be the set of all p -summable sequences on \mathbb{Z} with the standard norm $\|\cdot\|_p$. An infinite matrix $A := (a(j, j'))_{j, j' \in \mathbb{Z}} \in \mathcal{C}$ defines a bounded linear operator on ℓ^p , $1 \leq p \leq \infty$, in the sense that

$$(1.1) \quad Ac = \left(\sum_{j' \in \mathbb{Z}} a(j, j')c(j') \right)_{j \in \mathbb{Z}},$$

where $c = (c(j))_{j \in \mathbb{Z}} \in \ell^p$. Given a summable sequence $h = (h(j))_{j \in \mathbb{Z}} \in \ell^1$, define the *convolution operator* C_h on ℓ^p , $1 \leq p \leq \infty$, by

$$(1.2) \quad C_h : \ell^p \ni (b(j))_{j \in \mathbb{Z}} \mapsto \left(\sum_{k \in \mathbb{Z}} h(j - k)b(k) \right)_{j \in \mathbb{Z}} \in \ell^p.$$

Observe that the linear operator associated with an infinite matrix $A \in \mathcal{C}$ is dominated by a convolution operator in the sense that

$$(1.3) \quad |(Ac)(j)| \leq (C_h|c|)(j) := \sum_{j' \in \mathbb{Z}} h(j - j')|c(j')|, \quad j \in \mathbb{Z},$$

for any sequence $c = (c(j))_{j \in \mathbb{Z}} \in \ell^p$, $1 \leq p \leq \infty$, where $|c| = (|c(j)|)_{j \in \mathbb{Z}}$ and the sequence $(\sup_{j - j' = k} |a(j, j')|)_{k \in \mathbb{Z}}$ can be chosen to be the sequence $h = (h(j))_{j \in \mathbb{Z}}$ in (1.3). So infinite matrices in the set \mathcal{C} are said to be *convolution-dominated*.

Convolution-dominated infinite matrices were introduced by Gohberg, Kaashoek, and Woerdeman [12] as a generalization of Toeplitz matrices. They showed that the class \mathcal{C} equipped with the standard matrix multiplication and the above norm $\|\cdot\|_{\mathcal{C}}$

Received by the editors October 14, 2008 and, in revised form, November 30, 2009.

2010 *Mathematics Subject Classification*. Primary 47B35; Secondary 40E05, 65F05, 42C40, 47G30, 94A20.

©2010 American Mathematical Society
 Reverts to public domain 28 years from publication

is an inverse-closed Banach subalgebra of $\mathcal{B}(\ell^p)$ for $p = 2$. Here $\mathcal{B}(\ell^p)$, $1 \leq p \leq \infty$, is the space of all bounded linear operators on ℓ^p with the standard operator norm, and a subalgebra \mathcal{A} of a Banach algebra \mathcal{B} is said to be *inverse-closed* if when an operator $T \in \mathcal{A}$ has an inverse T^{-1} in \mathcal{B} , then $T^{-1} \in \mathcal{A}$ ([7, 11, 21]). The inverse-closed property for convolution-dominated infinite matrices was rediscovered by Sjöstrand [25] with a completely different proof and an application to a deep theorem about pseudodifferential operators. Recently Shin and Sun [23] generalized Gohberg, Kaashoek and Woerdeman's result and proved that the class \mathcal{C} is an inverse-closed Banach subalgebra of $\mathcal{B}(\ell^p)$ for any $1 \leq p \leq \infty$. The readers may refer to [5, 10, 20, 23, 25, 27] and the references therein for related results and various generalizations on the inverse-closed property for convolution-dominated infinite matrices.

Convolution-dominated infinite matrices arise and have been used in the study of spline approximation ([8, 9]), wavelets and affine frames ([6, 18]), Gabor frames and non-uniform sampling ([3, 14, 15, 26]), and pseudo-differential operators ([13, 16, 24, 25] and the references therein). Examples of convolution-dominated infinite matrices include the infinite matrix $(a(j - j'))_{j, j' \in \mathbb{Z}}$ associated with convolution operators and the infinite matrix $(a(j - j')e^{-2\pi\sqrt{-1}\theta j'(j-j')})_{i, j \in \mathbb{Z}}$ associated with twisted convolution operators, where $\theta \in \mathbb{R}$ and the sequence $a = (a(j))_{j \in \mathbb{Z}}$ satisfies $\sum_{j \in \mathbb{Z}} |a(j)| < \infty$ ([1, 14, 19, 27, 29]).

A convolution-dominated infinite matrix A is said to have ℓ^p -stability if there are two positive constants C_1 and C_2 such that

$$(1.4) \quad C_1 \|c\|_p \leq \|Ac\|_p \leq C_2 \|c\|_p \quad \text{for all } c \in \ell^p.$$

The ℓ^p -stability is one of basic assumptions for infinite matrices arising in the study of spline approximation, Gabor time-frequency analysis, nonuniform sampling, and algebra of pseudo-differential operators; see [1, 3, 6, 8, 9, 10, 14, 15, 16, 18, 19, 23, 24, 25, 26, 27, 29] and the references therein. **Practical criteria** for the ℓ^p -stability of a convolution-dominated infinite matrix will play important roles in the further study of those topics.

However, up to the knowledge of the author, little is known about practical criteria for the ℓ^p -stability of an infinite matrix. For an infinite matrix $A = (a(j - j'))_{j, j' \in \mathbb{Z}}$ associated with convolution operators, there is a very useful criterion for its ℓ^p -stability. It states that A has ℓ^p -stability if and only if the Fourier series $\hat{a}(\xi) := \sum_{j \in \mathbb{Z}} a(j)e^{-ij\xi}$ of the generating sequence $a = (a(j))_{j \in \mathbb{Z}} \in \ell^1$ does not vanish on the real line, i.e.,

$$(1.5) \quad \hat{a}(\xi) \neq 0 \quad \text{for all } \xi \in \mathbb{R}.$$

Applying this criterion for the ℓ^p -stability, one concludes that the spectrum $\sigma_p(C_a)$ of the convolution operator C_a as an operator on ℓ^p is independent of $1 \leq p \leq \infty$, i.e.,

$$(1.6) \quad \sigma_p(C_a) = \sigma_q(C_a) \quad \text{for all } 1 \leq p, q \leq \infty;$$

see [4, 17, 22, 23] and the references therein for the discussion on spectrum of various convolution operators. Applying the above criterion again, together with the classical Wiener's lemma ([29]), it follows that the inverse of an ℓ^p -stable convolution operator C_a is a convolution operator C_b associated with another summable sequence b .

For a convolution-dominated infinite matrix $A = (a(j, j'))_{j, j' \in \mathbb{Z}}$, a popular sufficient condition for its ℓ^1 -stability and ℓ^∞ -stability is that A is *diagonal-dominated*, i.e.,

$$(1.7) \quad \inf_{j \in \mathbb{Z}} \left(|a(j, j)| - \max \left(\sum_{j' \neq j} |a(j, j')|, \sum_{j' \neq j} |a(j', j)| \right) \right) > 0.$$

In this paper, we provide a practical criterion for the ℓ^p -stability of convolution-dominated infinite matrices. We show that a convolution-dominated infinite matrix A has ℓ^p -stability if and only if it has certain “diagonal-blocks-dominated” property (see Theorem 2.1 for the precise statement).

2. MAIN THEOREM

To state our criterion for the ℓ^p -stability of convolution-dominated infinite matrices, we introduce two concepts. Given an infinite matrix A , define the *truncation matrices* $A_s, s \geq 0$, by

$$A_s = (a(i, j)\chi_{(-s, s)}(i - j))_{i, j \in \mathbb{Z}},$$

where χ_E is the characteristic function on a set E . Given $y \in \mathbb{R}$ and $1 \leq N \in \mathbb{Z}$, define the operator χ_y^N on ℓ^p by

$$\chi_y^N : \ell^p \ni (c(j))_{j \in \mathbb{Z}} \mapsto (c(j)\chi_{(-N, N)}(j - y))_{j \in \mathbb{Z}} \in \ell^p.$$

The operator χ_y^N is a diagonal matrix $\text{diag}(\chi_{(-N, N)}(j - y))_{j \in \mathbb{Z}}$.

Theorem 2.1. *Let $1 \leq p \leq \infty$, and let A be a convolution-dominated infinite matrix in the class \mathcal{C} . Then the following statements are equivalent:*

- (i) *The infinite matrix A has ℓ^p -stability.*
- (ii) *There exist a positive constant C_0 and a positive integer N_0 such that*

$$(2.1) \quad \|\chi_n^{2N} A \chi_n^N c\|_p \geq C_0 \|\chi_n^N c\|_p, \quad c \in \ell^p,$$

hold for all integers $N \geq N_0$ and $n \in N\mathbb{Z}$.

- (iii) *There exist a positive integer N_0 and a positive constant α satisfying*

$$(2.2) \quad \alpha > 2(5 + 2^{1-p})^{1/p} \inf_{0 \leq s \leq N_0} \left(\|A - A_s\|_c + \frac{s}{N_0} \|A\|_c \right)$$

such that

$$(2.3) \quad \|\chi_n^{2N_0} A \chi_n^{N_0} c\|_p \geq \alpha \|\chi_n^{N_0} c\|_p, \quad c \in \ell^p,$$

hold for all $n \in N_0\mathbb{Z}$.

Taking $N_0 = 1$ in (2.2) and (2.3), we obtain a sufficient condition (2.4), which is a strong version of the diagonal-domination condition (1.7), for the ℓ^∞ -stability of a convolution-dominated infinite matrix.

Corollary 2.2. *Let $A = (a(j, j'))_{j, j' \in \mathbb{Z}}$ be a convolution-dominated infinite matrix in the class \mathcal{C} . If*

$$(2.4) \quad \inf_{j \in \mathbb{Z}} \left(|a(j, j)| - 2 \sum_{0 \neq k \in \mathbb{Z}} \sup_{j-j'=k} |a(j, j')| \right) > 0,$$

then A has ℓ^∞ -stability.

We say that an infinite matrix $A = (a(i, j))_{i, j \in \mathbb{Z}}$ is a *band matrix* if $a(i, j) = 0$ for all $i, j \in \mathbb{Z}$ satisfying $j > i + k$ or $j < i - k$. The quantity $2k + 1$ is the *bandwidth* of the matrix A . For a band matrix A with bandwidth $2k + 1$, $A - A_s$ is the zero matrix if $s > k$. Therefore for $N > k$,

$$\inf_{0 \leq s \leq N} \left(\|A - A_s\|_C + \frac{s}{N} \|A\|_C \right) \leq \frac{k}{N} \|A\|_C.$$

This, together with Theorem 2.1, gives the following sufficient condition for a band matrix to have ℓ^p -stability.

Corollary 2.3. *Let $1 \leq p \leq \infty$, and let A be a convolution-dominated band matrix in the class \mathcal{C} with bandwidth $2k + 1$. If there exists an integer $N_0 > k$ such that*

$$(2.5) \quad \|A\chi_n^{N_0} c\|_p \geq \alpha \|\chi_n^{N_0} c\|_p, \quad c \in \ell^p,$$

holds for some constant α strictly larger than $2(5 + 2^{1-p})^{1/p} k \|A\|_C / N_0$, then A has ℓ^p -stability.

If we further assume that the infinite matrix A in Corollary 2.3 has the form $A = (a(j - j'))_{j, j' \in \mathbb{Z}}$ for some finite sequence $a = (a(j))_{j \in \mathbb{Z}}$ satisfying $a(j) = 0$ for $|j| > k$, then $\|A\|_C = \sum_{|j| \leq k} |a(j)|$ and the condition (2.5) can reformulated as follows:

$$(2.6) \quad \|\tilde{A}_{N_0} c\|_p \geq \frac{\gamma k}{N_0} \left(\sum_{|j| \leq k} |a(j)| \right) \|c\|_p, \quad c \in \mathbb{R}^{2N_0+1},$$

holds for some $\gamma > 2(5 + 2^{1-p})^{1/p}$, where

$$(2.7) \quad \tilde{A}_{N_0} = (a(j - j'))_{-N_0-k \leq j \leq N_0+k, -N_0 \leq j' \leq N_0}$$

and

$$\|c\|_p = \begin{cases} (\sum_{j=-k_1}^{k_2} |c(j)|^p)^{1/p} & \text{if } 1 \leq p < \infty, \\ \sup_{-k_1 \leq j \leq k_2} |c(j)| & \text{if } p = \infty, \end{cases}$$

for $c = (c(-k_1), \dots, c(0), \dots, c(k_2))^T \in \mathbb{R}^{k_1+k_2+1}$. As a conclusion from (2.6) and (2.7), we see that if $A = (a(j - j'))_{j, j' \in \mathbb{Z}}$ does not have ℓ^p -stability, then for any large integer N ,

$$(2.8) \quad \inf_{0 \neq c \in \mathbb{R}^{2N+1}} \frac{\|\tilde{A}_N c\|_p}{\|c\|_p} \leq \frac{2(5 + 2^{1-p})^{1/p} k}{N} \left(\sum_{|j| \leq k} |a(j)| \right).$$

For the special case $p = 2$, the above inequality (2.8) can be interpreted as the minimal eigenvalue of $(\tilde{A}_N)^T \tilde{A}_N$ is less than or equal to $\frac{\sqrt{22}k^2}{N^2} (\sum_{|j| \leq k} |a(j)|)^2$, and it can also be rewritten as

$$(2.9) \quad \inf_{0 \neq P_N \in \Pi_N} \frac{\left(\int_{-\pi}^{\pi} |\hat{a}(\xi)|^2 |P_N(\xi)|^2 d\xi \right)^{1/2}}{\left(\int_{-\pi}^{\pi} |P_N(\xi)|^2 d\xi \right)^{1/2}} \leq \frac{\sqrt{22}k}{N} \left(\sum_{|j| \leq k} |a(j)| \right),$$

where $\hat{a}(\xi) = \sum_{j \in \mathbb{Z}} a(j) e^{-ij\xi}$ and Π_N is the set of all trigonometrical polynomials of degree at most N .

If the sequence $a = (a(j))_{j \in \mathbb{Z}}$ satisfies $a(0) = 1, a(-1) = -1$, and $a(j) = 0$ otherwise, then the bandwidth of the infinite matrix $A = (a(j - j'))_{j, j' \in \mathbb{Z}}$ is equal to 3, the norm $\|A\|_C$ of the associated infinite matrix A is equal to 2,

$$(2.10) \quad \tilde{A}_N = \begin{pmatrix} -1 & 0 & 0 & \cdots & 0 & 0 \\ 1 & -1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & -1 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & -1 \\ 0 & 0 & 0 & \cdots & 0 & 1 \end{pmatrix},$$

and

$$\inf_{0 \neq c \in \mathbb{R}^{2N+1}} \frac{\|\tilde{A}_N c\|_p}{\|c\|_p} \geq \frac{1}{N+1},$$

where the last inequality holds since the matrix

$$\tilde{B}_N := \begin{pmatrix} -1 & 0 & 0 & \cdots & 0 & 0 & 0 & 0 & \cdots & 0 & 0 \\ -1 & -1 & 0 & \cdots & 0 & 0 & 0 & 0 & \cdots & 0 & 0 \\ -1 & -1 & -1 & \cdots & 0 & 0 & 0 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ -1 & -1 & -1 & \cdots & -1 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 0 & \cdots & 0 & 0 & 1 & 1 & \cdots & 1 & 1 \\ 0 & 0 & 0 & \cdots & 0 & 0 & 0 & 1 & \cdots & 1 & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 0 & 0 & 0 & \cdots & 1 & 1 \\ 0 & 0 & 0 & \cdots & 0 & 0 & 0 & 0 & \cdots & 0 & 1 \end{pmatrix}$$

is a left inverse of the matrix \tilde{A}_N . Therefore the order N^{-1} in (2.8) cannot be improved in general, but the author believes that the bound constant $2(5+2^{1-p})^{1/p}$ in (2.2) and (2.8) is not optimal and could be improved.

3. PROOF

We say that a discrete subset Λ of \mathbb{R}^d is *relatively-separated* if

$$(3.1) \quad R(\Lambda) := \sup_{x \in \mathbb{R}^d} \sum_{\lambda \in \Lambda} \chi_{\lambda + [-1/2, 1/2]^d}(x) < \infty$$

([1, 23, 27]). Clearly, the set \mathbb{Z} of all integers is a relatively-separated subset of \mathbb{R} with

$$(3.2) \quad R(\mathbb{Z}) = 1.$$

Given a discrete set Λ , let $\ell^p(\Lambda)$ be the set of all p -summable sequences on the set Λ with standard norm $\|\cdot\|_{\ell^p(\Lambda)}$ or $\|\cdot\|_p$ for brevity.

Given two relatively-separated subsets Λ and Λ' of \mathbb{R}^d , define

$$\mathcal{C}(\Lambda, \Lambda') = \left\{ A := (a(\lambda, \lambda'))_{\lambda \in \Lambda, \lambda' \in \Lambda'} \mid \|A\|_{\mathcal{C}(\Lambda, \Lambda')} < \infty \right\},$$

where

$$\|A\|_{\mathcal{C}(\Lambda, \Lambda')} = \sum_{k \in \mathbb{Z}^d} \sup_{\lambda \in \Lambda, \lambda' \in \Lambda'} |a(\lambda, \lambda')| \chi_{k + [-1/2, 1/2]^d}(\lambda - \lambda').$$

It is obvious that

$$(3.3) \quad \mathcal{C}(\mathbb{Z}, \mathbb{Z}) = \mathcal{C}.$$

Given an infinite matrix $A = (a(\lambda, \lambda'))_{\lambda \in \Lambda, \lambda' \in \Lambda'}$, define its *truncation matrices* $A_s, s \geq 0$, by

$$A_s = \left(a(\lambda, \lambda') \chi_{(-s,s)^d}(\lambda - \lambda') \right)_{\lambda \in \Lambda, \lambda' \in \Lambda'}.$$

For any $y \in \mathbb{R}^d$ and a positive integer N , define the operator χ_y^N on $\ell^p(\Lambda)$ by

$$(3.4) \quad \chi_n^N : \ell^p(\Lambda) \ni (c(\lambda))_{\lambda \in \Lambda} \mapsto (c(\lambda) \chi_{(-N,N)^d}(\lambda - y))_{\lambda \in \Lambda} \in \ell^p(\Lambda).$$

In this section, we establish the following criterion for the ℓ^p -stability of infinite matrices in the class $\mathcal{C}(\Lambda, \Lambda')$, which is a slight generalization of Theorem 2.1 by (3.2) and (3.3).

Theorem 3.1. *Let $1 \leq p \leq \infty$, the subsets Λ, Λ' of \mathbb{R}^d be relatively-separated, and the infinite matrix A belong to $\mathcal{C}(\Lambda, \Lambda')$. Then the following statements are equivalent to each other:*

- (i) *The infinite matrix A has ℓ^p -stability, i.e., there exist positive constants C_1 and C_2 such that*

$$(3.5) \quad C_1 \|c\|_{\ell^p(\Lambda')} \leq \|Ac\|_{\ell^p(\Lambda)} \leq C_2 \|c\|_{\ell^p(\Lambda')} \quad \text{for all } c \in \ell^p(\Lambda').$$

- (ii) *There exist a positive constant C_0 and a positive integer N_0 such that*

$$(3.6) \quad \|\chi_n^{2N} A \chi_n^N c\|_{\ell^p(\Lambda)} \geq C_0 \|\chi_n^N c\|_{\ell^p(\Lambda')} \quad \text{for all } c \in \ell^p(\Lambda'),$$

where $N_0 \leq N \in \mathbb{Z}$ and $n \in N\mathbb{Z}^d$.

- (iii) *There exist a positive integer N_0 and a positive constant α satisfying*

$$(3.7) \quad \alpha > 2(5 + 2^{1-p})^{d/p} R(\Lambda)^{1/p} R(\Lambda')^{1-1/p} \inf_{0 \leq s \leq N_0} \left(\|A - A_s\|_{\mathcal{C}(\Lambda, \Lambda')} + \frac{ds}{N_0} \|A\|_{\mathcal{C}(\Lambda, \Lambda')} \right)$$

such that

$$(3.8) \quad \|\chi_n^{2N_0} A \chi_n^{N_0} c\|_{\ell^p(\Lambda)} \geq \alpha \|\chi_n^{N_0} c\|_{\ell^p(\Lambda')}$$

hold for all $c \in \ell^p(\Lambda')$ and $n \in N_0\mathbb{Z}$.

Using the above theorem, we obtain the following equivalence of ℓ^p -stability for infinite matrices having certain off-diagonal decay, which is established in [2, 28, 23] for $\gamma > d(d + 1), \gamma > 0$, and $\gamma \geq 0$ respectively.

Corollary 3.2. *Let Λ, Λ' be relatively-separated subsets of \mathbb{R}^d , and let $A = (a(\lambda, \lambda'))_{\lambda \in \Lambda, \lambda' \in \Lambda'}$ satisfy*

$$\|A\|_{\mathcal{C}_\gamma(\Lambda, \Lambda')} = \sum_{k \in \mathbb{Z}^d} (1 + |k|)^\gamma \sup_{\lambda \in \Lambda, \lambda' \in \Lambda'} |a(\lambda, \lambda')| \chi_{k+[-1/2, 1/2]^d}(\lambda - \lambda') < \infty,$$

where $\gamma > 0$. Then the ℓ^p -stability of the infinite matrix A are equivalent to each other for different $1 \leq p \leq \infty$.

Proof. Let $1 \leq p \leq \infty$ and let A have ℓ^p -stability. Then by Theorem 3.1 there exists a positive constant C_0 and a positive integer N_0 such that

$$(3.9) \quad \|\chi_n^{2N} A \chi_n^N c\|_{\ell^p(\Lambda)} \geq C_0 \|\chi_n^N c\|_{\ell^p(\Lambda')} \quad \text{for all } c \in \ell^p(\Lambda'),$$

where $N_0 \leq N \in \mathbb{Z}$ and $n \in N\mathbb{Z}^d$. From the equivalence of different norms on a finite-dimensional space, we have that

$$\begin{aligned} & ((2N)^d R(\Lambda))^{\min(1/q-1/p, 0)} \|\chi_n^N c\|_{\ell^p(\Lambda)} \leq \|\chi_n^N c\|_{\ell^q(\Lambda)} \\ & \leq ((2N)^d R(\Lambda))^{\max(1/q-1/p, 0)} \|\chi_n^N c\|_{\ell^p(\Lambda)} \quad \text{for all } c \in \ell^p(\Lambda), \end{aligned}$$

where $1 \leq p, q \leq \infty, 1 \leq N \in \mathbb{Z}$ and $n \in N\mathbb{Z}^d$ ([2, 23]). Therefore for $1 \leq q \leq \infty$,

$$(3.10) \quad \begin{aligned} \|\chi_n^{2N} A \chi_n^N c\|_{\ell^q(\Lambda)} &\geq C_0(2N)^{-d|1/p-1/q|} R(\Lambda')^{\min(1/p-1/q, 0)} \\ &\times R(\Lambda)^{-\max(1/p-1/q, 0)} \|\chi_n^N c\|_{\ell^q(\Lambda')} \quad \text{for all } c \in \ell^q(\Lambda'), \end{aligned}$$

where $N_0 \leq N \in \mathbb{Z}$ and $n \in N\mathbb{Z}^d$. We notice that

$$(3.11) \quad \begin{aligned} \inf_{0 \leq s \leq N} (\|A - A_s\|_{\mathcal{C}(\Lambda, \Lambda')} + \frac{ds}{N} \|A\|_{\mathcal{C}(\Lambda, \Lambda')}) &\leq \|A\|_{\mathcal{C}_\gamma(\Lambda, \Lambda')} \inf_{0 \leq s \leq N} (s^\gamma + \frac{ds}{N}) \\ &\leq (d+1) \|A\|_{\mathcal{C}_\gamma(\Lambda, \Lambda')} N^{-\gamma/(1+\gamma)}. \end{aligned}$$

Thus for $1 \leq q \leq \infty$ with $d|1/p - 1/q| < \gamma/(1 + \gamma)$, it follows from (3.10) and (3.11) that there exists a sufficiently large integer N_0 such that

$$(3.12) \quad \|\chi_n^{2N} A \chi_n^N c\|_{\ell^q(\Lambda)} \geq \alpha \|\chi_n^N c\|_{\ell^q(\Lambda')}$$

hold for all $c \in \ell^q(\Lambda'), N \geq N_0$ and $n \in N\mathbb{Z}^d$, where α is a positive constant larger than $2(5 + 2^{1-q})^{d/q} R(\Lambda)^{1/q} R(\Lambda')^{1-1/q} \inf_{0 \leq s \leq N_0} (\|A - A_s\|_{\mathcal{C}(\Lambda, \Lambda')} + \frac{ds}{N_0} \|A\|_{\mathcal{C}(\Lambda, \Lambda')})$. Then by Theorem 3.1, the infinite matrix A has ℓ^q -stability for all $1 \leq q \leq \infty$ with $d|1/q - 1/p| < \gamma/(1 + \gamma)$. Applying the above trick repeatedly, we prove the ℓ^q -stability of the infinite matrix A for any $1 \leq q \leq \infty$. \square

To prove Theorem 3.1, we first recall some basic properties for infinite matrices A in the class $\mathcal{C}(\Lambda, \Lambda')$ and its truncation matrices $A_s, s \geq 0$.

Lemma 3.3 ([23]). *Let $1 \leq p \leq \infty$, the subsets Λ, Λ' of \mathbb{R}^d be relatively-separated, A be an infinite matrix in the class $\mathcal{C}(\Lambda, \Lambda')$, and $A_s, s \geq 0$, be the truncation matrices of A . Then*

$$(3.13) \quad \|Ac\|_{\ell^p(\Lambda)} \leq R(\Lambda)^{1/p} R(\Lambda')^{1-1/p} \|A\|_{\mathcal{C}(\Lambda, \Lambda')} \|c\|_{\ell^p(\Lambda')} \quad \text{for all } c \in \ell^p(\Lambda'),$$

$$(3.14) \quad \lim_{s \rightarrow +\infty} \|A - A_s\|_{\mathcal{C}(\Lambda, \Lambda')} = 0,$$

$$(3.15) \quad \lim_{N \rightarrow +\infty} \inf_{0 \leq s \leq N} (\|A - A_s\|_{\mathcal{C}(\Lambda, \Lambda')} + \frac{ds}{N} \|A\|_{\mathcal{C}(\Lambda, \Lambda')}) = 0,$$

and

$$(3.16) \quad \|A_s\|_{\mathcal{C}} \leq \|A\|_{\mathcal{C}} \quad \text{for all } s \geq 0.$$

Let $\psi_0(x_1, \dots, x_d) = \prod_{i=1}^d \max(\min(2 - 2|x_i|, 1), 0)$ be a cut-off function on \mathbb{R}^d . Then

$$(3.17) \quad 0 \leq \chi_{[-1/2, 1/2]^d}(x) \leq \psi_0(x) \leq \chi_{(-1, 1)^d}(x) \leq 1 \quad \text{for all } x \in \mathbb{R}^d,$$

and

$$(3.18) \quad |\psi_0(x) - \psi_0(y)| \leq 2d \|x - y\|_\infty \quad \text{for all } x, y \in \mathbb{R}^d,$$

where $\|x\|_\infty = \max_{1 \leq i \leq d} |x_i|$ for $x = (x_1, \dots, x_d)$. Define the multiplication operator Ψ_n^N on $\ell^p(\Lambda)$ by

$$(3.19) \quad \Psi_n^N : \ell^p(\Lambda) \ni (c(\lambda))_{\lambda \in \Lambda} \mapsto \left(\psi_0\left(\frac{\lambda - n}{N}\right) c(\lambda) \right)_{\lambda \in \Lambda} \in \ell^p(\Lambda).$$

Applying (3.17) and (3.18) for the cut-off function ψ_0 , we obtain the following properties for the multiplication operators $\Psi_n^N, n \in N\mathbb{Z}$.

Lemma 3.4. *Let $1 \leq N \in \mathbb{Z}$, Λ be a relatively-separated subset of \mathbb{R}^d , and the multiplication operators $\Psi_n^N, n \in N\mathbb{Z}^d$, be as in (3.19). Then*

$$(3.20) \quad \|\Psi_n^N c\|_{\ell^p(\Lambda)} \leq \|\chi_n^N c\|_{\ell^p(\Lambda)} \quad \text{for all } c \in \ell^p(\Lambda),$$

where $1 \leq p \leq \infty$,

$$(3.21) \quad \|c\|_{\ell^p(\Lambda)} \leq \left(\sum_{n \in N\mathbb{Z}^d} \|\Psi_n^N c\|_{\ell^p(\Lambda)}^p \right)^{1/p} \leq 2^{d/p} \|c\|_{\ell^p(\Lambda)} \quad \text{for all } c \in \ell^p(\Lambda),$$

(3.22)

$$4^{d/p} \|c\|_{\ell^p(\Lambda)} \leq \left(\sum_{n \in N\mathbb{Z}^d} \|\Psi_n^{4N} c\|_{\ell^p(\Lambda)}^p \right)^{1/p} \leq (5 + 2^{1-p})^{d/p} \|c\|_{\ell^p(\Lambda)} \quad \text{for all } c \in \ell^p(\Lambda),$$

where $1 \leq p < \infty$, and

$$(3.23) \quad \|c\|_{\ell^\infty(\Lambda)} = \sup_{n \in N\mathbb{Z}^d} \|\Psi_n^N c\|_{\ell^\infty(\Lambda)} = \sup_{n \in N\mathbb{Z}^d} \|\Psi_n^{4N} c\|_{\ell^\infty(\Lambda)} \quad \text{for all } c \in \ell^\infty(\Lambda).$$

To prove Theorem 2.1, we also need the following result.

Lemma 3.5 ([23]). *Let $N \geq 1$, the subsets Λ, Λ' of \mathbb{R}^d be relatively-separated, A be an infinite matrix in the class $\mathcal{C}(\Lambda, \Lambda')$, A_N be the truncation matrix of A , and $\Psi_n^N, n \in N\mathbb{Z}^d$, be the multiplication operators in (3.19). Then*

$$(3.24) \quad \|\Psi_n^N A_N - A_N \Psi_n^N\|_{\mathcal{C}(\Lambda, \Lambda')} \leq \inf_{0 \leq s \leq N} \left(\|A_N - A_s\|_{\mathcal{C}(\Lambda, \Lambda')} + \frac{2ds}{N} \|A_s\|_{\mathcal{C}(\Lambda, \Lambda')} \right).$$

Now we start to prove Theorem 3.1.

Proof of Theorem 3.1. (i) \implies (ii): By the ℓ^p -stability of the infinite matrix A , there exists a positive constant C_0 (independent of $n \in N\mathbb{Z}^d$ and $1 \leq N \in \mathbb{Z}$) such that

$$(3.25) \quad \|A\chi_n^N c\|_{\ell^p(\Lambda)} \geq C_0 \|\chi_n^N c\|_{\ell^p(\Lambda')} \quad \text{for all } c \in \ell^p(\Lambda'),$$

where $n \in N\mathbb{Z}^d$ and $N \geq 1$. Noting that

$$(3.26) \quad \chi_n^{2N} A_N \psi_n^N = A_N \psi_n^N$$

and applying (3.13) yield

$$(3.27) \quad \begin{aligned} & \|A\chi_n^N c - \chi_n^{2N} A\chi_n^N c\|_{\ell^p(\Lambda)} \\ &= \|(I - \chi_n^{2N})(A - A_N)\chi_n^N c\|_{\ell^p(\Lambda)} \\ &\leq R(\Lambda)^{1/p} R(\Lambda')^{1-1/p} \|A - A_N\|_{\mathcal{C}(\Lambda, \Lambda')} \|\chi_n^N c\|_{\ell^p(\Lambda')}, \end{aligned}$$

where I is the identity operator. Combining the estimates in (3.25) and (3.27) proves that

$$(3.28) \quad \|\chi_n^{2N} A\chi_n^N c\|_{\ell^p(\Lambda)} \geq (C_0 - R(\Lambda)^{1/p} R(\Lambda')^{1-1/p} \|A - A_N\|_{\mathcal{C}(\Lambda, \Lambda')}) \|\chi_n^N c\|_{\ell^p(\Lambda')}$$

hold for all $c \in \ell^p(\Lambda')$, where $n \in N\mathbb{Z}^d$ and $N \geq 1$. The conclusion (ii) then follows from (3.14) and (3.28).

(ii) \implies (iii): The implication follows from (3.15).

(iii) \implies (i): Let $1 \leq p < \infty$. Take any $n \in N_0\mathbb{Z}^d$ and $c \in \ell^p(\Lambda')$. By the assumption (iii) for the infinite matrix A ,

$$(3.29) \quad \|\chi_n^{2N_0} A\Psi_n^{N_0} c\|_{\ell^p(\Lambda)} = \|\chi_n^{2N_0} A\chi_n^{N_0} \Psi_n^{N_0} c\|_{\ell^p(\Lambda)} \geq \alpha \|\Psi_n^{N_0} c\|_{\ell^p(\Lambda')}.$$

This together with (3.13) and (3.26) implies that

$$\begin{aligned}
 & \|A_{N_0} \Psi_n^{N_0} c\|_{\ell^p(\Lambda)} \\
 &= \|\chi_n^{2N_0} (A_{N_0} - A + A) \Psi_n^{N_0} c\|_{\ell^p(\Lambda)} \\
 &\geq \|\chi_n^{2N_0} A \chi_n^{N_0} \Psi_n^{N_0} c\|_{\ell^p(\Lambda)} - \|\chi_n^{2N_0} (A_{N_0} - A) \Psi_n^{N_0} c\|_{\ell^p(\Lambda)} \\
 (3.30) \quad &\geq (\alpha - R(\Lambda)^{1/p} R(\Lambda')^{1-1/p} \|A - A_{N_0}\|_{\mathcal{C}(\Lambda, \Lambda')}) \|\Psi_n^{N_0} c\|_{\ell^p(\Lambda')}.
 \end{aligned}$$

From (3.13) and (3.24) it follows that

$$\begin{aligned}
 & \|(\Psi_n^{N_0} A_{N_0} - A_{N_0} \Psi_n^{N_0}) c\|_{\ell^p(\Lambda)} \\
 &= \|(\Psi_n^{N_0} A_{N_0} - A_{N_0} \Psi_n^{N_0}) \Psi_n^{4N_0} c\|_{\ell^p(\Lambda)} \\
 &\leq R(\Lambda)^{1/p} R(\Lambda')^{1-1/p} \|\Psi_n^{N_0} A_{N_0} - A_{N_0} \Psi_n^{N_0}\|_{\mathcal{C}(\Lambda, \Lambda')} \|\Psi_n^{4N_0} c\|_{\ell^p(\Lambda')} \\
 &\leq R(\Lambda)^{1/p} R(\Lambda')^{1-1/p} \\
 (3.31) \quad &\times \inf_{0 \leq s \leq N_0} \left(\|A_{N_0} - A_s\|_{\mathcal{C}} + \frac{2ds}{N_0} \|A_{N_0}\|_{\mathcal{C}} \right) \|\Psi_n^{4N_0} c\|_{\ell^p(\Lambda')}.
 \end{aligned}$$

Combining (3.21), (3.22), (3.30) and (3.31), we get

$$\begin{aligned}
 2^{d/p} \|A_{N_0} c\|_{\ell^p(\Lambda)} &\geq \left(\sum_{n \in N_0 \mathbb{Z}} \|\Psi_n^{N_0} A_{N_0} c\|_{\ell^p(\Lambda)}^p \right)^{1/p} \\
 &\geq \left(\alpha - R(\Lambda)^{1/p} R(\Lambda')^{1-1/p} \|A - A_{N_0}\|_{\mathcal{C}(\Lambda, \Lambda')} \right) \left(\sum_{n \in N_0 \mathbb{Z}} \|\Psi_n^{N_0} c\|_{\ell^p(\Lambda')}^p \right)^{1/p} \\
 &\quad - R(\Lambda)^{1/p} R(\Lambda')^{1-1/p} \inf_{0 \leq s \leq N_0} \left(\|A_{N_0} - A_s\|_{\mathcal{C}(\Lambda, \Lambda')} + \frac{2ds}{N_0} \|A_{N_0}\|_{\mathcal{C}(\Lambda, \Lambda')} \right) \\
 &\quad \times \left(\sum_{n \in N_0 \mathbb{Z}} \|\Psi_n^{4N_0} c\|_{\ell^p(\Lambda')}^p \right)^{1/p} \\
 &\geq \left(\alpha - R(\Lambda)^{1/p} R(\Lambda')^{1-1/p} \|A - A_{N_0}\|_{\mathcal{C}(\Lambda, \Lambda')} - (5 + 2^{1-p})^{1/p} R(\Lambda)^{1/p} R(\Lambda')^{1-1/p} \right. \\
 &\quad \left. \times \inf_{0 \leq s \leq N_0} \left(\|A_{N_0} - A_s\|_{\mathcal{C}(\Lambda, \Lambda')} + \frac{2ds}{N_0} \|A_{N_0}\|_{\mathcal{C}(\Lambda, \Lambda')} \right) \right) \|c\|_{\ell^p(\Lambda')}.
 \end{aligned}$$

Therefore

$$\begin{aligned}
 \|Ac\|_{\ell^p(\Lambda)} &\geq \|A_{N_0} c\|_{\ell^p(\Lambda)} - \|(A - A_{N_0})c\|_{\ell^p(\Lambda)} \\
 &\geq 2^{-1/p} \left(\alpha - (1 + 2^{d/p}) R(\Lambda)^{1/p} R(\Lambda')^{1-1/p} \|A - A_{N_0}\|_{\mathcal{C}(\Lambda, \Lambda')} \right. \\
 &\quad \left. - (5 + 2^{1-p})^{d/p} R(\Lambda)^{1/p} R(\Lambda')^{1-1/p} \right. \\
 &\quad \left. \times \inf_{0 \leq s \leq N_0} \left(\|A_{N_0} - A_s\|_{\mathcal{C}(\Lambda, \Lambda')} + \frac{2ds}{N_0} \|A_{N_0}\|_{\mathcal{C}(\Lambda, \Lambda')} \right) \right) \|c\|_{\ell^p(\Lambda')} \\
 &\geq 2^{-d/p} \left(\alpha - 2(5 + 2^{1-p})^{1/p} R(\Lambda)^{1/p} \right. \\
 &\quad \left. \times R(\Lambda')^{1-1/p} \inf_{0 \leq s \leq N_0} \left(\|A - A_s\|_{\mathcal{C}(\Lambda, \Lambda')} + \frac{ds}{N_0} \|A\|_{\mathcal{C}(\Lambda, \Lambda')} \right) \right) \|c\|_{\ell^p(\Lambda')},
 \end{aligned}$$

and the conclusion (i) for $1 \leq p < \infty$ follows.

The conclusion (i) for $p = \infty$ can be proved by a similar argument. We omit the details here. \square

ACKNOWLEDGMENTS

The author thanks Professors Deguang Han, Zuhair M. Nashed, Xianliang Shi, and Wai-Shing Tang for their discussion and suggestions in preparing the manuscript.

REFERENCES

- [1] A. Aldroubi and K. Gröchenig, Nonuniform sampling and reconstruction in shift-invariant space, *SIAM Review*, **43**(2001), 585–620. MR1882684 (2003e:94040)
- [2] A. Aldroubi, A. Baskakov and I. Krishtal, Slanted matrices, Banach frames, and sampling, *J. Funct. Anal.*, **255**(2008), 1667–1691. MR2442078 (2010a:46059)
- [3] R. Balan, P. G. Casazza, C. Heil, and Z. Landau, Density, overcompleteness and localization of frames. I. Theory; II. Gabor systems, *J. Fourier Anal. Appl.*, **12**(2006), 105–143; 309–344. MR2224392 (2007b:42041); MR2235170 (2007b:42042)
- [4] B. A. Barnes, When is the spectrum of a convolution operator on L^p independent of p ? *Proc. Edinburgh Math. Soc.*, **33**(1990), 327–332. MR1057760 (91f:47046)
- [5] A. G. Baskakov, Wiener’s theorem and asymptotic estimates for elements of inverse matrices, *Funktsional. Anal. i Prilozhen.*, **24**(1990), 64–65; translation in *Funct. Anal. Appl.*, **24**(1990), 222–224. MR1082033 (92g:47049)
- [6] C. K. Chui, W. He, and J. Stöckler, Nonstationary tight wavelet frames. I. Bounded intervals; II. Unbounded intervals, *Appl. Comp. Harmonic Anal.*, **17**(2004), 141–197; **18**(2005), 25–66. MR2082157 (2005f:42082); MR2110512 (2005j:42026)
- [7] A. Connes, C^* algèbres et géométrie différentielle. *C. R. Acad. Sci. Paris Sér. A-B*, **290**(1980), A599–A604. MR572645 (81c:46053)
- [8] C. de Boor, A bound on the L_∞ -norm of the L_2 -approximation by splines in terms of a global mesh ratio, *Math. Comp.*, **30**(1976), 687–694. MR0425432 (54:13387)
- [9] S. Demko, Inverse of band matrices and local convergences of spline projections, *SIAM J. Numer. Anal.*, **14**(1977), 616–619. MR0455281 (56:13520)
- [10] G. Fendler, K. Gröchenig, and M. Leinert, Convolution-dominated operators on discrete groups, *Integral Equations Operator Theory*, **61**(2008), 493–509. MR2434338 (2009m:47085)
- [11] I. M. Gelfand, D. A. Raikov, and G. E. Silov, *Commutative Normed Rings*, Chelsea, New York, 1964. MR0205105 (34:4940)
- [12] I. Gohberg, M. A. Kaashoek, and H. J. Woerdeman, The band method for positive and strictly contractive extension problems: An alternative version and new applications, *Integral Equations Operator Theory*, **12**(1989), 343–382. MR998278 (90c:47022)
- [13] K. Gröchenig, Time-frequency analysis of Sjöstrand’s class, *Rev. Mat. Iberoamer.*, **22**(2006), 703–724. MR2294795 (2008b:35308)
- [14] K. Gröchenig and M. Leinert, Wiener’s lemma for twisted convolution and Gabor frames, *J. Amer. Math. Soc.*, **17**(2003), 1–18. MR2015328 (2004m:42037)
- [15] K. Gröchenig and M. Leinert, Symmetry and inverse-closedness of matrix algebras and functional calculus for infinite matrices, *Trans. Amer. Math. Soc.*, **358**(2006), 2695–2711. MR2204052 (2006k:47065)
- [16] K. Gröchenig and T. Strohmer, Pseudodifferential operators on locally compact abelian groups and Sjöstrand’s symbol class, *Journal für die reine und angewandte Mathematik*, **613**(2007), 121–146. MR2377132 (2009d:35360)
- [17] A. Hulanicki, On the spectrum of convolution operators on groups with polynomial growth, *Invent. Math.*, **17**(1972), 135–142. MR0323951 (48:2304)
- [18] S. Jaffard, Propriétés des matrices bien localisées près de leur diagonale et quelques applications, *Ann. Inst. Henri Poincaré*, **7**(1990), 461–476. MR1138533 (93h:47035)
- [19] R.-Q. Jia and C. A. Micchelli, Using the refinement equations for the construction of pre-wavelets. II. Powers of two, in *Curves and Surfaces (Chamonix-Mont-Blanc, 1990)*, Academic Press, Boston, MA, 1991, 209–246. MR1123739 (93e:65024)
- [20] V. G. Kurbatov, Algebras of difference and integral operators, *Funktsional. Anal. i Prilozhen.*, **24**(1990), 87–88. MR1069415 (91k:47067)
- [21] M. A. Naimark, *Normed Algebras*, Wolters-Noordhoff Publishing, Groningen, 1972. MR0438123 (55:11042)

- [22] T. Pytlik, On the spectral radius of elements in group algebras, *Bull. Acad. Polon. Sci. Ser. Sci. Math.*, **21**(1973), 899–902. MR0328476 (48:6818)
- [23] C. E. Shin and Q. Sun, Stability of localized operators, *J. Funct. Anal.*, **256**(2009), 2417–2439. MR2502521 (2010e:47012)
- [24] J. Sjöstrand, An algebra of pseudodifferential operators, *Math. Res. Lett.*, **1**(1994), 185–192. MR1266757 (95b:47065)
- [25] J. Sjöstrand, Wiener type algebras of pseudodifferential operators, Centre de Mathématiques, École Polytechnique, Palaiseau, France, Séminaire 1994–1995, 1995. MR1362552 (96j:47049)
- [26] Q. Sun, Non-uniform sampling and reconstruction for signals with finite rate of innovations, *SIAM J. Math. Anal.*, **38**(2006), 1389–1422. MR2286012 (2008b:94049)
- [27] Q. Sun, Wiener’s lemma for infinite matrices, *Trans. Amer. Math. Soc.*, **359**(2007), 3099–3123. MR2299448 (2008e:47036)
- [28] R. Tessera, Finding left inverse for operators on $\ell^p(\mathbb{Z}^d)$ with polynomial decay, preprint, 2008. arXiv:0801.1532
- [29] N. Wiener, Tauberian theorems, *Ann. of Math. (2)*, **33**(1932), 1–100. MR1503035

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CENTRAL FLORIDA, ORLANDO, FLORIDA 32816
E-mail address: qsun@mail.ucf.edu