

A STRENGTHENING AND A MULTIPARTITE GENERALIZATION OF THE ALON-BOPPANA-SERRE THEOREM

BOJAN MOHAR

(Communicated by Jim Haglund)

ABSTRACT. The Alon-Boppana theorem confirms that for every $\varepsilon > 0$ and every integer $d \geq 3$, there are only finitely many d -regular graphs whose second largest eigenvalue is at most $2\sqrt{d-1} - \varepsilon$. Serre gave a strengthening showing that a positive proportion of eigenvalues of any d -regular graph must be bigger than $2\sqrt{d-1} - \varepsilon$. We provide a multipartite version of this result. Our proofs are elementary and also work in the case when graphs are not regular. In the simplest, monopartite case, our result extends the Alon-Boppana-Serre result to non-regular graphs of minimum degree d and bounded maximum degree. The two-partite result shows that for every $\varepsilon > 0$ and any positive integers d_1, d_2, d , every n -vertex graph of maximum degree at most d , whose vertex set is the union of (not necessarily disjoint) subsets V_1, V_2 , such that every vertex in V_i has at least d_i neighbors in V_{3-i} for $i = 1, 2$, has $\Omega_\varepsilon(n)$ eigenvalues that are larger than $\sqrt{d_1-1} + \sqrt{d_2-1} - \varepsilon$. Finally, we strengthen the Alon-Boppana-Serre theorem by showing that the lower bound $2\sqrt{d-1} - \varepsilon$ can be replaced by $2\sqrt{d-1} + \delta$ for some $\delta > 0$ if graphs have bounded “global girth”. On the other side of the spectrum, if the odd girth is large, then we get an Alon-Boppana-Serre type theorem for the negative eigenvalues as well.

1. INTRODUCTION

After the breakthrough paper by Alon and Milman [2] in 1985, it became apparent that regular graphs, whose *spectral gap* (i.e. the difference between the largest and the second largest eigenvalue) is large, possess some extraordinary properties, like unusually fast expansion and resemblance to random graphs. This led to the definition of *Ramanujan graphs*. These are d -regular graphs whose second largest eigenvalue does not exceed the value $2\sqrt{d-1}$. Lubotzky, Phillips, and Sarnak [16], and independently Margulis [17], were the first to show that Ramanujan graphs exist. Their constructions are based on number theory and work when the degree d is equal to $p+1$ for some prime p . Later, several new constructions were discovered,

Received by the editors February 4, 2010.

2010 *Mathematics Subject Classification*. Primary 05C50.

Key words and phrases. Spectral radius, eigenvalue, Ramanujan graph, universal cover.

The author was supported in part by the Research Grant P1-0297 of ARRS (Slovenia), by an NSERC Discovery Grant (Canada) and by the Canada Research Chair program.

The author is on leave from IMFM & FMF, Department of Mathematics, University of Ljubljana, Ljubljana, Slovenia.

showing that Ramanujan graphs exist for every degree $d \geq 3$ which is of the form $p^k + 1$ for some integer $k \geq 1$ and some prime p ; see Morgenstern [19].

It is not immediately clear why the special choice of $2\sqrt{d-1}$ is taken when defining Ramanujan graphs. One reason is that this is the spectral radius of the infinite d -regular tree, which is the universal cover for all d -regular graphs. Another reason is the following result of Alon and Boppana (see [1]) which shows that this is the smallest number that makes sense.

Theorem 1.1 (Alon-Boppana). *For every $d \geq 2$ and every $\varepsilon > 0$, there are only finitely many d -regular graphs whose second largest eigenvalue is at most $2\sqrt{d-1} - \varepsilon$.*

Alternative proofs of Theorem 1.1 were given by Friedman [8] and by Nilli [20], who has recently further simplified her arguments in [21]. Actually, the proofs in [8, 21] imply a stronger version of Theorem 1.1 by making the same conclusion for more eigenvalues than just the second largest one. This strengthening is attributed to Serre [23] (see also [5, 7, 12]), who stated the following quantitative version of the Alon-Boppana theorem:

Theorem 1.2 (Serre). *For every positive integer d and every $\varepsilon > 0$, there exists a constant $c = c(d, \varepsilon)$ such that every d -regular graph of order n has at least cn eigenvalues that are larger than $2\sqrt{d-1} - \varepsilon$.*

In this paper we give a multipartite generalization of the Alon-Boppana theorem; see Theorem 5.1. The Ramanujan value $2\sqrt{d-1}$ is replaced by the spectral radius of the universal covering tree of the multipartite parameters (cf. Section 2 for definitions). Our proof has similarities with Nilli's proof [21], and it seems to be even simpler if restricted to the special case of d -regular graphs. The main step is based on the interlacing theorem and is entirely elementary.

Our proofs work also in the case when graphs are not regular. In the simplest, monpartite case, our result extends the Alon-Boppana-Serre result to non-regular graphs of minimum degree d and bounded maximum degree. A strengthening of this form has been obtained previously by Hoory [11]. In the next simplest two-partite case it is shown that for every $\varepsilon > 0$ and any positive integers d_1, d_2, d , every n -vertex graph of maximum degree at most d , whose vertex set is the union of (not necessarily disjoint) subsets V_1, V_2 , such that every vertex in V_i has at least d_i neighbors in V_{3-i} for $i = 1, 2$, has $\Omega_\varepsilon(n)$ eigenvalues that are larger than $\sqrt{d_1-1} + \sqrt{d_2-1} - \varepsilon$.

After submission of this paper, S. Cioabă informed us about some related work. Greenberg [10] obtained a generalized version of the Serre theorem in a similar form as ours, but only claiming that there are eigenvalues whose absolute value is larger than $2\sqrt{d-1} - \varepsilon$. Cioabă [4] improved Greenberg's work to the same form as given in Theorem 1.2. Greenberg's result also appears in [15].

In the last section we tailor the proofs to obtain a strengthening of the Alon-Boppana-Serre theorem by showing that the lower bound $2\sqrt{d-1} - \varepsilon$ can be replaced by $2\sqrt{d-1} + \delta$ for some $\delta > 0$ if graphs have bounded universal girth (see Section 6 for the definition). On the other side of the spectrum, if the *odd girth* (i.e. the length of a shortest odd cycle) is large, then we obtain an Alon-Boppana-Serre type theorem for the negative eigenvalues.

If G is a (finite) graph, we denote by $\lambda_i = \lambda_i(G)$ the i th largest eigenvalue of the adjacency matrix $A(G)$ of G , respecting multiplicities. The largest eigenvalue

of G , $\rho(G) = \lambda_1(G)$, is also referred to as the *spectral radius* of G . It follows from the Perron-Frobenius theorem (see, e.g. [13]) that $\rho(G)$ is an eigenvalue of G that has an eigenvector x whose coordinates are all non-negative. Moreover, if G is connected, then x is strictly positive.

If $r \geq 1$ is an integer, a set S of vertices of a graph G is said to be r -*apart* if any two vertices in S are at a distance at least $r + 1$ in G . We denote by $\alpha_r(G)$ the maximum cardinality of a vertex set in G that is r -apart. Note that $\alpha_1(G)$ is the usual independence number of the graph.

Let G be a graph, $v \in V(G)$, and let r be an integer. We denote by $G_r(v)$ the induced subgraph of G on vertices that are at a distance at most r from v . The subgraph $G_r(v)$ is called the r -*ball* around v in G .

We allow infinite graphs, but they will always be locally finite. In particular, the r -ball around any vertex of a graph G is always finite.

2. UNIVERSAL COVERS AND SUBCOVERS

Let $D = [d_{ij}]_{i,j=1}^t$ be a square matrix of order $t \geq 1$ whose entries d_{ij} are non-negative integers. For $i = 1, \dots, t$, we define the i^{th} *degree* in D as the integer $d_i = \sum_{j=1}^t d_{ij}$. Suppose that D further satisfies the following conditions:

- (D1) If $d_{ij} = 0$, then also $d_{ji} = 0$.
- (D2) The graph of D is connected; i.e., for every $i, k \in \{1, \dots, t\}$ there are integers m_1, m_2, \dots, m_s in $\{1, \dots, t\}$, where $m_1 = i$, $m_s = k$, and $d_{m_j m_{j+1}} > 0$ for $j = 1, \dots, s - 1$.
- (D3) For every sequence of (distinct) integers m_1, m_2, \dots, m_s in $\{1, \dots, t\}$, we have

$$d_{m_1 m_2} d_{m_2 m_3} \cdots d_{m_{s-1} m_s} d_{m_s m_1} = d_{m_1 m_s} d_{m_s m_{s-1}} \cdots d_{m_3 m_2} d_{m_2 m_1}.$$

Such a matrix is called a t -*partite degree matrix*.

Let $D = [d_{ij}]$ be a t -partite degree matrix. If a graph G admits a partition of its vertex set into t classes, $V(G) = U_1 \cup \dots \cup U_t$, such that every vertex in U_i has precisely d_{ij} neighbors in U_j , for all $i, j = 1, \dots, t$, then we say that D is a t -*partite degree matrix* for G . The corresponding partition $U_1 \cup \dots \cup U_t$ is said to be an *equitable partition* for D ; see, e.g. [9].

Lemma 2.1. *Let D be a t -partite degree matrix.*

- (a) *There exists a finite graph G whose degree matrix is D .*
- (b) *There exists a tree T_D whose degree matrix is D . The tree T_D is determined up to isomorphism.*

Proof. (a) First we remark that condition (D3) implies that the set of equalities $n_i d_{ij} = n_j d_{ji}$, $i, j \in \{1, \dots, t\}$, has a positive solution n_1, \dots, n_t . Since all d_{ij} are integers, there is a solution whose values n_i ($i = 1, \dots, t$) are positive integers. To obtain a graph G , we take vertex sets U_i of cardinalities n_i for $i = 1, \dots, t$, and join U_i and U_j so that the edges between them form a (d_{ij}, d_{ji}) -biregular bipartite graph. Then it is clear that D is a t -partite degree matrix for G .

(b) To get T_D , we just take what is known as the universal cover of the graph G obtained in part (a).

We add a short proof of existence of T_D that does not use property (D3), which is needed in (a). Let us first assume that $d_i \geq 2$ for $i = 1, \dots, t$. This case is

intuitively clear, and we leave the details of the proof for the reader. Note that T_D is always infinite in this case.

The rest of the proof is by induction on t . We may assume that $d_t \leq 1$. If $t = 1$, then T_D is either a single vertex (if $d_1 = 0$) or an edge (if $d_1 = 1$). If $t > 1$, then (D2) implies that $d_t = 1$ and the non-zero element in row t of D is not d_{tt} . Thus, there is a unique $j < t$ such that $d_{tj} = 1$. Let $p = d_{jt}$. By (D1), we conclude that $p > 0$ and all other elements in the column t of D are zero. Let D' be the submatrix of D obtained by deleting the last row and the last column. Since this operation acts like removing a vertex of degree 1 from a graph, D' still satisfies (D1)–(D2), and hence we can apply the induction hypothesis to find the tree $T_{D'}$. Finally, we obtain T_D by adding, to each vertex in V_j , p pendant edges. All new vertices are of degree 1 and form the class V_t in T_D . \square

The tree T_D is called the *universal cover* of the multipartite degree matrix D . Let $V_1 \cup \dots \cup V_t$ be the corresponding equitable partition of $V(T_D)$. If G is any graph whose t -partite degree matrix is D , there is a covering projection $\pi_G : T_D \rightarrow G$ which maps vertices in V_i onto the i th class of the equitable partition of $V(G)$.

Covering projections, universal covers and equitable partitions are regularly used in algebraic graph theory. In the sequel we shall introduce a weaker notion, distinguished by the prefix ‘sub’, in which only those properties that are important for our main results will be preserved.

A degree matrix D is said to be a *t -partite subdegree matrix* for a graph G if there is a graph homomorphism $\pi_G^D : T_D \rightarrow G$ which is *locally 1-1*; i.e., for each vertex $v \in V(T_D)$, π_G^D maps edges incident with v injectively to the edges incident with $\pi_G^D(v)$. The homomorphism $\pi_G^D : T_D \rightarrow G$ is called a *subuniversal projection* and the tree T_D is a *subuniversal cover* of G .

If $\pi_G^D : T_D \rightarrow G$ is a subuniversal projection, let $U_i = \pi_G^D(V_i) \subseteq V(G)$, $i = 1, \dots, t$. Then it is easy to see that the (not necessarily disjoint) vertex sets U_1, \dots, U_t satisfy the following condition: Every vertex in U_i has at least d_{ij} neighbors in U_j , for all $i, j = 1, \dots, t$. This gives a necessary condition for existence of a subuniversal projection. Unfortunately, this condition is not sufficient. But if we ask that every vertex in U_i has at least $d_{ij} + 1$ neighbors in U_j for all $i, j = 1, \dots, t$, then the existence of a subuniversal projection to G is easily verified.

Theorem 2.2. *Suppose that D is a subdegree matrix for a (possibly infinite) graph G , and let T_D be the corresponding subuniversal cover. Let $v \in V(G)$ and let $s \in V(T_D)$ be a vertex that is mapped to v via a subuniversal projection π_G^D . Then for every $r \geq 0$ the spectral radius of the r -ball in G is at least as large as the spectral radius of the corresponding r -ball in T_D , $\rho(G_r(v)) \geq \rho(T_{D,r}(s))$.*

Proof. The spectral radius of a connected graph H can be expressed as

$$(1) \quad \rho(H) = \limsup_{q \rightarrow \infty} (w_{2q}(H, u))^{1/(2q)},$$

where $w_{2q}(H, u)$ denotes the number of closed walks of length $2q$ in H starting at the vertex $u \in V(H)$. Every closed walk in $T_{D,r}(s)$ starting at s is projected by π_G^D to a closed walk in $G_r(v)$ starting at v . The projection of these walks is 1-1, since π_G^D is locally 1-1. Hence,

$$(2) \quad w_{2q}(G_r(v), v) \geq w_{2q}(T_{D,r}(s), s).$$

This inequality in combination with (1) implies that $\rho(G_r(v)) \geq \rho(T_{D,r}(s))$. \square

3. THE SPECTRAL RADIUS OF INFINITE TREES

If G is a connected infinite (locally finite) graph, we define its spectral radius $\rho(G)$ as

$$(3) \quad \rho(G) = \lim_{r \rightarrow \infty} \rho(G_r(v)),$$

where v is any vertex of G . It is easy to see that the limit exists (it may be infinite if the degrees of G have no finite upper bound) and that it is independent of the choice of v . The spectral radius of infinite graphs defined above coincides with the notion obtained through the spectral theory of linear operators in Hilbert spaces; we refer to [18] for an overview.

The monotonicity property of the spectral radius of finite graphs implies that for every connected finite graph H and any proper subgraph H' of H , we have $\rho(H') < \rho(H)$. Since G is connected, infinite, and locally finite, $G_r(v) \neq G_{r+1}(v) \neq G$ for every $r \geq 0$. This implies that

$$\rho(G_r(v)) < \rho(G_{r+1}(v)) < \rho(G).$$

Let us remark that the spectral radius of an infinite d -regular tree is equal to $2\sqrt{d-1}$, the value that appears in the definition of Ramanujan graphs. This was proved by Kesten [14]; see also Dynkin and Malyutov [6], Cartier [3], and Woess [24]. We will use the spectral radius of universal cover trees introduced in the previous section to replace the Ramanujan bound $2\sqrt{d-1}$ with the corresponding bound suitable for our multipartite generalization.

In the special case when the graph is the infinite d -regular tree, which shall be denoted by \mathbb{T}_d , it is easy to determine the precise rate of convergence in (3).

Theorem 3.1. *For every integer $d \geq 2$, we have*

$$\rho(\mathbb{T}_{d,r}) > 2\sqrt{d-1} \left(1 - \frac{\pi^2}{r^2} + O(r^{-3}) \right).$$

Proof. Let $w_q(G, v_0)$ denote the number of closed walks of length q . It will be convenient to consider the subtree \mathbb{T}'_d of \mathbb{T}_d which is equal to the connected component containing the vertex v_0 of the subgraph obtained after deleting an edge of \mathbb{T}_d incident with v_0 . The vertex v_0 has degree $d-1$ in \mathbb{T}'_d , while all other vertices still have degree d . The tree \mathbb{T}'_d has a natural projection onto the one-way-infinite path P_∞ (whose vertices we denote by the non-negative integers $0, 1, 2, \dots$) such that all vertices at distance i from v_0 are mapped onto the vertex i in P_∞ . Every closed walk (based at v_0) of length $2q$ in \mathbb{T}'_d is projected onto a closed walk in P_∞ based at the vertex 0. Moreover, the r -ball $\mathbb{T}'_{d,r}(v_0)$ in \mathbb{T}'_d is projected onto the path $P_{r+1} \subset P_\infty$ on vertices $0, 1, \dots, r$.

Whenever we walk away from v_0 in \mathbb{T}'_d , we have $d-1$ choices to do so. This implies that

$$(4) \quad w_{2q}(\mathbb{T}'_{d,r}, v_0) = (d-1)^q w_{2q}(P_{r+1}, 0).$$

When $q \rightarrow \infty$, the quantities raised to the power $1/(2q)$ tend to the spectral radii of the corresponding graphs, and we conclude that $\rho(\mathbb{T}'_{d,r}) = \sqrt{d-1} \rho(P_{r+1}) = 2\sqrt{d-1} \cos(\frac{\pi}{r+2}) = 2\sqrt{d-1} (1 - \frac{\pi^2}{r^2} + O(r^{-3}))$. Since $\mathbb{T}'_{d,r}$ is a proper finite subgraph of $\mathbb{T}_{d,r}$, this implies the (strict) inequality of the theorem. \square

The rate of convergence is likely the same for more general universal covers of finite graphs. We propose the following conjecture.

Conjecture 3.2. *For every multipartite degree matrix D there exists a constant $c = c(D)$ such that for every $s \in V(T_D)$ we have*

$$\rho(T_{D,r}(s)) \geq \rho(T_D) - cr^{-2}.$$

4. MULTIPARTITE RAMANUJAN GRAPHS

In this section we introduce a generalized notion of Ramanujan graphs. The following lemma shows that we cannot simply compare $\lambda_2(G)$ with $\rho(T_D)$, as is the case for d -regular graphs.

Lemma 4.1. *If D is a t -partite degree matrix, then all eigenvalues of D are real and their algebraic multiplicity is equal to their geometric multiplicity. If D is a multipartite degree matrix of a finite graph G , then every eigenvalue of D is also an eigenvalue of G . Moreover, $\rho(G) = \rho(D)$.*

Proof. Let n_1, \dots, n_t be a positive solution of the system $n_i d_{ij} = n_j d_{ji}$, $i, j \in \{1, \dots, t\}$, which was shown to exist in the proof of Lemma 2.1(a). If R is the diagonal matrix of order t whose entry R_{ii} is equal to $n_i^{1/2}$ ($i = 1, \dots, t$), then RDR^{-1} is a symmetric matrix. This implies the first part of the lemma.

To verify the second part, let λ be an eigenvalue of D , and let $y = (y_i \mid i = 1, \dots, t)$ be an eigenvector for λ . Let $V_1 \cup \dots \cup V_t$ be the partition of $V(G)$ corresponding to the degree matrix D . If we set $x_v = y_i$ for every $v \in V_i$, then it is easy to see that $x = (x_v \mid v \in V(G))$ is an eigenvector of the adjacency matrix of G for the eigenvalue λ .

To prove the last claim, observe that the eigenvalue $\rho(D)$ has a positive eigenvector by the Perron-Frobenius theorem. Its lift in G is a positive eigenvector of G for the eigenvalue $\rho(D)$. Again, by applying the Perron-Frobenius theorem, we conclude that this eigenvector corresponds to the largest eigenvalue of G . \square

Let D be a t -partite degree matrix. Let k be the largest integer such that $\lambda_k(D) \geq \rho(T_D)$. Note that k exists since $\lambda_1(D) \geq \rho(T_D)$. We say that a finite graph G with degree matrix (resp. subdegree matrix) D is *D -Ramanujan* (resp. *D^+ -Ramanujan*) if $\lambda_{k+1}(G) \leq \rho(T_D)$. We believe that there is an abundance of generalized Ramanujan graphs and propose the following conjectures (in which we assume that the minimum degree of D is at least 2).

Conjecture 4.2. *If there exists a D -Ramanujan graph for a multipartite degree matrix D , then there exist infinitely many D -Ramanujan graphs.*

Conjecture 4.3. *If there exists a D^+ -Ramanujan graph for a multipartite degree matrix D , then there exist infinitely many D^+ -Ramanujan graphs.*

Conjecture 4.4. *If D is a degree matrix of order $t \geq 2$ and $\lambda_2(D) < \rho(T_D)$, then there exist infinitely many D -Ramanujan graphs.*

One cannot exclude the possibility that there exist infinitely many D -Ramanujan graphs for every degree matrix D , but our knowledge is too limited at this point to propose this as a conjecture.

5. A GENERALIZED ALON-BOPPANA-SERRE THEOREM

Theorem 5.1. *Let D be a multipartite degree matrix, and let $\rho_D = \rho(T_D)$.*

(a) *For every $\varepsilon > 0$ there exists an integer $r = r(D, \varepsilon)$ such that for every integer $k \geq 1$ and for every graph G , if D is a subdegree matrix of G and $\alpha_{2r+1}(G) \geq k$, then $\lambda_k(G) \geq \rho_D - \varepsilon$.*

(b) *For $\varepsilon > 0$ and every positive integer Δ , there exists a constant $c = c(D, \Delta, \varepsilon) > 0$ such that every graph G of order n , of maximum degree at most Δ and with subdegree matrix D has at least cn eigenvalues that are larger than $\rho_D - \varepsilon$.*

Proof. (a) Let $r = r(D, \varepsilon)$ be the smallest integer such that $\rho(T_{D,r}) \geq \rho_D - \varepsilon$, and let k and G be as specified. Since $\alpha_{2r+1}(G) \geq k$, there are vertices v_1, \dots, v_k that are $(2r + 2)$ -apart. The r -balls $G_r(v_1), \dots, G_r(v_k)$ around these vertices are not only pairwise disjoint but also form an induced subgraph of G . By the eigenvalue interlacing property for induced subgraphs, we know that

$$\lambda_k(G) \geq \lambda_k(G_r(v_1) \cup \dots \cup G_r(v_k)) \geq \min\{\rho(G_r(v_i)) \mid 1 \leq i \leq k\}.$$

By Theorem 2.2 and by our choice of r , we have

$$\rho(G_r(v_i)) \geq \rho(T_{D,r}(s_i)) \geq \rho_D - \varepsilon.$$

This completes the proof of (a).

(b) This part follows from (a). It is just to be noted that any $(2r + 1)$ -ball in G contains at most $B = \frac{\Delta}{\Delta-2}(\Delta - 1)^{2r+1}$ vertices. Thus, $\alpha_{2r+1}(G) \geq n/B$, and hence part (a) applies with $c = B^{-1}$. \square

It is worth mentioning that the condition involving $\alpha_{2r+1}(G)$ in Theorem 5.1(a) is necessary if we only assume that D is a subdegree matrix. Simple examples showing this are provided by the family of all complete graphs K_n whose second largest eigenvalue is always equal to -1 or by the family of all complete bipartite graphs $K_{m,n}$ whose second eigenvalue is 0 .

For the special case when $D = [d]$, Theorem 3.1 gives the precise description for the values $r(d, \varepsilon)$ and $c(d, \Delta, \varepsilon)$ in Theorem 5.1. By Theorem 3.1,

$$r = r(d, \varepsilon) = \pi \left(\frac{2\sqrt{d-1}}{\varepsilon} \right)^{1/2} (1 + O(d^{-1/4}\varepsilon^{1/2}))$$

and

$$c(d, \Delta, \varepsilon) = \frac{\Delta}{\Delta-2}(\Delta - 1)^{-(2r+1)}$$

will do the job.

As an example, let us consider the following special case. The bipartite degree matrix

$$(5) \quad D = \begin{bmatrix} 0 & d_1 \\ d_2 & 0 \end{bmatrix}$$

involves, in particular, all bipartite graphs with bipartition $V = A \cup B$ whose degrees in A are at least d_1 and whose degrees in B are at least d_2 . The spectral radius of T_D is (cf. [18])

$$\rho(T_D) = \sqrt{d_1 - 1} + \sqrt{d_2 - 1}.$$

Thus, only finitely many bipartite (d_1, d_2) -biregular graphs have their k th eigenvalue ($k \geq 2$) smaller than $\rho(T_D) - \varepsilon$. Theorem 5.1 suggests the following strengthening, which we will prove directly by using Theorem 2.2.

Corollary 5.2. *Let $d_1 \leq d_2 \leq d$ be positive integers, and let \mathcal{G}_{d_1, d_2}^d be the set of all graphs whose maximum vertex degree is at most d and whose vertex set is the union of (not necessarily disjoint) subsets U_1, U_2 , such that every vertex in U_i has at least d_i neighbors in U_{3-i} for $i = 1, 2$. For every $\varepsilon > 0$, every n -vertex graph $G \in \mathcal{G}_{d_1, d_2}^d$ has $\Omega_\varepsilon(n)$ eigenvalues larger than $\sqrt{d_1 - 1} + \sqrt{d_2 - 1} - \varepsilon$.*

Proof. We claim that there exists a subuniversal projection π_G^D , where D is the degree matrix given in (5). The tree T_D is (d_1, d_2) -biregular. We map a vertex v_0 of degree d_1 in T_D onto the vertex $u \in U_1$. After fixing v_0 , we extend the mapping to a locally 1-1 homomorphism in a greedy fashion (by taking the breadth-first search order of vertices of T_D starting at v_0) so that vertices of degree d_i are mapped to U_i , $i = 1, 2$.

Let v_1 be a neighbor of v_0 in T_D and let $v = \pi_G^D(v_1) \in U_2$. Theorem 2.2 shows that for large enough $r = r(d_1, d_2, \varepsilon)$,

$$(6) \quad \rho(G_r(u)) \geq \rho(T_{D,r}(v_0)) \geq \rho(T_D) - \varepsilon,$$

$$(7) \quad \rho(G_r(v)) \geq \rho(T_{D,r}(v_1)) \geq \rho(T_D) - \varepsilon.$$

Since the maximum degree of G is bounded by d , the $(2r + 1)$ -balls in G have a bounded number of vertices, say at most B . Therefore, $\alpha_{2r+1}(G) \geq n/B$, and so there are at least this many pairwise non-adjacent induced r -balls around vertices in G . As before, the eigenvalue interlacing theorem and (6)–(7) imply that linearly many eigenvalues of G are larger than

$$\rho(T_D) - \varepsilon = \sqrt{d_1 - 1} + \sqrt{d_2 - 1} - \varepsilon. \quad \square$$

6. GLOBAL GIRTH AND RAMANUJAN GRAPHS

All known Ramanujan graphs are Cayley graphs, and their girth increases with their order. We shall use the method of this paper to explain why the girth cannot be bounded. Actually, we shall prove that a small girth condition implies that d -regular graphs are “far from being Ramanujan”; see Theorem 6.2 below.

Let G be a graph. A closed walk $v_1 v_2 \dots v_k v_1$ is *retracting-free* if $v_{i-1} \neq v_{i+1}$ for $i = 1, \dots, k$ (where $v_0 = v_k$ and $v_{k+1} = v_1$). It is easy to see that if G is a finite graph with minimum degree at least 2, then for every vertex v of G there exists a retracting-free closed walk through v .

Let $g(v)$ be the length of a shortest retracting-free closed walk through v . The *universal girth* of G , denoted by $m(G)$, is the smallest integer k such that every vertex in G has a retracting-free closed walk of length k . Let us observe that $m(G)$ is at most the least common multiple of the values $g(v)$, $v \in V(G)$. Also, if G is vertex-transitive, then $m(G)$ is equal to the girth of G .

Let $\mathbb{X}_{d,g}$ be the graph obtained from the $(d - 2)g$ -regular tree T by expanding each vertex $v \in V(T)$ into the cycle C_v of length g , such that each vertex of C_v is incident with $d - 2$ of the edges of T incident with v . See Figure 1, which shows the case of $d = 4$ and $g = 4$. The graph $\mathbb{X}_{d,g}$ is the Cayley graph of the free product of $d - 2$ copies of \mathbb{Z}_2 and one copy of \mathbb{Z}_g (with the natural generating set).

Paschke [22] determined the spectral radius of $\mathbb{X}_{d,g}$:

Theorem 6.1 (Paschke). *For $d \geq 3$ and $g \geq 3$, the graph $\mathbb{X}_{d,g}$ has spectral radius*

$$\min_{s>0} (d - 2) \phi \left(\frac{1 + \cosh sg}{\sinh sg \sinh s} \right) + 2 \cosh s > 2\sqrt{d - 1},$$




FIGURE 1. Expanding a vertex into a cycle

where $\phi(t) = \frac{\sqrt{1+t^2}-1}{t}$.

Paschke [22] used this result to provide a non-trivial lower bound on the spectral radius of infinite vertex-transitive graphs of the given girth g . He showed that a vertex transitive d -regular graph containing a g -cycle has spectral radius at least $\rho(\mathbb{X}_{d,g})$. The formula in Theorem 6.1 gives a lower bound of the form

$$2\sqrt{d-1} + \frac{2(d-2)}{(d-1)^{(g+1)/2}} h(d, g),$$

where h is a function such that for every $g \geq 3$, $\lim_{d \rightarrow \infty} h(d, g) = 1$, and for every $d \geq 3$, $\lim_{g \rightarrow \infty} h(d, g) = 1$.

Now, we strengthen the Alon-Boppana-Serre theorem by showing that the lower bound $2\sqrt{d-1} - \varepsilon$ can be replaced by $2\sqrt{d-1} + \delta$ for some $\delta > 0$ if graphs have a bounded universal girth.

Theorem 6.2. *For every $\Delta \geq d \geq 3$ and every $g \geq 3$, there exist $\delta > 0$ and $c > 0$ such that every n -vertex graph G with minimum degree at least d , maximum degree at most Δ and universal girth at most g has at least $\lceil cn \rceil$ eigenvalues that are larger than $2\sqrt{d-1} + \delta$.*

Proof. The proof follows the same pattern as in the proof of Theorem 5.1, except that we use the graph $\mathbb{X}_{d,m}$, where $m = m(G) \leq g$ is the universal girth of G , playing the role of the universal cover \mathbb{T}_d . Here, we have to take r large enough so that $\rho((\mathbb{X}_{d,m})_r) \geq 2\sqrt{d-1} + \delta$. Such an r exists because of (3) and since $\rho(\mathbb{X}_{d,m}) > \rho(\mathbb{T}_d) = 2\sqrt{d-1}$. \square

It is straightforward to generalize the proof of Theorem 6.2 to the setting of degree matrices. What we need is just an analogue of the Paschke theorem. However, we do not intend to dig into the details in this paper.

7. THE OTHER SIDE OF THE SPECTRUM

As shown in the previous section, small universal girth yields improved lower bounds on large eigenvalues, so Ramanujan graphs must have growing girth. On the other hand, large girth has some further consequences. In particular, it shows that the negative eigenvalues satisfy the Alon-Boppana-Serre property as well.

Let us first formulate the monopartite version for the negative eigenvalues. It involves the notion of the *odd girth* of the graph, meaning the length of a shortest cycle of odd length in the graph. (If G is bipartite, then the odd girth is ∞ .) This

result was obtained earlier by Friedman [8] and Nilli [21]; it also appears in Cioabă [4] (with a slightly weaker estimate of ε).

Theorem 7.1. *For every $\Delta \geq d \geq 2$ and $g \geq 3$, there exists a positive constant $c = c(d, \Delta, g) > 0$ such that every graph G of order n , of minimum degree d , maximum degree at most Δ , and with odd girth at least g has at least cn eigenvalues that are larger than $2\sqrt{d-1}(1-\varepsilon)$ and has at least cn eigenvalues that are smaller than $-2\sqrt{d-1}(1-\varepsilon)$, where $\varepsilon = (\frac{2\pi}{g})^2 + O(g^{-3})$.*

Proof. (Sketch) The proof is essentially the same as the proof of Theorem 5.1(b), where we take $r = \lfloor \frac{1}{2}g \rfloor - 1$ and apply the estimate of Theorem 3.1. The assumption that the odd girth is more than $2r+1$ shows that the r -balls in G contain no cycles of odd length. In particular, they are bipartite, and hence their spectrum is symmetric with respect to 0. Thus, knowing that the spectral radius λ is large, we conclude that the smallest eigenvalue $-\lambda$ is large in absolute value. Now, we can use the interlacing theorem for the smallest eigenvalues of G compared to the eigenvalues of the induced subgraph of G consisting of disjoint r -balls around $\lceil cn \rceil$ vertices that are $(2r+2)$ -apart. \square

The generalized version of Theorem 7.1 holds as well. The proof is the same, except that we do not provide an explicit estimate on ε in terms of the odd girth.

Theorem 7.2. *Let D be a multipartite degree matrix, and let $\rho_D = \rho(T_D)$. For every $\varepsilon > 0$ and every positive integer Δ , there exists an integer $g = g(D, \varepsilon)$ and a positive constant $c = c(D, \Delta, \varepsilon) > 0$ such that every graph G of order n , of maximum degree at most Δ , with subdegree matrix D , and with odd girth at least g has at least cn eigenvalues that are smaller than $-\rho_D + \varepsilon$.*

REFERENCES

1. N. Alon, Eigenvalues and expanders, *Combinatorica* 6 (1986) 83–96. MR875835 (88e:05077)
2. N. Alon, V. D. Milman, λ_1 , isoperimetric inequalities for graphs, and superconcentrators, *J. Combin. Theory Ser. B* 38 (1985) 73–88. MR782626 (87b:05092)
3. P. Cartier, Harmonic analysis on trees, in “Harmonic analysis on homogeneous spaces” (Proc. Sympos. Pure Math., Vol. XXVI, Williams College, Williamstown, Mass., 1972), AMS, 1973, pp. 419–424. MR0338272 (49:3038)
4. S. M. Cioabă, Eigenvalues of graphs and a simple proof of a theorem of Greenberg, *Linear Algebra Appl.* 416 (2006) no. 2–3, 776–782. MR2242462 (2007h:05097)
5. G. Davidoff, P. Sarnak, A. Valette, *Elementary number theory, group theory, and Ramanujan graphs*, Cambridge University Press, Cambridge, 2003. MR1989434 (2004f:11001)
6. E. B. Dynkin, M. B. Malyutov, Random walks on groups with a finite number of generators, *Soviet Math. Doklady* 2 (1961) 399–402. MR0131904 (24:A1751)
7. K. Feng, W.-C. Winnie Li, Spectra of hypergraphs and applications, *J. Number Theory* 60 (1996) 1–22. MR1405722 (97f:05128)
8. J. Friedman, Some geometric aspects of graphs and their eigenfunctions, *Duke Math. J.* 69 (1993) 487–525. MR1208809 (94b:05134)
9. C. Godsil, G. Royle, *Algebraic Graph Theory*, Springer, 2001. MR1829620 (2002f:05002)
10. Y. Greenberg, Spectra of graphs and their covering trees (in Hebrew), Ph.D. thesis, Hebrew University of Jerusalem, 1995.
11. S. Hoory, A lower bound on the spectral radius of the universal cover of a graph, *J. Combin. Theory, Ser. B* 93 (2005) 33–43. MR2102266 (2005k:05144)
12. S. Hoory, N. Linial, A. Wigderson, Expander graphs and their applications, *Bull. Amer. Math. Soc. (N.S.)* 43 (2006) 439–561. MR2247919 (2007h:68055)
13. R. A. Horn, C. R. Johnson, *Matrix Analysis*, Cambridge Univ. Press, Cambridge, 1985. MR832183 (87e:15001)

14. H. Kesten, Symmetric random walks on groups, *Trans. Amer. Math. Soc.* 92 (1959) 336–354. MR0109367 (22:253)
15. A. Lubotzky, T. Nagnibeda, Not every uniform tree covers Ramanujan graphs, *J. Combin. Theory, Ser. B* 74 (1998) 202–212. MR1654133 (2000g:05052)
16. A. Lubotzky, R. Phillips, P. Sarnak, Ramanujan graphs, *Combinatorica* 8 (1988) 261–277. MR963118 (89m:05099)
17. G. A. Margulis, Explicit group-theoretic constructions of combinatorial schemes and their applications in the construction of expanders and concentrators (in Russian), *Problemy Peredachi Informatsii* 24 (1988) 51–60; Engl. transl. in *Problems Inform. Transmission* 24 (1988) 39–46. MR939574 (89f:68054)
18. B. Mohar, W. Woess, A survey on spectra of infinite graphs, *Bull. London Math. Soc.* 21 (1989) 209–234. MR986363 (90d:05162)
19. M. Morgenstern, Existence and explicit constructions of $q + 1$ regular Ramanujan graphs for every prime power q , *J. Combin. Theory Ser. B* 62 (1994) 44–62. MR1290630 (95h:05089)
20. A. Nilli, On the second eigenvalue of a graph, *Discrete Math.* 91 (1991) 207–210. MR1124768 (92j:05124)
21. A. Nilli, Tight estimates for eigenvalues of regular graphs, *Electron. J. Combin.* 11 (2004) Note 9, 4 pp. MR2056091
22. W. Paschke, Lower bound for the norm of a vertex-transitive graph, *Math. Z.* 213 (1993) 225–239. MR1221715 (94j:05085)
23. J.-P. Serre, Répartition asymptotique des valeurs propres de l'opérateur de Hecke T_p , *J. Amer. Math. Soc.* 10 (1997) 75–102. MR1396897 (97h:11048)
24. W. Woess, *Random walks on infinite graphs and groups*, Cambridge University Press, Cambridge, 2000. MR1743100 (2001k:60006)

DEPARTMENT OF MATHEMATICS, SIMON FRASER UNIVERSITY, BURNABY, BRITISH COLUMBIA,
V5A 1S6, CANADA

E-mail address: mohar@sfu.ca