

UNIFORM ALGEBRAS GENERATED BY HOLOMORPHIC AND CLOSE-TO-HARMONIC FUNCTIONS

GAUTAM BHARALI AND SUSHIL GORAI

(Communicated by Franc Forstneric)

ABSTRACT. The initial motivation for this paper is to discuss a more concrete approach to an approximation theorem of Axler and Shields, which says that the uniform algebra on the closed unit disc \mathbb{D} generated by z and h , where h is a nowhere-holomorphic harmonic function on \mathbb{D} that is continuous up to $\partial\mathbb{D}$, equals $\mathcal{C}(\overline{\mathbb{D}})$. The abstract tools used by Axler and Shields make harmonicity of h an essential condition for their result. We use the concepts of plurisubharmonicity and polynomial convexity to show that, in fact, the same conclusion is reached if h is replaced by $h + R$, where R is a non-harmonic perturbation whose Laplacian is “small” in a certain sense.

1. INTRODUCTION AND STATEMENT OF RESULTS

This paper is motivated by the following result of Axler and Shields [1] (in what follows, \mathbb{D} will denote the open unit disc in \mathbb{C} centered at the origin):

Result 1.1 ([1], Theorem 4). Let h be a function in $\mathcal{C}(\overline{\mathbb{D}})$ that is harmonic but nowhere holomorphic on \mathbb{D} . Then, $[z, h]_{\overline{\mathbb{D}}} = \mathcal{C}(\overline{\mathbb{D}})$.

Recall that $[z, h]_{\overline{\mathbb{D}}}$ denotes the uniform algebra on $\overline{\mathbb{D}}$ generated by z and h . Axler and Shields use results that are abstract and of extremely general scope, such as the Bishop Antisymmetric Decomposition, to deduce their theorems. Harmonicity plays a very central role in their approach, and it is difficult to answer even this simple question: *to what extent can we allow harmonicity to fail, by adding a small perturbation R to h , and yet recover the conclusion of Result 1.1 with $h+R$ replacing h ?*

Axler and Shields themselves imply that they tried to prove Result 1.1 without the use of their deep result on the $L^\infty(\mathbb{D})$ -subalgebra $H^\infty(\mathbb{D})[h]$ (which is where harmonicity plays a key role) but to no avail (see page 636 of [1] for their statement). Hence, there is an interest in a more explicit approach even to Result 1.1.

Received by the editors January 20, 2010 and, in revised form, May 13, 2010 and June 15, 2010.

2010 *Mathematics Subject Classification*. Primary 30E10, 32E20, 32U05, 46J15.

Key words and phrases. Harmonic function, plurisubharmonic function, polynomially convex.

The first author is supported by the DST via the Fast Track grant SR/FTP/MS-12/2007.

The second author is supported by CSIR-UGC fellowship 09/079(2063). Support is also provided by the UGC under DSA-SAP, Phase IV.

©2010 American Mathematical Society
Reverts to public domain 28 years from publication

We are able, using plurisubharmonic functions and polynomial convexity in a simple way, to prove an Axler-Shields-type result which states that $[z, h + R]_{\mathbb{D}} = \mathcal{C}(\overline{\mathbb{D}})$, where R is a small (in an appropriate sense) non-harmonic perturbation. Furthermore, taking $R = 0$ in our result reproduces the conclusion of Result 1.1, thus providing a different approach to the Axler-Shields theorem.

The central result of this article is:

Theorem 1.2. *Let $h : \overline{\mathbb{D}} \rightarrow \mathbb{C}$ be a function that is harmonic on \mathbb{D} and belongs to $\mathcal{C}(\overline{\mathbb{D}})$. Let $R \in \mathcal{C}^2(\mathbb{D}) \cap \mathcal{C}(\overline{\mathbb{D}})$, and suppose that R is a non-harmonic perturbation of h that is small in the following sense:*

- a) *The set $\{z \in \mathbb{D} : \partial_{\bar{z}}(h + R)(z) = 0\}$ has zero Lebesgue measure.*
- b) *The Laplacian of R has the bound*

$$(1.1) \quad |\Delta R(z)| \leq C \frac{|\partial_{\bar{z}}(h + R)(z)|^2}{\sup_{\overline{\mathbb{D}}} |h + R|} \quad \forall z \in \mathbb{D},$$

for some constant $C \in (0, 1)$.

Then, $[z, h + R]_{\mathbb{D}} = \mathcal{C}(\overline{\mathbb{D}})$.

Remark 1.3. Observe that we can recover Result 1.1 from the above theorem. First, $R \equiv 0$ certainly satisfies (b). Note, furthermore, that $\partial_{\bar{z}}h$ is anti-holomorphic on \mathbb{D} . Its zeros in \mathbb{D} thus form a discrete subset of \mathbb{D} , hence a set of zero Lebesgue measure. All the hypotheses of Theorem 1.2 are satisfied, and hence $[z, h]_{\mathbb{D}} = \mathcal{C}(\overline{\mathbb{D}})$.

Before proceeding to the proof, let us glance at the central ideas involved. The proof may be summarised as follows (in what follows, given a function f and a set $S \subset \text{dom}(f)$, $\text{Gr}_S(f)$ will denote the set $\text{graph}(f) \cap (S \times \mathbb{C})$, i.e. the portion of the graph of f whose projection onto the first coordinate is S):

- We start with a construction that goes back to Hörmander and Wermer [5]: we define the function

$$\psi_r(z, w) := |w - (h + R)(rz)|^2, \quad (z, w) \in \overline{D(0; r^{-1})} \times \mathbb{C}, \quad r \in (0, 1),$$

which vanishes precisely on the graph of $(h + R)(r \cdot)$. We use the condition (1.1) to show that ψ_r is plurisubharmonic in $\Delta_r := D(0; r^{-1}) \times D(0; \rho)$, where $\rho > 0$ is large enough to contain the aforementioned graph.

- From the last fact and the fact that each Δ_r , $r \in (0, 1)$, is Runge, we realise that $\text{Gr}_{\overline{\mathbb{D}}}((h + R)(r \cdot))$ is polynomially convex. But because $(h + R)(r \cdot) \rightarrow (h + R)$ uniformly on $\overline{\mathbb{D}}$ as $r \uparrow 1$, we deduce the same for $\text{Gr}_{\overline{\mathbb{D}}}(h + R)$.
- Knowing that $\text{Gr}_{\overline{\mathbb{D}}}(h + R)$ is polynomially convex, the first condition on R allows us to appeal to a variation on a theorem of Wermer [6, Theorem 1]. Wermer's original theorem would have required us to demand that $h, R \in \mathcal{C}^1(\overline{\mathbb{D}})$. However, with very slight modifications to Wermer's proof, we can appeal to the resulting theorem to infer that $[z, h + R]_{\mathbb{D}} = \mathcal{C}(\overline{\mathbb{D}})$.

The main idea needed for the aforementioned variation on Wermer's theorem has been remarked upon in [6]. However, it might be of interest to the reader to see the relevant lemmas carefully restated to suit the present setting (i.e. with lower boundary regularity). Hence, we shall discuss this variation in Section 2. The proof of Theorem 1.2 will be presented in Section 3.

2. TECHNICAL RESULTS

We begin this section with a technical, but essentially elementary, result. We must first explain some notation. Given a domain $\Omega \subset \mathbb{C}^d$ and a real-valued function $F \in \mathcal{C}^2(\Omega)$, the *Levi form of F at z* , denoted by $\mathcal{L}F(z; \cdot)$, is the quadratic form given by

$$\mathcal{L}F(z; V) := \sum_{j,k=1}^d \frac{\partial^2 F}{\partial z_j \partial \bar{z}_k}(z) v_j \bar{v}_k \quad \forall V = (v_1, \dots, v_d) \in \mathbb{C}^d.$$

Lemma 2.1. *Let Ω be a domain in \mathbb{C} and let $f \in \mathcal{C}^2(\Omega)$. Define the function $\psi(z, w) := |w - f(z)|^2$, $(z, w) \in \Omega \times \mathbb{C}$. Then, for the Levi form $\mathcal{L}\psi(z, w; V)$, $V = (V_1, V_2) \in \mathbb{C}^2$, we have*

$$(2.1) \quad \mathcal{L}\psi(z, w; V) \geq \left(2\operatorname{Re} \left(\overline{(f(z) - w)} \partial_{z\bar{z}}^2 f(z) \right) + |\partial_{z\bar{z}} f(z)|^2 \right) |V_1|^2.$$

In particular, if $f = h + R$, where $h \in \operatorname{harm}(\Omega)$, we have

$$(2.2) \quad \mathcal{L}\psi(z, w; V) \geq \left(|\partial_{z\bar{z}}(h + R)(z)|^2 + 2\operatorname{Re} \left(\overline{(h(z) + R(z) - w)} \partial_{z\bar{z}}^2 R(z) \right) \right) |V_1|^2.$$

Proof. We compute:

$$\begin{aligned} \partial_{z\bar{z}}^2 \psi(z, w) &= 2\operatorname{Re}(\partial_{z\bar{z}}^2 f(z) \cdot \overline{(f(z) - w)}) + |\partial_z f(z)|^2 + |\partial_{\bar{z}} f(z)|^2, \\ \partial_{z\bar{w}}^2 \psi(z, w) &= -\partial_z f(z), \\ \partial_{w\bar{w}}^2 \psi(z, w) &= 1. \end{aligned}$$

Now, using the above calculation, we have the Levi form $\mathcal{L}\psi(z, w; V)$, with $V = (V_1, V_2) \in \mathbb{C}^2$, as (we denote the function mapping $(z, w) \mapsto w$ by w):

$$\begin{aligned} \mathcal{L}\psi(\cdot; V) &= (2\operatorname{Re}(\partial_{z\bar{z}}^2 f \cdot \overline{(f - w)}) + |\partial_z f|^2 + |\partial_{\bar{z}} f|^2) |V_1|^2 - 2\operatorname{Re}(\partial_z f \cdot V_1 \bar{V}_2) + |V_2|^2 \\ &= |\partial_z f \cdot V_1 - V_2|^2 + (2\operatorname{Re}(\partial_{z\bar{z}}^2 f \cdot \overline{(f - w)}) + |\partial_{\bar{z}} f|^2) |V_1|^2 \\ &\geq (2\operatorname{Re}(\partial_{z\bar{z}}^2 f \cdot \overline{(f - w)}) + |\partial_{\bar{z}} f|^2) |V_1|^2. \end{aligned}$$

The second inequality follows by replacing f by $h + R$ and noting that $\partial_{z\bar{z}}^2 h = 0$. \square

We now present the following variation on [6, Theorem 1]. We need to clarify some notation needed in its proof: given a compact subset $K \Subset \mathbb{C}^d$, we define

$$\mathcal{P}(K) := \text{the class of uniform limits on } K \text{ of holomorphic polynomials in } \mathbb{C}^d.$$

Theorem 2.2 (A variation on Theorem 1 of [6]). *Let $f : \mathbb{D} \rightarrow \mathbb{C}$ be a continuous function. Assume $\operatorname{Gr}_{\mathbb{D}}(f)$ is polynomially convex. Define*

$$\mathcal{S} := \{z \in \mathbb{D} : \text{there exists a } \mathbb{C}\text{-open neighbourhood } V_z \ni z \text{ such that } f \text{ has continuous first-order partial derivatives on } V_z\},$$

and let $W := \{z \in \mathcal{S} : \partial_{z\bar{z}} f(z) \neq 0\}$. If $\mathbb{D} \setminus W$ has zero Lebesgue measure, then $[z, f]_{\mathbb{D}} = \mathcal{C}(\mathbb{D})$.

Remark 2.3. Wermer’s original result requires that $f \in \mathcal{C}^1(\mathbb{D})$ and that $\partial_{z\bar{z}} f$ be non-vanishing. But immediately after the proof of [6, Theorem 1], it is stated that the hypothesis of [6, Theorem 1] can be weakened by letting the condition $\partial_{z\bar{z}} f(z) \neq 0$ fail on a non-empty subset of zero Lebesgue measure. The key to Theorem 2.2 is that f can also be allowed to be non-differentiable on this exceptional set. We justify this below.

Sketch of the proof of Theorem 2.2. Since the proof involves minor modifications to the original, we shall be brief. The notation $\mu \perp [z, f]_{\overline{\mathbb{D}}}$ will denote a complex measure $\mu \in \mathcal{C}(\overline{\mathbb{D}})^*$ representing a bounded linear functional on $\mathcal{C}(\overline{\mathbb{D}})$ that annihilates $[z, f]_{\overline{\mathbb{D}}}$.

Wermer's proof uses the following fact, which occurs as a part of Bishop's proof of [2, Theorem 4]:

(*) (Bishop) For any complex measure $\mu \in \mathcal{C}(\overline{\mathbb{D}})^*$, let

$$H_\mu(a) := \int_{\mathbb{C}} \frac{1}{z-a} d\mu(z).$$

If $H_\mu = 0$ *m*-a.e., then $\mu = 0$ (*m* denotes the planar Lebesgue measure on \mathbb{C}).

Wermer's strategy consists of the following two parts:

- (a) Use the Oka-Weil theorem to construct, for each $a \in \overline{\mathbb{D}}$, an open neighbourhood $\mathcal{D} \supset \text{Gr}_{\overline{\mathbb{D}}}(f)$ and a function $h \in \mathcal{O}(\mathcal{D})$ (which depend on a) such that:
- $\exists R > 0$ such that $h(\text{Gr}_{\overline{\mathbb{D}}}(f)) \subset E_R \cup \{0\}$, where E_R denotes the complement in \mathbb{C} of the closed sector $\{re^{i\theta} : 0 \leq r \leq R, |\theta| \leq \pi/4\}$;
 - h is non-vanishing on $\text{Gr}_{\overline{\mathbb{D}}}(f) \setminus \{(a, f(a))\}$;
 - $\exists h_1 \in \mathcal{O}(\mathcal{D})$ satisfying $h(z, w) = (z-a)h_1(z, w) \forall (z, w) \in \mathcal{D}$.
- (b) Apply (for a fixed $a \in \overline{\mathbb{D}}$) the dominated convergence theorem to the measure $\mu \perp [z, f]_{\overline{\mathbb{D}}}$ and the sequence

$$\{h_1(\cdot, f)P_n \circ h(\cdot, f)\}_{n \in \mathbb{Z}_+} \subset \mathcal{P}(\overline{\mathbb{D}})$$

(where the sequence $\{P_n\}_{n \in \mathbb{Z}_+}$ is as given by [6, Lemma 3]) to conclude that $H_\mu(a) = 0$. This implies that $H_\mu \equiv 0$ because the above argument works for each $a \in \overline{\mathbb{D}}$.

Since $H_\mu \equiv 0$ for every $\mu \perp [z, f]_{\overline{\mathbb{D}}}$, it follows from (*) that $[z, f]_{\overline{\mathbb{D}}} = \mathcal{C}(\overline{\mathbb{D}})$.

Observe that the inference $H_\mu \equiv 0$ (for $\mu \perp [z, f]_{\overline{\mathbb{D}}}$) is stronger than is necessary for the desired conclusion. This suggests the following *modified* two-step strategy:

- (a') Construct the objects $(\mathcal{D}, h, E_R, h_1)$ having exactly the same properties as in Part (a) above, but *only associated to each* $a \in W$.
- (b') Repeat Part (b) of Wermer's strategy for all those points $a \in \overline{\mathbb{D}}$ for which Wermer's dominated-convergence-theorem argument, showing $H_\mu(a) = 0$, still makes sense (call the complement of all such points \mathcal{E}).

It is not hard to see that $\mathcal{E} = (\overline{\mathbb{D}} \setminus W) \cup (\tilde{E} \cap \overline{\mathbb{D}})$, where:

$$\tilde{E} := \left\{ a \in \mathbb{C} : \int_{\mathbb{C}} |z-a|^{-1} d|\mu|(z) = \infty \right\}.$$

The set \tilde{E} has zero Lebesgue measure, which is a well-known fact about finite, positive Borel measures in general. By *exactly the same* considerations as in Part (b) and from the obvious fact that $\mu \perp [z, f]_{\overline{\mathbb{D}}} \implies H_\mu(a) = 0 \forall a \notin \overline{\mathbb{D}}$, our modified strategy gives us

$$\mu \perp [z, f]_{\overline{\mathbb{D}}} \implies H_\mu(a) = 0 \forall a \in \mathbb{C} \setminus \mathcal{E}.$$

Since, by hypothesis, $\mathfrak{m}(\mathcal{E}) = 0$, we infer from (*) that each $\mu \perp [z, f]_{\overline{\mathbb{D}}}$ is just the zero measure. Hence, $[z, f]_{\overline{\mathbb{D}}} = \mathcal{C}(\overline{\mathbb{D}})$. \square

3. THE PROOF OF THEOREM 1.2

We recall a standard notation that we shall use in our proof. Given a domain $\Omega \subset \mathbb{C}^d$ and a compact subset $K \Subset \Omega$, we define the $\mathcal{O}(\Omega)$ -hull of K as

$$\widehat{K}_\Omega := \{z \in \Omega : |f(z)| \leq \sup_K |f| \ \forall f \in \mathcal{O}(\Omega)\}.$$

Proof of Theorem 1.2. We begin with a preliminary observation. The estimate (1.1) may be rewritten as

$$|\Delta R(z)| \leq \frac{|\partial_{\bar{z}}(h + R)(z)|^2}{\sup_{\mathbb{D}} |h + R| + (\frac{1}{C} - 1) \sup_{\mathbb{D}} |h + R|},$$

whence we can certainly find a constant $\delta_0 > 0$ such that

$$\sup_{\mathbb{D}} |h + R| + (\frac{1}{C} - 1) \sup_{\mathbb{D}} |h + R| \geq \sup_{\mathbb{D}} |h + R| + \delta_0.$$

Hence, for the remainder of this proof, we may assume that

$$(3.1) \quad |\Delta R(z)| \leq \frac{|\partial_{\bar{z}}(h + R)(z)|^2}{\sup_{\mathbb{D}} |h + R| + \delta_0} \ \forall z \in \mathbb{D}.$$

For each $r \in (0, 1)$, let us define

$$\psi_r(z, w) := |w - (h + R)(rz)|^2, \ (z, w) \in \overline{D(0; r^{-1})} \times \mathbb{C}.$$

The Levi-form computations (2.1) and (2.2), taken together with the estimate (3.1) on ΔR , establish that

$$(3.2) \quad \psi_r \text{ is plurisubharmonic in } \Delta_r := D(0; r^{-1}) \times D(0; M + 2\delta_0) \ \forall r \in (0, 1),$$

where $M := \sup_{\mathbb{D}} |h + R|$. Given these preliminaries, we can complete the proof in two steps.

Step I. Polynomial convexity of $\text{Gr}_{\mathbb{D}}(h + R)$.

Since $(h + R)$ is uniformly continuous on $\overline{\mathbb{D}}$, it follows that:

For each $\varepsilon > 0$, $\exists \delta(\varepsilon) > 0$ such that

$$(3.3) \quad 0 < (1 - r) \leq \delta(\varepsilon) \implies |(h + R)(rz) - (h + R)(z)| < \varepsilon \ \forall z \in \overline{\mathbb{D}}.$$

Consider a point $p = (z_0, w_0) \in \overline{\mathbb{D}} \times D(0; M + 2\delta_0) \setminus \text{Gr}_{\mathbb{D}}(h + R)$. Then, by definition, $\psi_1(z_0, w_0) =: \varepsilon_p^2 > 0$. Write $r(p) := 1 - \delta(\varepsilon_p/3)$, where $\delta(\varepsilon_p/3)$ is as given by (3.3). Then, (3.3) tells us:

$$\begin{aligned} |w_0 - (h + R)(r(p)z_0)| &\geq |w_0 - (h + R)(z_0)| - |(h + R)(z_0) - (h + R)(r(p)z_0)| \\ &> 2\varepsilon_p/3, \\ \psi_{r(p)}(z, w) &= |(h + R)(z) - (h + R)(r(p)z)|^2 \\ &< \varepsilon_p^2/9 \ \forall (z, w) \in \text{Gr}_{\mathbb{D}}(h + R). \end{aligned}$$

By the last two estimates, we have just shown that

$$(3.4) \quad \psi_{r(p)}(p) > 4\varepsilon_p^2/9 \ \text{and} \ \psi_{r(p)}(z, w) < \varepsilon_p^2/9 \ \forall (z, w) \in \text{Gr}_{\mathbb{D}}(h + R).$$

Let us now write $K := \text{Gr}_{\mathbb{D}}(h+R)$. We claim that $p \notin \widehat{K}$. To do so, we invoke a well-known result of Hörmander [4, Theorem 4.3.4] which states that if $\Omega \subset \mathbb{C}^d$, $d \geq 2$, is a pseudoconvex domain and $K \Subset \Omega$ is a compact subset, then the hull \widehat{K}_{Ω} can also be expressed as

$$\widehat{K}_{\Omega} = \{z \in \mathbb{C}^d : U(z) \leq \sup_K U \quad \forall U \in \text{psh}(\Omega)\}.$$

It thus follows from (3.4) that $p \notin \widehat{K}_{\Delta_{r(p)}}$. Note that each $\Delta_{r(p)}$ is Runge. We know therefore that $\widehat{K}_{\Delta_{r(p)}} = \widehat{K}$. Since p was arbitrarily chosen, we have just shown that

$$p \notin \widehat{K} \quad \forall p \in \overline{\mathbb{D}} \times D(0; M + 2\delta_0) \setminus K.$$

Of course, it is easy to see that no point in $\mathbb{C}^2 \setminus (\overline{\mathbb{D}} \times \overline{D(0; M)})$ can belong to \widehat{K} . Hence, $K = \text{Gr}_{\mathbb{D}}(h+R)$ is polynomially convex.

Step II. Completing the proof.

We appeal to Theorem 2.2 with $(h+R)$ playing the role of f . In the terminology of Theorem 2.2,

$$\overline{\mathbb{D}} \setminus W = \partial\mathbb{D} \cup \{z \in \mathbb{D} : \partial_{\bar{z}}(h+R)(z) = 0\},$$

which, by hypothesis, has zero Lebesgue measure. We have already established that $\text{Gr}_{\mathbb{D}}(h+R)$ is polynomially convex. Thus, $(h+R)$ satisfies all the conditions stated in Theorem 2.2, and we conclude that $[z, h+R]_{\overline{\mathbb{D}}} = \mathcal{C}(\overline{\mathbb{D}})$. \square

Added in Proof. It was brought to our notice that stronger results subsuming Result 1.1 had been established by Chirka [3] in 1969. However, there are gaps in the proofs of [3, Theorem 4] and [3, Theorem 5], on which Chirka's results rely. The most significant gap is the one in the proof of Theorem 4, which is false as stated. The proof appears to presume that (in the notation of [3, Theorem 4]) \widehat{K}_A is always connected. In short: [1] gives the earliest *complete* proof of Result 1.1 that we are aware of. That said, we feel that the *basic ideas* in [3] could still (bypassing Theorems 4 and 5 entirely) be made to work. \square

ACKNOWLEDGEMENT

We thank the referee for pointing out ways in which the arguments in our first draft could be simplified and for suggesting that we examine whether the boundary regularity of the functions in Theorem 1.2 could be lowered.

REFERENCES

- [1] S. Axler and A. Shields, *Algebras generated by analytic and harmonic functions*, Indiana Univ. Math. J. **36** (1987), no. 3, 631-638. MR905614 (88h:46102)
- [2] E. Bishop, *A minimal boundary for function algebras*, Pacific J. Math. **9** (1959), 629-642. MR0109305 (22:191)
- [3] E. M. Čirka, *Approximation by holomorphic functions on smooth submanifolds in \mathbb{C}^n* (Russian), Mat. Sb. **78** (120) (1969), 101-123; AMS translation: Math. USSR Sbornik **7** (1969), 95-114. MR0239121 (39:480)
- [4] L. Hörmander, *An Introduction to Complex Analysis in Several Variables*, 3rd ed., North-Holland, Amsterdam, 1990. MR1045639 (91a:32001)

- [5] L. Hörmander and J. Wermer, *Uniform approximation on compact sets in \mathbb{C}^n* , Math. Scand. **23** (1968), 5-21. MR0254275 (40:7484)
- [6] J. Wermer, *Polynomially convex disks*, Math. Ann. **158** (1965), 6-10. MR0174968 (30:5158)

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE, BANGALORE – 560012, INDIA
E-mail address: `bharali@math.iisc.ernet.in`

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE, BANGALORE – 560012, INDIA
E-mail address: `sushil@math.iisc.ernet.in`