

A NOTE ON DISCRETENESS OF F -JUMPING NUMBERS

KARL SCHWEDE

(Communicated by Irena Peeva)

ABSTRACT. Suppose that R is a ring essentially of finite type over a perfect field of characteristic $p > 0$ and that $\mathfrak{a} \subseteq R$ is an ideal. We prove that the set of F -jumping numbers of $\tau_b(R; \mathfrak{a}^t)$ has no limit points under the assumption that R is normal and \mathbb{Q} -Gorenstein – we make no assumption as to whether the \mathbb{Q} -Gorenstein index is divisible by p . Furthermore, we also show that the F -jumping numbers of $\tau_b(R; \Delta, \mathfrak{a}^t)$ are discrete under the more general assumption that $K_R + \Delta$ is \mathbb{R} -Cartier.

1. INTRODUCTION

The test ideal is an important and subtle object associated to ideals \mathfrak{a} in positive characteristic rings R . It measures the singularities of both the ambient ring and the elements of the ideal; see [HY03]. While the test ideal was initially introduced in the celebrated theory of tight closure of Hochster and Huneke (see [HH90]), more recent interest in the test ideal has been in regards to its connection with the multiplier ideal – a fundamental invariant of higher dimensional algebraic geometry in characteristic zero; see for example [Tak06] or [MY09].

Given a normal ring R essentially of finite type over a perfect field of characteristic $p > 0$, an ideal $\mathfrak{a} \subseteq R$ and a real number $t \geq 0$, one can form the (big) test ideal $\tau_b(R; \mathfrak{a}^t)$ – an object which measures both algebraic and arithmetic properties of R and \mathfrak{a} . Inspired by the test ideal’s close relation with the multiplier ideal $\mathcal{J}(R, \mathfrak{a}^t)$, people have studied the numbers t_i where $\tau_b(R; \mathfrak{a}^{t_i})$ changes. That is, people have studied the *F-jumping numbers* (see [MTW05]), real numbers which are by definition the $t_i > 0$ such that for every $\varepsilon > 0$,

$$\tau_b(R; \mathfrak{a}^{t_i - \varepsilon}) \neq \tau_b(R; \mathfrak{a}^{t_i}).$$

One easy to observe fact about multiplier ideals is that their jumping numbers are discrete and rational, at least when R is \mathbb{Q} -Gorenstein and normal; see [ELSV04]. Here, by discrete we mean that the set of jumping numbers with respect to a fixed ideal have no limit points. Because of this, various groups have recently worked to show that the F -jumping numbers of the test ideal are also discrete and rational; see [Har06], [BMS08], [BMS09], [KLZ09], and [BSTZ10]. In the most recent-mentioned work, the author, along with M. Blickle, S. Takagi, and W. Zhang, showed that

Received by the editors April 8, 2010 and, in revised form, October 4, 2010.

2000 *Mathematics Subject Classification*. Primary 13A35, 14F18, 14B05.

Key words and phrases. Test ideal, jumping number, \mathbb{Q} -Gorenstein, multiplier ideal.

The author was partially supported by a National Science Foundation postdoctoral fellowship and by NSF grant DMS-1064485/0969145.

the F -jumping numbers of test ideals formed a discrete set of rational numbers when R is normal and \mathbb{Q} -Gorenstein with index not divisible by $p > 0$. Recall that the *index* of a \mathbb{Q} -Gorenstein ring R is the smallest natural number n where $\omega_R^{(n)} = \mathcal{O}_{\text{Spec } R}(nK_R)$ is locally free.

The most fundamental case left open is the case when R is \mathbb{Q} -Gorenstein but of arbitrary index; see [BSTZ10, Question 6.1]. We answer this question at least for discreteness.

Theorem 3.5. *Suppose that R is a normal domain essentially of finite type over an F -finite field. Further suppose that $\mathfrak{a} \subseteq R$ is an ideal and Δ is an \mathbb{R} -divisor on $X = \text{Spec } R$ such that $K_X + \Delta$ is \mathbb{R} -Cartier (for example, this holds if $\Delta = 0$ and R is \mathbb{Q} -Gorenstein). Then, as t varies, the F -jumping numbers of $\tau_b(R; \Delta, \mathfrak{a}^t)$ have no limit points – they are discrete.*

We also point out why the existing proofs of *rationality* do not seem to work in the case that R is \mathbb{Q} -Gorenstein with index divisible by p .

Recently, in [DH09], de Fernex and Hacon gave a definition of the multiplier ideal without the \mathbb{Q} -Gorenstein assumption and asked the question of whether discreteness and rationality of the F -jumping numbers still holds in this context. Following this, Urbinati showed that rationality need not hold but gave some evidence that discreteness may hold in general; see [Urb10]. This suggests that one should not expect rationality to hold in positive characteristic either.

2. DEFINITION OF THE TEST IDEAL

We only give a very brief description of the big test ideal in this paper. Please see [BSTZ10] for a more detailed description of the test ideal.

First we fix some notation. Given a ring R of characteristic $p > 0$ and M an R -module, we set $F_*^e M$ to be the R -module which agrees with M as an additive group but where the R -module structure is defined by the rule $r.m = r^{p^e}m$. Also recall that R is said to be *F -finite* if $F_*^e R$ is a finitely generated R -module.

Convention. Throughout this paper, all rings will be assumed to be F -finite.

Recall that an \mathbb{R} -divisor on a normal scheme X is a formal linear combination of prime Weil divisors D_i with real coefficients. An \mathbb{R} -divisor D is called \mathbb{R} -*Cartier* if it is equal to an \mathbb{R} -linear combination of Cartier divisors.

We now define the test ideal $\tau_b(R; \Delta, \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m})$. We work in this greater generality because when proving our main theorem, we perturb our initial triple $(R, \Delta, \mathfrak{a}^t)$ to a new triple $(R, \Delta', \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m})$ which has the same test ideal.

Definition 2.1 ([HH90], [Hoc07], [Sch10]). Suppose that R is an F -finite normal domain, $\Delta \geq 0$ is an \mathbb{R} -divisor on $X = \text{Spec } R$, $\mathfrak{a}, \mathfrak{b}_1, \dots, \mathfrak{b}_m \subseteq R$ are non-zero ideals and $t, s_1, \dots, s_m \geq 0$ are real numbers. Then the *big test ideal* $\tau_b(R; \Delta, \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m})$ is defined to be the unique smallest non-zero ideal $J \subseteq R$ such that

$$(1) \quad \phi \left(F_*^e (\mathfrak{a}^{\lceil t(p^e - 1) \rceil} \mathfrak{b}_1^{\lceil s_1(p^e - 1) \rceil} \dots \mathfrak{b}_m^{\lceil s_m(p^e - 1) \rceil} J) \right) \subseteq J$$

for every $e \geq 0$ and every $\phi \in \text{Hom}_R(F_*^e R(\lceil (p^e - 1)\Delta \rceil), R)$. This ideal always exists in the context described.

Remark 2.2. In the case that $K_X + \Delta$ is \mathbb{Q} -Cartier, the *big test ideal* is known to equal the (*finitistic*) *test ideal* (which we will not define here); see [Tak04] and [BSTZ10] for details.

If all $\mathfrak{b}_i = R$, then we denote the associated big test ideal by $\tau_b(R; \Delta, \mathfrak{a}^t)$. Likewise if $\Delta = 0$, then we denote the associated big test ideal using the notation $\tau_b(R; \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m})$. Finally, if the $\mathfrak{b}_i = (f_i)$ are principal, we denote the associated big test ideal by $\tau_b(R; \Delta, \mathfrak{a}^t f_1^{s_1} \dots f_m^{s_m})$

Remark 2.3. Given a non-zero element $c \in \tau_b(R; \Delta, \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m})$ (such an element is called a *big sharp test element*), we note that

$$(2) \quad \tau_b(R; \Delta, \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m}) = \sum_{e \geq 0} \sum_{\phi} \phi \left(F_*^e (c \mathfrak{a}^{\lceil t(p^e - 1) \rceil} \mathfrak{b}_1^{\lceil s_1(p^e - 1) \rceil} \dots \mathfrak{b}_m^{\lceil s_m(p^e - 1) \rceil}) \right),$$

where the inner sum is over $\phi \in \text{Hom}_R(F_*^e R(\lceil (p^e - 1)\Delta \rceil), R)$. To see this, simply note that the right side satisfies equation (1), and it is by definition the smallest ideal containing c satisfying equation (1); note $\mathfrak{a}^0 = \mathfrak{b}_i^0 = R$.

Suppose that $X = \text{Spec } R$ is normal. Then given $\phi \in \text{Hom}_R(F_*^e R, R) \cong F_*^e \mathcal{O}_X((1-p^e)K_X)$, we may view ϕ as determining an effective Weil divisor linearly equivalent to $(1-p^e)K_X$.

Definition 2.4. We use D_ϕ to denote the Weil divisor associated to ϕ in this way.

Given an \mathbb{R} -divisor $\Delta \geq 0$ on X , one has an inclusion

$$(3) \quad \text{Hom}_R(F_*^e R(\lceil (p^e - 1)\Delta \rceil), R) \subseteq \text{Hom}_R(F_*^e R, R).$$

The following lemma gives a nice interpretation of this submodule.

Lemma 2.5. *An element $\phi \in \text{Hom}_R(F_*^e R, R)$ is contained inside the submodule $\text{Hom}_R(F_*^e R(\lceil (p^e - 1)\Delta \rceil), R)$ if and only if $D_\phi \geq (p^e - 1)\Delta$.*

Proof. Because all the modules are reflexive, the statement can be reduced to the case when R is a discrete valuation ring and $\Delta = s \text{ div}(x)$, where x is the parameter for the DVR R and $s \geq 0$ is a real number. In this case, the inclusion from equation (3) can be identified with the multiplication map $R \rightarrow R$ which sends 1 to $x^{\lceil s(p^e - 1) \rceil}$. Thus, $\phi \in \text{Hom}_R(F_*^e R, R) \cong R$ is contained inside $\text{Hom}_R(F_*^e R(\lceil (p^e - 1)\Delta \rceil), R) \cong x^{\lceil s(p^e - 1) \rceil} R$ if and only if $D_\phi \geq \lceil s(p^e - 1) \rceil \text{ div}(x) = \lceil (p^e - 1)\Delta \rceil$. However, since D_ϕ is integral, it is harmless to remove the round-up $\lceil \cdot \rceil$. \square

3. DISCRETENESS OF F -JUMPING NUMBERS

In this section we prove our main result. We accomplish this by perturbing our triples $(R, \Delta, \mathfrak{a}^t)$ in order to reduce the discreteness statement to the case where the (log) \mathbb{Q} -Gorenstein index is not divisible by $p > 0$. First we need a lemma.

Lemma 3.1. *Suppose that $(X = \text{Spec } R, \Delta, \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m})$ is a triple and that $\Delta = \Gamma + b \text{ div}(f)$ for some $f \in R \setminus \{0\}$ and non-negative number $b \in \mathbb{R}$. Then*

$$\tau_b(R; \Delta, \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m}) = \tau_b(R; \Gamma, f^b \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m}).$$

This type of statement is essentially obvious for multiplier ideals, but because of certain issues surrounding the construction of test ideals we have thus-far presented, it is somewhat less obvious in this context. However, it is still quite straightforward, especially from the definition of the generalized test ideal by Hara-Yoshida-Takagi (the proof in that case uses the theory tight closure); see [HY03] and [Tak04]. We provide a short proof here, certainly acknowledging that this statement is obvious to experts.

Proof. Choose c to be a big sharp test element for both $(X, \Delta, \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m})$ and $(X, \Gamma, f^b \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m})$. Then we know that the test ideal $\tau_b(R; \Gamma, f^b \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m})$ equals

$$\begin{aligned} & \sum_{\phi, D_\phi \geq (p^e-1)\Gamma} \phi \left(F_*^e c f^{\lceil b(p^e-1) \rceil} \mathfrak{a}^{\lceil t(p^e-1) \rceil} \mathfrak{b}_1^{\lceil s_1(p^e-1) \rceil} \dots \mathfrak{b}_m^{\lceil s_m(p^e-1) \rceil} \right) \\ &= \sum_{\phi, D_\phi \geq (p^e-1)\Gamma + \text{div } f^{\lceil b(p^e-1) \rceil}} \phi \left(F_*^e c \mathfrak{a}^{\lceil t(p^e-1) \rceil} \mathfrak{b}_1^{\lceil s_1(p^e-1) \rceil} \dots \mathfrak{b}_m^{\lceil s_m(p^e-1) \rceil} \right) \\ &\subseteq \sum_{\phi, D_\phi \geq (p^e-1)\Gamma + \lceil b(p^e-1) \text{ div}(f) \rceil} \phi \left(F_*^e c \mathfrak{a}^{\lceil t(p^e-1) \rceil} \mathfrak{b}_1^{\lceil s_1(p^e-1) \rceil} \dots \mathfrak{b}_m^{\lceil s_m(p^e-1) \rceil} \right) \\ &\subseteq \sum_{\phi, D_\phi \geq (p^e-1)\Gamma + b(p^e-1) \text{ div}(f)} \phi \left(F_*^e c \mathfrak{a}^{\lceil t(p^e-1) \rceil} \mathfrak{b}_1^{\lceil s_1(p^e-1) \rceil} \dots \mathfrak{b}_m^{\lceil s_m(p^e-1) \rceil} \right) \\ &= \tau_b(R; \Delta, \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m}), \end{aligned}$$

and so $\tau_b(R; \Gamma, f^b \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m}) \subseteq \tau_b(R; \Delta, \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m})$. For the converse inclusion, observe first that

$$\text{div}(f^{\lceil b(p^e-1) \rceil}) - b(p^e-1) \text{ div}(f) \leq \text{div}(f).$$

Thus, since cf is also a test element,

$$\begin{aligned} \tau_b(R, \Delta, \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m}) &= \sum_{\phi, D_\phi \geq (p^e-1)\Delta} \phi \left(F_*^e c f \mathfrak{a}^{\lceil t(p^e-1) \rceil} \mathfrak{b}_1^{\lceil s_1(p^e-1) \rceil} \dots \mathfrak{b}_m^{\lceil s_m(p^e-1) \rceil} \right) \\ &= \sum_{\phi, D_\phi \geq (p^e-1)\Gamma + b(p^e-1) \text{ div}(f)} \phi \left(F_*^e c f \mathfrak{a}^{\lceil t(p^e-1) \rceil} \mathfrak{b}_1^{\lceil s_1(p^e-1) \rceil} \dots \mathfrak{b}_m^{\lceil s_m(p^e-1) \rceil} \right) \\ &= \sum_{\phi, D_\phi \geq (p^e-1)\Gamma + b(p^e-1) \text{ div}(f) + \text{div}(f)} \phi \left(F_*^e c \mathfrak{a}^{\lceil t(p^e-1) \rceil} \mathfrak{b}_1^{\lceil s_1(p^e-1) \rceil} \dots \mathfrak{b}_m^{\lceil s_m(p^e-1) \rceil} \right) \\ &\subseteq \sum_{\phi, D_\phi \geq (p^e-1)\Gamma + \text{div}(f^{\lceil b(p^e-1) \rceil})} \phi \left(F_*^e c \mathfrak{a}^{\lceil t(p^e-1) \rceil} \mathfrak{b}_1^{\lceil s_1(p^e-1) \rceil} \dots \mathfrak{b}_m^{\lceil s_m(p^e-1) \rceil} \right) \\ &= \sum_{\phi, D_\phi \geq (p^e-1)\Gamma} \phi \left(F_*^e c f^{\lceil b(p^e-1) \rceil} \mathfrak{a}^{\lceil t(p^e-1) \rceil} \mathfrak{b}_1^{\lceil s_1(p^e-1) \rceil} \dots \mathfrak{b}_m^{\lceil s_m(p^e-1) \rceil} \right) \\ &= \tau_b(R, \Gamma, f^b \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m}), \end{aligned}$$

and so $\tau_b(R; \Delta, \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m}) \subseteq \tau_b(R; \Gamma, f^b \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m})$, as desired. \square

We also need a very special case of Skoda's theorem.

Lemma 3.2 ([HT04, Theorem 4.1]). *Suppose that $X = \text{Spec } R$, $\Delta > 0$, $\mathfrak{a} \subseteq R$ and $t \geq 0$ is as above. Further suppose that $f \in R$ is a non-zero element. Then*

$$\tau_b(X; \Delta + \text{div}(f), \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m}) = f \tau_b(X; \Delta, \mathfrak{a}^t \mathfrak{b}_1^{s_1} \dots \mathfrak{b}_m^{s_m}).$$

Proof. We leave the proof to reader; see [HT04, Theorem 4.2] and [BSTZ10, Lemma 3.26]. \square

Now we can prove the following result.

Theorem 3.3. *Suppose that R is an F -finite normal domain and further suppose that $(X = \text{Spec } R, \Delta, \mathfrak{a}^t)$ is a triple where $K_X + \Delta$ is \mathbb{R} -Cartier. Then for each point $x \in X$, there exists an open set $U = \text{Spec } R' = \text{Spec } R[h^{-1}]$ containing $x \in X$*

with the following properties: There exists an effective \mathbb{Q} -divisor Γ on U , elements $f_1, \dots, f_m \in R' \setminus \{0\}$ and non-negative real numbers b_1, \dots, b_m such that

- (1) $K_U + \Gamma$ is \mathbb{Q} -Cartier with index not divisible by $p > 0$, and, furthermore, $(p^e - 1)(K_U + \Gamma) \sim 0$ for some integer $e > 0$.
- (2) The F -jumping numbers of $\tau_b(U, \Gamma, f_1^{b_1} \dots f_m^{b_m} (\mathfrak{a}R')^t)$ are the same as the F -jumping numbers of $\tau_b(U, \Delta|_U, (\mathfrak{a}R')^t)$ (both sets of jumping numbers are with respect to t).

Proof. Choose a non-zero section ϕ of $\text{Hom}_R(F_*^e R, R)$ and set $\Gamma = \frac{1}{p^e - 1} D_\phi$. It follows that $K_X + \Gamma$ satisfies condition (1) on X . Therefore, $(K_X + \Delta) - (K_X + \Gamma) = \Delta - \Gamma$ is \mathbb{R} -Cartier, and so we may write $\Delta - \Gamma = d_1 D_1 + \dots + d_m D_m$ for some integral effective Cartier divisors D_i and real numbers $d_i \in \mathbb{R}$. We choose our open set $U = \text{Spec } R[h^{-1}] = \text{Spec } R'$ to be any such set containing $x \in X$ where all of the $D_i|_U$ are principal divisors.

Now write $D_i|_U = \text{div}(f_i)$ for some $f_i \in R' \setminus 0$ and also by abuse of notation denote $\Gamma := \Gamma|_U$. Choose natural numbers l_i such that $b_i := l_i + d_i > 0$ for all i and set $g := f_1^{l_1} \dots f_m^{l_m} \in R'$. Notice that $(\Delta|_U + \text{div}(g)) - \Gamma = b_1 \text{div}(f_1) + \dots + b_m \text{div}(f_m)$.

By Lemma 3.2, the F -jumping numbers of $\tau_b(U; \Delta|_U, (\mathfrak{a}R')^t)$ and the F -jumping numbers of $\tau_b(U; \Delta|_U + \text{div}(g), (\mathfrak{a}R')^t)$ coincide. Now using Lemma 3.1 we have

$$\begin{aligned} & \tau_b(U; \Delta|_U + \text{div}(g), (\mathfrak{a}R')^t) \\ &= \tau_b(U; \Gamma + b_1 \text{div}(f_1) + \dots + b_m \text{div}(f_m), (\mathfrak{a}R')^t) \\ &= \tau_b(U; \Gamma, f_1^{b_1} \dots f_m^{b_m} (\mathfrak{a}R')^t), \end{aligned}$$

which proves the theorem. \square

Remark 3.4. If, in Theorem 3.3, $K_X + \Delta$ is \mathbb{Q} -Cartier, then one needs only a single $f_1^{b_1}$ (and no other $f_i^{b_i}$). However, if the index of $K_X + \Delta$ is divisible by $p > 0$, then it follows by construction that b_1 will be a rational number with its denominator divisible by $p > 0$.

We are now in a position to prove the discreteness of the F -jumping numbers in the case that X is essentially of finite type over a field. The proof idea follows the usual lines.

Theorem 3.5. Suppose that R is a normal domain essentially of finite type over an F -finite field. Further suppose that $\mathfrak{a} \subseteq R$ is an ideal and Δ is an \mathbb{R} -divisor on $X = \text{Spec } R$ such that $K_X + \Delta$ is \mathbb{R} -Cartier (for example, this holds if $\Delta = 0$ and R is \mathbb{Q} -Gorenstein). Then, as t varies, the F -jumping numbers of $\tau_b(R; \Delta, \mathfrak{a}^t)$ have no limit points – they are discrete.

Proof. By [BSTZ10, Proposition 3.28], it is sufficient to answer this question on a finite affine cover of X . Therefore, we reduce to the case that X is one of the charts from Theorem 3.3. In particular, it is sufficient to prove our result for triples of the form $\tau_b(R; \Gamma, f_1^{b_1} \dots f_m^{b_m} \mathfrak{a}^t)$, where $(p^e - 1)(K_X + \Gamma) \sim 0$ for some $e > 0$. Using [BSTZ10, Lemma 4.2, Proposition 3.28], one can further assume that R is of finite type over an F -finite field of characteristic $p > 0$. One then has two options:

- (a) Mimic the proof of the main result of [BSTZ10, Section 4]. In other words, use the methods of F -adjunction (as worked out in [Sch09a] and [BSTZ10]) to reduce to the case where R is a polynomial ring and then use

degree bounding methods similar to those found in [BMS08]. Note that in [BSTZ10] one worked with triples $(R, \Delta, \mathfrak{a}^t)$ and not with the more complicated objects $(R, \Gamma, f_1^{b_1} \dots f_m^{b_m} \mathfrak{a}^t)$, but the methods are easily generalized to our setting.

- (b) Use the new language of [Bli09, Section 4]. We claim that the algebra of p^{-e} -linear maps associated to the triple $(R, \Gamma, f_1^{b_1} \dots f_m^{b_m})$, as in [Sch09b, Remark 3.10], is “gauge bounded” (see [Bli09, Definition 4.7]). To see this claim, note that by [Sch09a, Lemma 3.9] or [Sch09b, Remark 4.4] the Cartier-algebra associated to (R, Γ) is finitely generated and thus gauge bounded by [Bli09, Proposition 4.8]. It then follows from [Bli09, Proposition 4.13] that the Cartier-algebra associated to $(R, \Gamma, f_1^{b_1} \dots f_m^{b_m})$ is also gauge bounded, as claimed. To finish the proof, apply [Bli09, Theorem 4.14].

In either case, the result follows easily from the theories previously developed. \square

4. ON THE QUESTION OF RATIONALITY

Note that the usual way to prove the rationality of the F -jumping numbers employs the following theorem. First recall that a pair (X, Δ) is called *log \mathbb{Q} -Gorenstein with index n* if $n(K_X + \Delta)$ is Cartier and $n > 0$ is the smallest integer with this property.

Theorem 4.1 ([BMS08], [BSTZ10]). *Suppose (X, Δ) is log \mathbb{Q} -Gorenstein with index n such that n divides $(p^e - 1)$ for some fixed $e > 0$. Further suppose that \mathfrak{a} is an ideal sheaf of X . Then if t_0 is an F -jumping number of $\tau(X; \Delta, \mathfrak{a}^t)$, then $p^e t_0$ is also an F -jumping number.*

However, without the “index not divisible by p ” assumption, this theorem is false. Consider the following example (which in some sense is typical by Remark 3.4).

Example 4.2. Set $X = \mathbb{A}_k^1 = \text{Spec } k[x]$, $\Delta = \frac{1}{p} \text{div}(x)$ and $\mathfrak{a} = (x)$. Then the F -jumping numbers of $(X, \Delta, \mathfrak{a}^t) = (X, (x)^{1/p} \mathfrak{a}^t)$ with respect to t are

$$\frac{p-1}{p}, \frac{2p-1}{p}, \frac{3p-1}{p}, \dots$$

In particular, p (or p^e) times any of them is not an F -jumping number.

ACKNOWLEDGEMENTS

The author would like to thank Manuel Blickle and Kevin Tucker for several stimulating and valuable conversations. The author would also like to thank the referee for numerous helpful suggestions.

REFERENCES

- [Bli09] M. Blickle: *Test ideals via algebras of p^{-e} -linear maps*, arXiv:0912.2255. To appear in the Journal of Algebraic Geometry.
- [BMS08] M. Blickle, M. Mustaţă, and K. Smith: *Discreteness and rationality of F -thresholds*, Michigan Math. J. **57** (2008), 43–61. MR2492440 (2010c:13003)
- [BMS09] M. Blickle, M. Mustaţă, and K. E. Smith: *F -thresholds of hypersurfaces*, Trans. Amer. Math. Soc. **361** (2009), no. 12, 6549–6565. MR2538604 (2011a:13006)
- [BSTZ10] M. Blickle, K. Schwede, S. Takagi, and W. Zhang: *Discreteness and rationality of F -jumping numbers on singular varieties*, Math. Ann. **347** (2010), no. 4, 917–949. MR2658149

- [DH09] T. De Fernex and C. Hacon: *Singularities on normal varieties*, Compos. Math. **145** (2009), no. 2, 393–414. MR2501423 (2010c:14013)
- [ELSV04] L. Ein, R. Lazarsfeld, K. E. Smith, and D. Varolin: *Jumping coefficients of multiplier ideals*, Duke Math. J. **123** (2004), no. 3, 469–506. MR2068967 (2005k:14004)
- [Har06] N. Hara: *F -pure thresholds and F -jumping exponents in dimension two*, Math. Res. Lett. **13** (2006), no. 5–6, 747–760. With an appendix by Paul Monsky. MR2280772 (2007m:14032)
- [HT04] N. Hara and S. Takagi: *On a generalization of test ideals*, Nagoya Math. J. **175** (2004), 59–74. MR2085311 (2005g:13009)
- [HY03] N. Hara and K.-I. Yoshida: *A generalization of tight closure and multiplier ideals*, Trans. Amer. Math. Soc. **355** (2003), no. 8, 3143–3174 (electronic). MR1974679 (2004i:13003)
- [Hoc07] M. Hochster: *Foundations of tight closure theory*, lecture notes from a course taught at the University of Michigan, fall 2007.
- [HH90] M. Hochster and C. Huneke: *Tight closure, invariant theory, and the Briançon-Skoda theorem*, J. Amer. Math. Soc. **3** (1990), no. 1, 31–116. MR1017784 (91g:13010)
- [KLZ09] M. Katzman, G. Lyubeznik, and W. Zhang: *On the discreteness and rationality of F -jumping coefficients*, J. Algebra **322** (2009), no. 9, 3238–3247. MR2567418 (2011c:13005)
- [MTW05] M. Mustaţă, S. Takagi, and K. Watanabe: *F -thresholds and Bernstein-Sato polynomials*, European Congress of Mathematics, Eur. Math. Soc., Zürich, 2005, pp. 341–364. MR2185754 (2007b:13010)
- [MY09] M. Mustaţă and K.-I. Yoshida: *Test ideals vs. multiplier ideals*, Nagoya Math. J. **193** (2009), 111–128. MR2502910 (2010a:13010)
- [Sch09a] K. Schwede: *F -adjunction*, Algebra Number Theory **3** (2009), no. 8, 907–950. MR2587408 (2011b:14006)
- [Sch09b] K. Schwede: *Test ideals in non- Q -Gorenstein rings*, arXiv:0906.4313, to appear in Trans. Amer. Math. Soc.
- [Sch10] K. Schwede: *Centers of F -purity*, Math. Z. **265** (2010), no. 3, 687–714. MR2644316 (2011e:13011)
- [Tak04] S. Takagi: *An interpretation of multiplier ideals via tight closure*, J. Algebraic Geom. **13** (2004), no. 2, 393–415. MR2047704 (2005c:13002)
- [Tak06] S. Takagi: *Formulas for multiplier ideals on singular varieties*, Amer. J. Math. **128** (2006), no. 6, 1345–1362. MR2275023 (2007i:14006)
- [Urb10] S. Urbinati: *Discrepancies of non- \mathbb{Q} -Gorenstein varieties*, arXiv:1001.2930. To appear in the Michigan Mathematical Journal.

DEPARTMENT OF MATHEMATICS, THE PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK,
PENNSYLVANIA 16802

E-mail address: schwede@math.psu.edu