

## EIGENFUNCTION EXPANSIONS IN $\mathbb{R}^n$

TODOR GRAMCHEV, STEVAN PILIPOVIC, AND LUIGI RODINO

(Communicated by Richard Rochberg)

ABSTRACT. The main goal of this paper is to extend in  $\mathbb{R}^n$  a result of Seeley on eigenfunction expansions of real analytic functions on compact manifolds. As a counterpart of an elliptic operator in a compact manifold, we consider in  $\mathbb{R}^n$  a selfadjoint, globally elliptic Shubin type differential operator with spectrum consisting of a sequence of eigenvalues  $\lambda_j, j \in \mathbb{N}$ , and a corresponding sequence of eigenfunctions  $u_j, j \in \mathbb{N}$ , forming an orthonormal basis of  $L^2(\mathbb{R}^n)$ . Elements of Schwartz  $\mathcal{S}(\mathbb{R}^n)$ , resp. Gelfand-Shilov  $S_{1/2}^{1/2}$  spaces, are characterized through expansions  $\sum_j a_j u_j$  and the estimates of coefficients  $a_j$  by the power function, resp. exponential function of  $\lambda_j$ .

### 1. INTRODUCTION AND STATEMENT OF THE RESULT

We shall give a version in  $\mathbb{R}^n$  of some results, already known on compact manifolds, concerning eigenfunction expansions. Broadly speaking, the aim is to relate the regularity of a function with the decay properties of the sequence of the Fourier coefficients. More precisely, we want to reproduce in  $\mathbb{R}^n$  the classical results of [15], Section 10, for Sobolev regularity and [16] for analytic functions on a compact manifold taking into account Weyl asymptotics for eigenvalues.

Our basic example of an operator will be the harmonic oscillator appearing in Quantum Mechanics,

$$(1.1) \quad H = -\Delta + |x|^2,$$

whose eigenfunctions are the Hermite functions

$$(1.2) \quad h_\alpha(x) = H_\alpha(x) e^{-|x|^2/2}, \quad \alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n,$$

where  $H_\alpha(x)$  is the  $\alpha$ -th Hermite polynomial; cf. [5]. See for example [14], [13], [11], for related Hermite expansions as well as [7], [18] for connections with a degenerate harmonic oscillator.

---

Received by the editors October 11, 2010.

2010 *Mathematics Subject Classification.* Primary 35S05; Secondary 46F05, 35B65.

*Key words and phrases.* Shubin-type operators, Gelfand–Shilov spaces.

The first author was partially supported by a PRIN project of MIUR, Italy and GNAMPA, INDAM.

The second author was supported by the project 144016, Serbia.

The third author was partially supported by a PRIN project of MIUR, Italy and GNAMPA, INDAM.

Here we shall consider a more general class of operators with polynomial coefficients in  $\mathbb{R}^n$ :

$$(1.3) \quad P = \sum_{|\alpha|+|\beta|\leq m} c_{\alpha\beta}x^\beta D_x^\alpha, \quad D^\alpha = (-i)^{|\alpha|}\partial_x^\alpha,$$

studied by Shubin [17] in the frame of a global pseudo-differential calculus; see also [8], [1], [12]. Let us recall, in short, some definitions and results from [17], Chapter IV.

First, global ellipticity for  $P$  in (1.3) is defined by imposing

$$(1.4) \quad p_m(x, \xi) = \sum_{|\alpha|+|\beta|=m} c_{\alpha\beta}x^\beta \xi^\alpha \neq 0 \quad \text{for } (x, \xi) \neq (0, 0).$$

This condition is obviously satisfied by  $H$  in (1.1). For these operators, the counterpart of the standard Sobolev spaces comprises the spaces

$$(1.5) \quad Q^s(\mathbb{R}^n) = \{u \in \mathcal{S}'(\mathbb{R}^n), \|u\|_{Q^s} := \sum_{|\alpha|+|\beta|\leq s} \|x^\beta \partial_x^\alpha u\|_{L^2(\mathbb{R}^n)} < +\infty\},$$

where  $\mathcal{S}'(\mathbb{R}^n)$  is the class of the tempered distributions of Schwartz and  $s \in \mathbb{N}$ . Under the global ellipticity assumption (1.4),  $P : Q^m(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$  is a Fredholm operator. The finite-dimensional null-space  $\text{Ker } P$  is given by functions in the Schwartz space  $\mathcal{S}(\mathbb{R}^n)$ .

Following Gelfand and Shilov, it is natural to consider as a global counterpart in  $\mathbb{R}^n$  of the real analytic class the inductive (respectively, projective) Gelfand-Shilov classes  $S_\nu^\mu(\mathbb{R}^n)$  (respectively,  $\Sigma_\nu^\mu(\mathbb{R}^n)$ ),  $\mu > 0$ ,  $\nu > 0$ ,  $\mu + \nu \geq 1$  (respectively  $\mu + \nu > 1$ ), defined as the set of all  $u \in \mathcal{S}(\mathbb{R}^n)$  for which there exist  $A > 0, C > 0$  (respectively, for every  $A > 0$  there exists  $C > 0$ ) such that

$$|x^\beta \partial_x^\alpha u(x)| \leq CA^{-|\alpha|-|\beta|}(\alpha!)^\mu(\beta!)^\nu, \quad \alpha \in \mathbb{N}^n, \beta \in \mathbb{N}^n;$$

see [6], [11] and Chapter 6 in [12]. In the sequel we shall limit our attention to  $S_\mu^\mu(\mathbb{R}^n)$ ,  $\mu \geq 1/2$  (respectively,  $\Sigma_\mu^\mu(\mathbb{R}^n)$ ,  $\mu > \frac{1}{2}$ ). We recall that  $u \in S_\mu^\mu(\mathbb{R}^n)$  iff there exist  $A > 0, C > 0$  (respectively, for every  $A > 0$  one can find  $C > 0$ ) such that

$$(1.6) \quad \sum_{|\alpha|+|\beta|=s} \|x^\beta \partial_x^\alpha u\|_{L^2(\mathbb{R}^n)} \leq CA^{-s}(s!)^\mu, \quad s \in \mathbb{N}.$$

It was shown recently that every solution  $u \in \mathcal{S}'(\mathbb{R}^n)$  of  $Pu = 0$  belongs to  $S_{1/2}^{1/2}(\mathbb{R}^n)$  provided (1.4) holds; see [3], [4] for details and more general results.

We assume, as in Seeley [16], that  $P$  is a normal operator (i.e.,  $P^*P = PP^*$ ) satisfying the global ellipticity condition (1.4). This guarantees the existence of a basis of orthonormal eigenfunctions  $u_j$ ,  $j \in \mathbb{N}$ , with eigenvalues  $\lambda_j$ ,  $\lim_{j \rightarrow \infty} |\lambda_j| = +\infty$  (see Seeley [16] and Shubin [17]). Hence, given  $u \in L^2(\mathbb{R}^n)$  or  $u \in \mathcal{S}'(\mathbb{R}^n)$ , we can expand

$$(1.7) \quad u = \sum_{j=1}^\infty a_j u_j,$$

where the Fourier coefficient  $a_j \in \mathbb{C}$  is defined by

$$(1.8) \quad a_j = (u, u_j)_{L^2(\mathbb{R}^n)}, \quad j = 1, 2, \dots,$$

with convergence in  $L^2(\mathbb{R}^n)$  or  $\mathcal{S}'(\mathbb{R}^n)$  for (1.7).

In view of [3] the eigenfunctions  $u_j$  belong to  $S_{1/2}^{1/2}(\mathbb{R}^n)$ .

We state the first main result.

**Theorem 1.1.** *Suppose that  $P$  is globally elliptic (cf. (1.3), (1.4)) and normal. Then:*

- (i)  $u \in Q^s(\mathbb{R}^n) \iff \sum_{j=1}^{\infty} |a_j|^2 |\lambda_j|^{2s/m} < \infty \iff \sum_{j=1}^{\infty} |a_j|^2 j^{s/n} < \infty, s \in \mathbb{N}.$
- (ii)  $u \in \mathcal{S}(\mathbb{R}^n) \iff |a_j| = O(|\lambda_j|^{-s}), j \rightarrow \infty \iff |a_j| = O(j^{-s}), j \rightarrow \infty$  for all  $s \in \mathbb{N}.$

Next, we show the global analogue to Seeley’s theorem in [16].

**Theorem 1.2.** *Let  $P$  be as before and let  $\mu \geq 1/2$  (respectively,  $\mu > 1/2$ ). Then we have:*

$$u \in S_{\mu}^{\mu}(\mathbb{R}^n) \iff \sum_{j=1}^{\infty} |a_j|^2 e^{\epsilon |\lambda_j|^{1/(m\mu)}} < \infty$$

for some  $\epsilon > 0 \iff \sum_{j=1}^{\infty} |a_j|^2 e^{\epsilon j^{1/(m\mu)}} < \infty$  for some  $\epsilon > 0 \iff$  there exist  $C > 0, \epsilon > 0$  such that

$$(1.9) \quad |a_j| \leq C e^{-\epsilon j^{1/(2n\mu)}}, \quad j \in \mathbb{N}$$

(respectively,  $u \in \Sigma_{\mu}^{\mu}(\mathbb{R}^n) \iff \sum_{j=1}^{\infty} |a_j|^2 e^{\epsilon |\lambda_j|^{1/(m\mu)}} < \infty$  for all  $\epsilon > 0 \iff \sum_{j=1}^{\infty} |a_j|^2 e^{\epsilon j^{1/(m\mu)}} < \infty$  for all  $\epsilon > 0 \iff$  for every  $\epsilon > 0$  there exist  $C > 0$  such that

$$(1.10) \quad |a_j| \leq C e^{-\epsilon j^{1/(2n\mu)}}, \quad j \in \mathbb{N}.$$

*Remark 1.3.* Choosing as  $P$  the harmonic oscillator  $H$  in (1.1), with eigenfunctions  $h_{\alpha}(x)$  as in (1.2), we recapture the results on the Hermite expansions related to Gelfand-Shilov type spaces for  $n = 1$  whereas for  $n \geq 2$ , taking into account the multiplicity of the eigenvalue  $\lambda_{\alpha} = \sum_{j=1}^n (2\alpha_j + 1)$  for  $h_{\alpha}, \alpha \in \mathbb{N}^n$ , we obtain as a particular case of Theorem 1.2 the characterization:  $u \in S_{1/2}^{1/2}(\mathbb{R}^n)$  iff

$$|a_{\alpha}| \leq C e^{-\epsilon |\alpha|}, \quad \alpha \in \mathbb{N}^n,$$

for positive constants  $C$  and  $\epsilon$ , where  $u = \sum_{\alpha \in \mathbb{N}^n} a_{\alpha} h_{\alpha}$ ; cf. [11], [13] and the references therein.

## 2. PROOF OF THEOREM 1.1

It is not restrictive to assume that  $P$  is positive, with  $\lambda_j > 0$ ; cf. [16]. We need some preliminary results from [17]. Namely, concerning asymptotics of eigenvalues, Theorem 30.1 in [17] and Proposition 4.6.4 in [12] give the following lemma.

**Lemma 2.1.** *Let  $P$  be globally elliptic of order  $m > 0$ , cf. (1.3), (1.4), and strictly positive. Then for the eigenvalues  $\lambda_j, j = 1, 2, \dots$ , we have*

$$\lambda_j \sim C j^{m/(2n)} \quad \text{as } j \rightarrow +\infty,$$

for a positive constant  $C$ .

Now, for  $P$  as before and  $r \in \mathbb{R}, r \neq 0$ , introduce the  $r$ -th power

$$(2.1) \quad P^r u = \sum_{j=1}^{\infty} \lambda_j^r a_j u_j,$$

with  $a_j, u_j$  as in (1.7), (1.8). The operator  $P^r$  is well defined as a map  $\mathcal{S}'(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$ .

**Lemma 2.2.** *Let  $u \in \mathcal{S}'(\mathbb{R}^n)$ . Then  $u \in Q^s(\mathbb{R}^n)$  if and only if  $P^{s/m}u \in L^2(\mathbb{R}^n)$ ,  $s \in \mathbb{N}$ . The norms  $\|u\|_{Q^s}$  and  $\|P^{s/m}u\|_{L^2(\mathbb{R}^n)}$  are equivalent.*

In fact,  $P^{s/m}$  is an elliptic operator of order  $s$  in the pseudo-differential calculus of [17], cf. Section 4.3 in [12], and consequently  $P^{s/m}u \in L^2(\mathbb{R}^n)$  corresponds to  $u \in Q^s(\mathbb{R}^n)$  with equivalence of norms; cf. Proposition 2.1.9 and Theorem 2.1.12 in [12].

We may now prove (i) in Theorem 1.1. From (2.1) and the Parseval identity:

$$\|P^{s/m}u\|_{L^2(\mathbb{R}^n)}^2 = \left\| \sum_{j=1}^{\infty} \lambda_j^{s/m} a_j u_j \right\|_{L^2(\mathbb{R}^n)}^2 = \sum_{j=1}^{\infty} \lambda_j^{2s/m} |a_j|^2.$$

In view of Lemma 2.1 we have  $\lambda_j^{2s/m} \sim C' j^{s/n}$ , and therefore from Lemma 2.2:

$$c_1 \|u\|_{Q^s}^2 \leq \sum_{j=1}^{\infty} j^{s/n} |a_j|^2 \leq c_2 \|u\|_{Q^s}^2$$

for suitable positive constants  $c_1, c_2$ . This gives (i). On the other hand,  $\mathcal{S}(\mathbb{R}^n) = \bigcap_{s \in \mathbb{N}} Q^s(\mathbb{R}^n)$ ; hence (ii) follows from (i). Note also that by Lemma 2.2 we may generalize the definition of  $Q^s(\mathbb{R}^n)$  to all  $s \in \mathbb{R}$ , and (i) extends obviously to these spaces. Finally, we observe that the preceding arguments and the statement of Theorem 1.1 remain valid for any globally elliptic normal pseudo-differential operator in [17].

### 3. PROOF OF THEOREM 1.2

We shall follow the argument of [16], pages 737-738. Namely, we shall use the following adapted version of the celebrated theorem of the iterates of [10].

**Lemma 3.1.** *Let  $P$  be globally elliptic (cf. (1.3), (1.4)) of order  $m$ . Let  $\mu \geq 1/2$  and  $u \in \mathcal{S}'(\mathbb{R}^n)$ . Then  $u \in S_\mu^\mu(\mathbb{R}^n)$  if and only if for some  $C > 0$ ,*

$$(3.1) \quad \|P^M u\|_{L^2(\mathbb{R}^n)} \leq C^{M+1} (M!)^{\mu m} \quad \text{for all } M \in \mathbb{N}.$$

A short proof of Lemma 3.1 will be given in Section 4; for more details, we refer to the forthcoming paper [2]. By applying Lemma 3.1, in the sequel we may then take the estimates (3.1) as an equivalent definition of the class  $S_\mu^\mu(\mathbb{R}^n)$ , fixing as  $P$  the operator in Theorem 1.1. On the other hand, assuming without loss of generality that  $u \in \mathcal{S}(\mathbb{R}^n)$ , we have:

$$\|P^M u\|_{L^2(\mathbb{R}^n)}^2 = \left\| \sum_{j=1}^{\infty} a_j P^M u_j \right\|_{L^2(\mathbb{R}^n)}^2 = \left\| \sum_{j=1}^{\infty} \lambda_j^M a_j u_j \right\|_{L^2(\mathbb{R}^n)}^2 = \sum_{j=1}^{\infty} \lambda_j^{2M} |a_j|^2,$$

in view of (1.7), (1.8) and Parseval's identity. It follows from Lemma 2.1 that

$$(3.2) \quad C_1 \|P^M u\|_{L^2(\mathbb{R}^n)}^2 \leq \sum_{j=1}^{\infty} j^{mM/n} |a_j|^2 \leq C_2 \|P^M u\|_{L^2(\mathbb{R}^n)}^2$$

for suitable positive constants  $C_1, C_2$ . Assume now that the estimate (1.9) is satisfied. Then from the first estimate in (3.2) we have for some  $C > 0, \epsilon > 0$ ,

$$(3.3) \quad \|P^M u\|_{L^2(\mathbb{R}^n)}^2 \leq C \sum_{j=1}^{\infty} j^{mM/n} e^{-2\epsilon j^{1/(2n\mu)}} \leq \tilde{C} \sup_j j^{mM/n} e^{-\epsilon j^{1/(2n\mu)}}$$

with

$$\tilde{C} = C \sum_{j=1}^{\infty} e^{-\epsilon j^{1/(2n\mu)}}.$$

Now observe that the identity

$$e^{\omega j^{1/(2n\mu)}} = \sum_{M=0}^{\infty} \frac{\omega^M j^{M/(2n\mu)}}{M!}$$

implies that for any  $\omega > 0$  and  $M \in \mathbb{N}$ :

$$(3.4) \quad j^{M/(2n\mu)} e^{-\omega j^{1/(2n\mu)}} \leq \omega^{-M} M!.$$

Taking the  $2\mu m$ -th power of both sides of (3.4) and applying it in the last estimate in (3.3) with  $\epsilon = 2\mu m\omega$ , we obtain

$$\|P^M u\|_{L^2(\mathbb{R}^n)}^2 \leq \tilde{C} (\omega^{-M} M!)^{2\mu m},$$

which gives (3.1) for some  $C > 0$ . Similarly, assuming (3.1) and using the second estimate in (3.2), we deduce (1.9). The same computations give the other equivalences in Theorem 1.2.

#### 4. PROOF OF LEMMA 3.1

We shall use the estimates (1.6) as the definition of  $S_\mu^\mu(\mathbb{R}^n)$ . It is then easy to show that  $u \in S_\mu^\mu(\mathbb{R}^n)$  implies (3.1). In the opposite direction, we assume (3.1) and prove (1.6). Write for short

$$(4.1) \quad |u|_s = \sum_{|\alpha|+|\beta|=s} \|x^\beta \partial_x^\alpha u\|_{L^2(\mathbb{R}^n)}.$$

The following interpolation result for the semi-norms  $|u|_s$  is needed in the case when  $m \geq 2$ , the integer  $m$  being the order of  $P$ .

**Proposition 4.1.** *There exists a constant  $C > 0$  such that for any  $s \in \mathbb{N}$ , with  $s = pm + r, p \in \mathbb{N}, 0 < r < m$ , and for all  $\epsilon > 0$ ,*

$$(4.2) \quad |u|_s \leq \epsilon |u|_{(p+1)m} + C \epsilon^{-\frac{r}{m-r}} |u|_{pm} + C^s (s!)^{1/2} \|u\|_{L^2(\mathbb{R}^n)}.$$

The proof of Proposition 4.1 is omitted for brevity. A corresponding result for the homogeneous Sobolev spaces is well known; see for example [10], Lemma 3.3, and subsequent remarks. A novelty with respect to Sobolev spaces is the last term on the right-hand side of (4.2): the factor  $(s!)^{1/2}$  comes from the symbolic calculus of [17], Section 24; see also [12], Sections 1.7, 1.8.

Since  $\mu \geq 1/2$ , Proposition 4.1, with  $\epsilon = 1$ , say, implies that we may limit ourselves to checking (1.6) for  $s = pm, p = 0, 1, \dots$ . Namely, we shall prove that the sequence

$$(4.3) \quad \sigma_p(u, \lambda) = (pm)!^{-\mu} \lambda^{-p} |u|_{pm}, \quad p = 0, 1, \dots,$$

is bounded if  $\lambda$  is sufficiently large. To this end, we use the following proposition.

**Proposition 4.2.** *Let  $P$  be as in Lemma 3.1. Then for  $\lambda > 0$  large enough, we have for all  $p \in \mathbb{N}$ :*

$$(4.4) \quad \sigma_{p+1}(u, \lambda) \leq [(pm + 1) \dots (pm + m)]^{-\mu} \sigma_p(Pu, \lambda) + \sigma_p(u, \lambda) + \sigma_{p-1}(u, \lambda) + \sigma_0(u, \lambda).$$

In Propositions 4.1, 4.2 and in the sequel we may assume  $u \in \mathcal{S}(\mathbb{R}^n)$ . Note that  $\sigma_0(u, \lambda) = |u|_0 = \|u\|_{L^2(\mathbb{R}^n)}$ .

**Proof of Proposition 4.2.** We recall that  $P : Q^m(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$  is Fredholm. Assuming  $\text{Ker } P = \{0\}$  for simplicity, we have for a constant  $C > 0$ :

$$(4.5) \quad \|u\|_{Q^s} = \sum_{|\alpha|+|\beta| \leq s} \|x^\beta D_x^\alpha u\|_{L^2(\mathbb{R}^n)} \leq C \|Pu\|_{L^2(\mathbb{R}^n)}, \quad u \in \mathcal{S}(\mathbb{R}^n).$$

Consider then for  $|\alpha| + |\beta| = (p + 1)m$  the term  $\|x^\beta D_x^\alpha u\|_{L^2(\mathbb{R}^n)}$  and write

$$x^\beta D_x^\alpha u = x^{\beta-\delta} x^\delta D_x^{\alpha-\gamma} D_x^\gamma u,$$

where we fix  $\gamma \leq \alpha, \delta \leq \beta$  so that  $|\gamma| + |\delta| = pm$  and  $|\alpha - \gamma| + |\beta - \delta| = m$ . Therefore we may estimate, using (4.5):

$$(4.6) \quad \begin{aligned} \|x^\beta D_x^\alpha u\|_{L^2(\mathbb{R}^n)} &\leq \|x^{\beta-\delta} D_x^{\alpha-\gamma} (x^\delta D_x^\gamma u)\|_{L^2(\mathbb{R}^n)} + \|x^{\beta-\delta} [x^\delta, D_x^{\alpha-\gamma}] D_x^\gamma u\|_{L^2(\mathbb{R}^n)} \\ &\leq C \|P(x^\delta D_x^\gamma u)\|_{L^2(\mathbb{R}^n)} + \|x^{\beta-\delta} [x^\delta, D_x^{\alpha-\gamma}] D_x^\gamma u\|_{L^2(\mathbb{R}^n)} \\ &\leq I_1 + I_2 + I_3 \end{aligned}$$

with

$$(4.7) \quad \begin{aligned} I_1 &= C \|x^\delta D_x^\gamma (Pu)\|_{L^2(\mathbb{R}^n)}, \quad I_2 = C \|[P, x^\delta D_x^\gamma] u\|_{L^2(\mathbb{R}^n)}, \\ I_3 &= \|x^{\beta-\delta} [x^\delta, D_x^{\alpha-\gamma}] D_x^\gamma u\|_{L^2(\mathbb{R}^n)}. \end{aligned}$$

Summing up  $I_1, I_2, I_3$  over all  $(\alpha, \beta)$  with  $|\alpha| + |\beta| = (p + 1)m$ , we can estimate correspondingly  $|u|_{(p+1)m}$  and  $\sigma_{p+1}(u, \lambda)$  from (4.1), (4.3); for short:

$$(4.8) \quad |u|_{(p+1)m} \leq J_1 + J_2 + J_3, \quad \sigma_{p+1}(u, \lambda) \leq Y_1 + Y_2 + Y_3.$$

Since  $J_1 \leq C|Pu|_p$  for a new constant  $C$ , then for  $\lambda$  sufficiently large,

$$(4.9) \quad Y_1 = ((p + 1)m)!^{-\mu} \lambda^{-p-1} J_1 \leq [(pm + 1) \dots (pm + m)]^{-\mu} \sigma_p(Pu, \lambda).$$

To treat  $Y_2$ , by using the expression of  $P$  in (1.3), we compute

$$[P, x^\delta D_x^\gamma] = \sum_{|\tilde{\alpha}|+|\tilde{\beta}| \leq m} c_{\alpha\beta} [x^{\tilde{\beta}} D_x^{\tilde{\alpha}}, x^\delta D_x^\gamma]$$

with

$$\begin{aligned} [x^{\tilde{\beta}} D_x^{\tilde{\alpha}}, x^\delta D_x^\gamma] &= \sum_{0 \neq \tau \leq \tilde{\alpha}, \tau \leq \delta} C_{1\tilde{\alpha}\delta\tau} x^{\delta+\tilde{\beta}-\tau} D_x^{\gamma+\tilde{\alpha}-\tau} \\ &\quad - \sum_{0 \neq \tau \leq \tilde{\beta}, \tau \leq \gamma} C_{2\tilde{\beta}\gamma\tau} x^{\delta+\tilde{\beta}-\tau} D_x^{\gamma+\tilde{\alpha}-\tau}, \end{aligned}$$

where  $|C_{1\tilde{\alpha}\delta\tau}|$  and  $|C_{2\tilde{\beta}\gamma\tau}|$  can be estimated by  $C_3(pm)^{|\tau|}$ . The constants here and in the sequel do not depend on  $p$ . Hence

$$(4.10) \quad \|[P, x^\delta D_x^\gamma] u\|_{L^2(\mathbb{R}^n)} \leq C_4 \sum_{|\tilde{\alpha}|+|\tilde{\beta}| \leq m} \sum_{\tau} (pm)^{|\tau|} \|x^{\delta+\tilde{\beta}-\tau} D_x^{\gamma+\tilde{\alpha}-\tau} u\|_{L^2(\mathbb{R}^n)},$$

with  $0 \neq \tau \leq \tilde{\alpha}, \tau \leq \delta$  or  $0 \neq \tau \leq \tilde{\beta}, \tau \leq \gamma$ . Set  $s = |\delta + \tilde{\beta} - \tau| + |\gamma + \tilde{\alpha} - \tau| = (pm + |\tilde{\alpha}| + |\tilde{\beta}| - 2|\tau|)$ . Since  $|\tilde{\alpha}| + |\tilde{\beta}| \leq m$  and  $0 < |\tau| \leq m$ , we have  $(p - 1)m \leq s < (p + 1)m$ . Note also that  $s \leq (p + 1)m - 2|\tau|$ ; hence in (4.10) we can estimate  $|\tau| \leq ((p + 1)m - s)/2$ . We may then write

$$(4.11) \quad J_2 \leq C_5(J'_2 + (pm)^{m/2}|u|_{pm} + J''_2)$$

with

$$(4.12) \quad \begin{aligned} J'_2 &= \sum_{pm < s < (p+1)m} (pm)^{((p+1)m-s)/2} |u|_s, \\ J''_2 &= \sum_{(p-1)m \leq s < pm} (pm)^{((p+1)m-s)/2} |u|_s. \end{aligned}$$

We estimate  $|u|_s$  in (4.12) by Proposition 4.1. Namely, in the expression of  $J'_2$  we apply (4.2) with

$$\epsilon = (pm)^{-((p+1)m-s)/2} (4mC_5)^{-1}$$

and we get

$$(4.13) \quad J'_2 \leq (4C_5)^{-1} |u|_{(p+1)m} + C_6(pm)^{m/2} |u|_{pm} + C_7^{p+1} ((p + 1)m)!^{1/2} |u|_0.$$

Similarly

$$(4.14) \quad J''_2 \leq C_8(pm)^{m/2} |u|_{pm} + C_9(pm)^m |u|_{(p-1)m} + C_9^{p+1} ((p + 1)m)!^{1/2} |u|_0.$$

Applying (4.13), (4.14) in (4.11) and taking  $\lambda$  sufficiently large, we conclude

$$Y_2 = ((p + 1)m)!^{-\mu} \lambda^{-p-1}, \quad J_2 \leq \frac{1}{4} (\sigma_{p+1}(u, \lambda) + \sigma_p(u, \lambda) + \sigma_{p-1}(u, \lambda) + \sigma_0(u, \lambda)).$$

The same arguments give identical estimates for  $Y_3$ , and summing up with (4.9) in (4.8) we obtain (4.4). Proposition 4.2 is therefore proved.

**End of the proof of Lemma 3.1.** Rewrite the inequality (4.4) with  $\sigma_q(u, \lambda)$  in the left-hand side,  $q = p, p - 1, \dots$ . Applying these estimates successively to the terms  $\sigma_p(u, \lambda), \sigma_{p-1}(u, \lambda), \dots$  in the right-hand side of (4.4), we deduce, for  $\lambda$  sufficiently large,

$$(4.15) \quad \sigma_{p+1}(u, \lambda) \leq \sum_{q=0}^p [(qm + 1) \dots (qm + m)]^{-\mu} \sigma_q(Pu, \lambda) + \sigma_0(u, \lambda).$$

We then obtain by induction

$$(4.16) \quad \sigma_p(u, \lambda) \leq \sum_{r=0}^p \binom{p}{r} (rm)!^{-\mu} \sigma_0(P^r u, \lambda).$$

Namely, given (4.16) as granted for  $q \leq p$ , we want to prove it for  $\sigma_{p+1}(u, \lambda)$ . As a first step we apply (4.15), and then we estimate  $\sigma_q(Pu, \lambda), q \leq p$ , in the right-hand side of (4.15) by the inductive assumptions. Easy computations (see for example [9]) give (4.16) for  $\sigma_{p+1}(u, \lambda)$ .

Finally, let us apply the inequality (3.1) with  $M = r = 0, 1, \dots, p$  to the right-hand side of (4.16). Standard factorial and binomial estimates show that  $\sigma_p(u, \lambda) \leq C^{p+1}$  for a new constant  $C > 0$ . Enlarging  $\lambda$ , we conclude that the sequence  $\sigma_p(u, \lambda)$  in (4.3) is bounded. Lemma 3.1 is proved.

*Remark 4.3.* The results in [4] lead to a natural conjecture that one can extend the result of Theorem 1.2 for  $S_{mt/(m+k)}^{kt/(k+m)}(\mathbb{R}^n)$ ,  $t \geq 1$ , by means of eigenfunction expansions for anisotropic elliptic Shubin type operators modelled by  $(-\Delta)^m + x^{2k}$ .

## REFERENCES

1. P. Boggiatto, E. Buzano and L. Rodino, *Global hypoellipticity and spectral theory*. Math. Res. **92**, Akademie Verlag, Berlin, 1996. MR1435282 (97m:35040)
2. D. Calvo and L. Rodino, *Iterates of operators and Gelfand-Shilov functions*. To appear in Int. Transf. Spec. Funct.
3. M. Capiello, T. Gramchev and L. Rodino, *Super-exponential decay and holomorphic extensions for semilinear equations with polynomial coefficients*. J. Funct. Anal. **237** (2006), 634–654. MR2230353 (2007e:35075)
4. M. Capiello, T. Gramchev and L. Rodino, *Entire extensions and exponential decay for semilinear elliptic equations*. J. Anal. Math. **111** (2010), 339–367.
5. A. Erdélyi, W. Magnus, F. Oberhettinger and F.G. Tricomi, *Higher Transcendental Functions*, Vols. 1-3, McGraw-Hill, New York, 1953, 1955. MR0058756 (15:419i), MR0066496 (16:586c)
6. I.M. Gelfand and G.E. Shilov, *Generalized functions II*. Academic Press, New York, 1968. MR0230128 (37:5693)
7. T. Gramchev, S. Pilipović and L. Rodino, *Global Regularity and Stability in S-Spaces for Classes of Degenerate Shubin Operators*. Pseudo-Differential Operators: Complex Analysis and Partial Differential Equations Operator Theory: Advances and Applications **205** (2010), 81-90. MR2664575
8. B. Helffer, *Théorie spectrale pour des opérateurs globalement elliptiques*. Astérisque **112**, Société Mathématique de France, Paris, 1984. MR743094 (86d:35151)
9. H. Komatsu, *A proof of Kotake and Narashiman's theorem*. Proc. Japan Acad. **38** (1962), 615-618. MR0152909 (27:2880)
10. T. Kotake and M.S. Narasimhan, *Regularity theorems for fractional powers of a linear elliptic operator*. Bull. Soc. Math. France **90** (1962), 449-471. MR0149329 (26:6819)
11. M. Langenbruch, *Hermite functions and weighted spaces of generalized functions*. Manuscripta Math. **119** (2006), 269–285. MR2207851 (2007c:46039)
12. F. Nicola and L. Rodino, *Global pseudo-differential calculus on Euclidean spaces*, Birkhäuser, Basel, 2010. MR2668420
13. S. Pilipovic, *Tempered ultradistributions*. Boll. Unione Mat. Ital. VII. Ser. B **2** (1988), 235–251. MR946067 (89f:46088)
14. M. Reed and B. Simon, *Methods of modern mathematical physics*, Vol. 1, Academic Press, San Diego, CA, 1972. MR0493419 (58:12429a)
15. R.T. Seeley, *Integro-differential operators on vector bundles*, Trans. Amer. Math. Soc. **117** (1965), 167-204. MR0173174 (30:3387)
16. R.T. Seeley, *Eigenfunction expansions of analytic functions*, Proc. Amer. Math. Soc. **21** (1969), 734-738. MR0240835 (39:2180)
17. M. Shubin, *Pseudodifferential operators and spectral theory*. Springer Series in Soviet Mathematics, Springer Verlag, Berlin, 1987. MR883081 (88c:47105)
18. M.W. Wong, *The heat equation for the Hermite operator on the Heisenberg group*. Hokkaido Math. J. **34** (2005), 393–404. MR2159004 (2006c:35112)

DIPARTIMENTO DI MATEMATICA E INFORMATICA, UNIVERSITÀ DI CAGLIARI, VIA OSPEDALE 72, 09124 CAGLIARI, ITALY

*E-mail address:* `todor@unica.it`

INSTITUTE OF MATHEMATICS, UNIVERSITY OF NOVI SAD, TRG. D. OBRADOVICA 4, 21000 NOVI SAD, SERBIA

*E-mail address:* `stevan.pilipovic@uns.dmi.ac.rs`

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI TORINO, VIA CARLO ALBERTO 10, 10123 TORINO, ITALY

*E-mail address:* `luigi.rodino@unito.it`