PROCEEDINGS OF THE AMERICAN MATHEMATICAL SOCIETY Volume 140, Number 9, September 2012, Pages 3289–3295 S 0002-9939(2012)11182-7 Article electronically published on January 20, 2012

VOLUME FORMULAS FOR A SPHERICAL TETRAHEDRON

JUN MURAKAMI

(Communicated by Jianguo Cao)

ABSTRACT. The present paper gives two concrete formulas for the volume of an arbitrary spherical tetrahedron that is in a 3-dimensional spherical space of constant curvature +1. One formula is given in terms of dihedral angles, and another one is given in terms of edge lengths.

INTRODUCTION

The calculation of the volume of an arbitrary tetrahedron in a 3-space of non-zero constant curvature is rather hard, and the first result was given in [1] in 1999 for hyperbolic tetrahedra. The papers [5] and [4] gave another formula for hyperbolic tetrahedra, which is implicitly based on the quantum 6j-symbol. Moreover, it was stated in [5] that an adequate analytic continuation of the obtained formula is also applicable for a spherical tetrahedron. But the formula is given by multivalued functions, and it is not described which stratum we should select for actual computation. On the other hand, volumes of spherical tetrahedra of special shapes were given by many people in the past, and the most recent work is [2], which gives a formula for a spherical tetrahedron having a small symmetry.

In the present paper, volume formulas for a spherical tetrahedron T of general shape are given in Theorems 1.1 and 1.2. The formula in Theorem 1.1 is given in terms of dihedral angles, and the formula in Theorem 1.2 is given in terms of edge lengths. These formulas are obtained by improving those in [5] and [4], and, by using the Schläfli differential equality, it is shown that the new formulas actually give the volume of T modulo $2\pi^2$. Note that $2\pi^2$ is the volume of S^3 with radius 1, which is the universal cover of any 3-dimensional spherical space of constant curvature +1. Since T can be included in a 3-dimensional hemisphere, the volume of T is less than π^2 and so we can actually compute the volume of T from the formulas in Theorems 1.1 and 1.2.

1. Volume formulas

1.1. Volume formula in terms of dihedral angles. Let T be a spherical tetrahedron and let $\theta_1, \theta_2, \dots, \theta_6$ be its dihedral angles at edges e_1, e_2, \dots, e_6 , respectively, given in Figure 1. We assume that $0 < \theta_j < \pi$ for $j = 1, 2, \dots, 6$. Let

©2012 American Mathematical Society

Received by the editors November 27, 2010 and, in revised form, March 29, 2011.

²⁰¹⁰ Mathematics Subject Classification. Primary 51M25; Secondary 52A38, 26B15.

Key words and phrases. Tetrahedron, volume, spherical space.

This research was partially supported by Grant-in-Aid for Scientific Research(C) 22540236 from JSPS.



FIGURE 1. Edges of T

$$\begin{aligned} a_1 &= e^{i\theta_1}, \ a_2 &= e^{i\theta_2}, \ \cdots, \ a_6 &= e^{i\theta_6}, \ \text{and} \\ L(a_1, a_2, \cdots, a_6, z) \\ &= \frac{1}{2} \left(\text{Li}_2(z) + \text{Li}_2(a_1^{-1} a_2^{-1} a_4^{-1} a_5^{-1} z) + \text{Li}_2(a_1^{-1} a_3^{-1} a_4^{-1} a_6^{-1} z) \right. \\ &\quad + \text{Li}_2(a_2^{-1} a_3^{-1} a_5^{-1} a_6^{-1} z) - \text{Li}_2(-a_1^{-1} a_2^{-1} a_3^{-1} z) - \text{Li}_2(-a_1^{-1} a_5^{-1} a_6^{-1} z) \\ &\quad - \text{Li}_2(-a_2^{-1} a_4^{-1} a_6^{-1} z) - \text{Li}_2(-a_3^{-1} a_4^{-1} a_5^{-1} z) + \sum_{j=1}^3 \log a_j \log a_{j+3} \Big), \end{aligned}$$

where $\text{Li}_2(z)$ is the dilogarithm function defined by analytic continuation of the following integral:

(1.1)
$$\operatorname{Li}_{2}(x) = -\int_{0}^{x} \frac{\log(1-t)}{t} dt \quad \text{for a real number } x < 1.$$

The analytic continuation of the right-hand side integral defines a multi-valued complex function $\text{li}_2(z)$, and let $\text{Li}_2(z)$ be the principal branch of $\text{li}_2(z)$ which is the analytic continuation of (1.1) on the region $\mathbb{C} \setminus \{x \in \mathbb{R} \mid x \geq 1\}$. We also fix the principal branch of the log function as usual by the branch cut along the negative real axis.

We define the auxiliary parameter z_0 as

(1.2)
$$z_0 = \frac{-q_1 + \sqrt{q_1^2 - 4q_0q_2}}{2q_2},$$

where

 $q_0 = a_1 \, a_4 + a_2 \, a_5 + a_3 \, a_6 + a_1 \, a_2 \, a_6 + a_1 \, a_3 \, a_5 + a_2 \, a_3 \, a_4$

 $+ a_4 a_5 a_6 + a_1 a_2 a_3 a_4 a_5 a_6,$

$$q_{1} = -(a_{1} - a_{1}^{-1})(a_{4} - a_{4}^{-1}) - (a_{2} - a_{2}^{-1})(a_{5} - a_{5}^{-1}) - (a_{3} - a_{3}^{-1})(a_{6} - a_{6}^{-1}),$$

$$q_{2} = a_{1}^{-1}a_{4}^{-1} + a_{2}^{-1}a_{5}^{-1} + a_{3}^{-1}a_{6}^{-1} + a_{1}^{-1}a_{2}^{-1}a_{6}^{-1} + a_{1}^{-1}a_{3}^{-1}a_{5}^{-1} + a_{2}^{-1}a_{3}^{-1}a_{4}^{-1} + a_{5}^{-1}a_{6}^{-1} + a_{1}^{-1}a_{5}^{-1}a_{6}^{-1} + a_{1}^{-1}a_{2}^{-1}a_{3}^{-1}a_{4}^{-1}a_{5}^{-1}a_{6}^{-1} + a_{1}^{-1}a_{5}^{-1}a_{6}^{-1} + a_{1}^{-1}a_{5}^{-1}a_{6}$$

Then z_0 is a solution of

(1.3)
$$\exp\left(2\,z\,\frac{\partial L}{\partial z}\right) = 1$$

where

$$\exp\left(2z\frac{\partial L}{\partial z}\right) = \frac{(a_1\,a_2\,a_3+z)\,(a_1\,a_5\,a_6+z)\,(a_2\,a_4\,a_6+z)\,(a_3\,a_4\,a_5+z)}{(1-z)\,(a_1\,a_2\,a_4\,a_5-z)\,(a_1\,a_3\,a_4\,a_6-z)\,(a_2\,a_3\,a_5\,a_6-z)}.$$

3290

Now we state the main result of this paper.

Theorem 1.1. Let T be a spherical tetrahedron with dihedral angles $\theta_1, \theta_2, \dots, \theta_6$ at the edges $e_1, e_2, \dots e_6$ given in Figure 1. Let $a_j = e^{i\theta_j}$ for $j = 1, 2, \dots, 6$ and let Vol(T) be the volume of T. Then

$$\operatorname{Vol}(T) = -\operatorname{Re}(L(a_1, a_2, \cdots, a_6, z_0)) + \pi \left(\arg(-q_2) + \frac{1}{2} \sum_{j=1}^6 \theta_j \right) - \frac{3}{2} \pi^2$$

mod $2\pi^2$,

where $\operatorname{Re}(z)$ is the real part of z and z_0 , q_2 given by (1.2).

1.2. Volume formula in terms of edge lengths. Let T be a spherical tetrahedron with edge lengths l_1, l_2, \dots, l_6 at the edges $e_1, e_2, \dots e_6$, respectively, given in Figure 1. Let $b_j = e^{i l_j}$ for $j = 1, 2, \dots, 6$ and let $\widetilde{L}(b_1, b_2, b_3, b_4, b_5, b_6, z) = L(-b_4^{-1}, -b_5^{-1}, -b_6^{-1}, -b_2^{-1}, -b_3^{-1}, z)$. Then the following formula holds.

Theorem 1.2. For a spherical tetrahedron T as above,

$$\operatorname{Vol}(T) = \operatorname{Re}\left(\widetilde{L}(b_1, b_2, \cdots, b_6, \widetilde{z}_0)\right) - \pi \operatorname{arg}(-\widetilde{q}_2) - \sum_{j=1}^6 l_j \left. \frac{\partial \operatorname{Re}\left(\widetilde{L}(b_1, b_2, \cdots, b_6, z)\right)}{\partial l_j} \right|_{z=\widetilde{z}_0} - \frac{1}{2} \pi^2 \mod 2\pi^2,$$

where \tilde{z}_0 and \tilde{q}_2 are obtained from z_0 and q_2 in (1.2) by substituting a_j with $-b_{j\pm 3}^{-1}$ for $j = 1, 2, \dots, 6$.

2. Proof of the formulas

2.1. **Gram matrices.** Let T be a spherical tetrahedron with dihedral angles θ_1 , \cdots , θ_6 as before. Let G be the Gram matrix of T defined by

$$G = \begin{pmatrix} 1 & -\cos\theta_1 & -\cos\theta_2 & -\cos\theta_6 \\ -\cos\theta_1 & 1 & -\cos\theta_3 & -\cos\theta_5 \\ -\cos\theta_2 & -\cos\theta_3 & 1 & -\cos\theta_4 \\ -\cos\theta_6 & -\cos\theta_5 & -\cos\theta_4 & 1 \end{pmatrix}.$$

An actual computation shows that the discriminant in (1.2) is given by

(2.1)
$$q_1^2 - 4 q_0 q_2 = 16 \det G,$$

which is positive since T is spherical. It is known¹ that

(2.2)
$$\cos l_j = \frac{c_{pq}}{\sqrt{c_{pp} c_{qq}}},$$

and so we have

(2.3)
$$\exp(2\,i\,l_j) = \frac{2\,c_{pq}^2 - c_{pp}\,c_{qq} + 2\,i\,c_{pq}\,\sqrt{\det G}\,\sin\theta_j}{c_{pp}\,c_{qq}}$$

¹Formula (2.2) comes from the formula on p. 7, l.4, of [2] applied to the dual tetrahedron T^* . It is a spherical version of the formula just below (5.1) in [6].

JUN MURAKAMI

by using the formula (5.1) in [6] that is $c_{pq}^2 - c_{pp} c_{qq} = -\det G \sin^2 \theta_j$. Here p and q denote the row and column of $G = (g_{ab})$ such that $g_{p'q'} = -\cos \theta_j$, $\{p,q\} = \{1,2,3,4\} \setminus \{p',q'\}$ and c_{ab} is the cofactor of G, i.e. $c_{ab} = (-1)^{a+b} \det G_{ab}$ where G_{ab} is the submatrix obtained from G by deleting its a-th row and b-th column.

2.2. Some functions and their properties. Before proving the formulas, we introduce some functions and investigate their properties. Let T be an abstract tetrahedron, let $\theta_1, \theta_2, \dots, \theta_6$ be its dihedral angles at the edges e_1, e_2, \dots, e_6 as before, and let

$$D_s = \{(\theta_1, \theta_2, \cdots, \theta_6) \in (0, \pi)^6 \subset \mathbb{R}^6 \mid \theta_1, \theta_2, \cdots, \theta_6\}$$

correspond to the dihedral angles of a spherical tetrahedron}.

Let $a_j = e^{i\theta_j}$ for $j = 1, 2, \cdots, 6$,

$$\begin{split} \Delta_0(x,y,z) &= \\ &-\frac{1}{4} \Big(\operatorname{Li}_2(-xy^{-1}z^{-1}) + \operatorname{Li}_2(-x^{-1}yz^{-1}) + \operatorname{Li}_2(-x^{-1}y^{-1}z) + \operatorname{Li}_2(-xyz) \Big), \\ &\Delta(a_1,a_2,\cdots,a_6) = \Delta_0(a_1,a_2,a_3) + \Delta_0(a_1,a_5,a_6) + \Delta_0(a_2,a_4,a_6) \\ &+ \Delta_0(a_3,a_4,a_5) - \frac{1}{2} \sum_{j=1}^6 \big(\log a_j \big)^2, \\ &U(a_1,a_2,\cdots,a_6,z) = L(a_1,a_2,\cdots,a_6,z) + \Delta(a_1,a_2,\cdots,a_6) \end{split}$$

and

 $V(a_1, a_2, a_3, a_4, a_5, a_6)$

$$= -U(a_1, a_2, a_3, a_4, a_5, a_6, z_0) + \pi i \left(\log z_0 - \sum_{j=1}^6 \log a_j \right) - \frac{13}{6} \pi^2.$$

Lemma 2.1. The function $\Delta(a_1, a_2, \dots, a_6)$ is analytic on D_s and the imaginary part of $4a_j \frac{\partial \Delta}{\partial a_i}$ is given by

$$\operatorname{Im}\left(4\,a_j\,\frac{\partial\Delta}{\partial a_j}\right) = -2\,\pi.$$

Proof. We prove this for the case j = 1. For the function Δ ,

$$a_1 \frac{\partial \Delta}{\partial a_1} = a_1 \frac{\partial \Delta_0(a_1, a_2, a_3)}{\partial a_1} + a_1 \frac{\partial \Delta_0(a_1, a_5, a_6)}{\partial a_1} - \log a_1$$

and

$$a_1 \frac{\partial \Delta_0(a_1, a_p, a_q)}{\partial a_1} = \frac{1}{4} \left(\log(1 + \frac{a_1}{a_p a_q}) - \log(1 + \frac{a_p}{a_1 a_q}) - \log(1 + \frac{a_q}{a_1 a_p}) + \log(1 + a_1 a_p a_q) \right)$$

3292

for $\{p,q\} = \{2,3\}, \{5,6\}$. The imaginary part $\operatorname{Im} \log(1 + e^{i\theta})$ is given by

$$\operatorname{Im}\log(1+e^{i\theta}) = \begin{cases} \frac{\theta}{2} & \text{if } -\pi < \theta < \pi, \\ \frac{\theta}{2} - \pi & \text{if } \pi < \theta < 3\pi. \end{cases}$$

Let θ_u , θ_v , θ_w be three dihedral angles at three edges having a vertex in common. Then they satisfy

(2.4) $0 < \theta_u + \theta_v - \theta_w$, $\theta_u - \theta_v + \theta_w$, $-\theta_u + \theta_v + \theta_w < \pi$, $\pi < \theta_u + \theta_v + \theta_w < 3\pi$. Hence $\Delta_0(a_1, a_p, a_q)$ is analytic on D_s and we have

$$\operatorname{Im}\left(a_1 \frac{\partial \Delta_0(a_1, a_p, a_q)}{\partial a_1}\right) = \frac{\theta_1}{2} - \frac{\pi}{4}, \qquad \operatorname{Im}\left(4 a_1 \frac{\partial \Delta}{\partial a_1}\right) = -2 \pi.$$

Moreover, Δ is analytic on D_s because none of the imaginary parts of the log terms of Δ attain either π or $-\pi$ on D_s .

Lemma 2.2. The function $L(a_1, a_2, \dots, a_6, z_0(a_1, a_2, \dots, a_6))$ is analytic on D_s , and so $U(a_1, a_2, \dots, a_6, z_0(a_1, a_2, \dots, a_6))$ is analytic on D_s .

Proof. We know that $|z_0| < 1$ because, for q_0 , q_1 , q_2 in (1.2), q_1 is a real number and $q_0q_2 = q_0\overline{q_0} = |q_0|^2$ is a positive real number, and $q_1^2 - 4q_0q_2$ is also a positive real number by (2.1). This implies that, for $w \in \mathbb{C}$ with |w| = 1, $|w z_0| < 1$. Noting that $\text{Li}_2(z)$ is analytic on the unit open disk $\{z \in \mathbb{C} \mid |z| < 1\}$, all the dilog terms of L are analytic on D_s since $|a_1| = \cdots = |a_6| = 1$.

Lemma 2.3. The differential $\frac{\partial U}{\partial z}$ satisfies $z_0 \frac{\partial U}{\partial z}\Big|_{z=z_0} = \pi i$.

Proof. Since $\frac{\partial U}{\partial z} = \frac{\partial L}{\partial z}$ and z_0 is a solution of equation (1.3), $z_0 \frac{\partial U}{\partial z}\Big|_{z=z_0} = k \pi i$ for some integer constant k because U is analytic on D_s by the above lemma. Let $T_{\frac{\pi}{2}}$ be the regular spherical tetrahedron with edge lengths $\pi/2$. Then $\theta_j = \pi/2$, $a_j = i$ for $j = 1, \dots, 6$, $z_0 = (i+1)/2$ and

$$z_0 \left. \frac{\partial U}{\partial z} \right|_{z=z_0} = \frac{1}{2} \left(-4 \log \frac{1-i}{2} + 4 \log \frac{1+i}{2} \right) = \pi i.$$

Hence $z_0 \frac{\partial U}{\partial z}\Big|_{z=z_0} = \pi i$ for all the spherical tetrahedra.

Now we show the following proposition for V corresponding to the Schläfli differential equality

(2.5)
$$d\operatorname{Vol}(T) = \sum_{j=1}^{6} \frac{l_j}{2} d\theta_j,$$

which is a fundamental tool for analyzing the volume. For example, see [3]. **Proposition 2.4.** The function V satisfies $\frac{\partial V}{\partial \theta_j} = l_j/2$ for $j = 1, 2, \cdots, 6$. Proof. Let $\varphi = \exp\left(4a_1 \frac{\partial \Delta}{\partial a_1}\right)$ and $\psi = \exp\left(2a_1 \frac{\partial L}{\partial a_1}\Big|_{z=z_0}\right)$. Then $\varphi = \frac{(a_1 + a_2 a_3)(a_1 a_2 a_3 + 1)(a_1 + a_5 a_6)(a_1 a_5 a_6 + 1)}{(a_1 a_2 + a_3)(a_1 a_3 + a_2)(a_1 a_5 + a_6)(a_1 a_6 + a_5)},$ $\psi = \frac{(a_1 a_2 a_4 a_5 - z_0)(a_1 a_3 a_4 a_6 - z_0)}{a_4 (a_1 a_2 a_3 + z_0)(a_1 a_5 a_6 + z_0)}.$ An actual computation and (2.3) show that

$$\exp\left(4a_1\frac{\partial U}{\partial a_1}\right)\Big|_{z=z_0} = \varphi\psi^2 = \frac{2c_{34}^2 - c_{33}c_{44} + 2ic_{34}\sqrt{\det G}\sin\theta_1}{c_{33}c_{44}} = \exp(2l_1i).$$

Hence we get $a_1 \frac{\partial U}{\partial a_1}\Big|_{z=z_0} = i (l_1 + k \pi)/2$ for some integer constant k because U is analytic on D_s by Lemma 2.2. For the tetrahedron $T_{\frac{\pi}{2}}$ given in the proof of Lemma 2.3, $l_1 = \pi/2$ and $a_1 \frac{\partial U}{\partial a_1}\Big|_{z=z_0} = -3 \pi i/4$, which means that k = -2 and $a_1 \frac{\partial U}{\partial a_1}\Big|_{z=z_0} = i (l_1 - 2\pi)/2$. According to $\frac{\partial U}{\partial \theta_1} = i a_1 \frac{\partial U}{\partial a_1}$, we have

(2.6)
$$\frac{\partial U}{\partial \theta_1}\Big|_{z=z_0} = \frac{1}{2} (2\pi - l_1).$$

Therefore

$$\frac{\partial}{\partial \theta_1} \left(-U(a_1, \cdots, a_6, z_0(a_1, \cdots, a_6)) + \pi i \left(\log z_0 - \sum_{j=1}^6 \log a_j \right) \right)$$
$$= \frac{l_1}{2} - \frac{\partial z_0}{\partial \theta_1} \frac{\partial U}{\partial z} \Big|_{z=z_0} + \pi i \frac{\partial z_0}{\partial \theta_1} \frac{1}{z_0}.$$

Since $\frac{\partial U}{\partial z}\Big|_{z=z_0} = i \pi/z_0$ by Lemma 2.3, we get $\frac{\partial V}{\partial \theta_1} = l_1/2$.

2.3. **Proof of the formula in terms of dihedral angles.** We first give a formula using complex analytic functions.

Proposition 2.5. Let T be a spherical tetrahedron with dihedral angles $\theta_1, \theta_2, \dots, \theta_6$ at the edges $e_1, e_2, \dots e_6$ as in Figure 1. Let $a_j = e^{i\theta_j}$ for $j = 1, 2, \dots, 6$ as before and let Vol(T) be the volume of T. Then

 $Vol(T) = V(a_1, a_2, a_3, a_4, a_5, a_6) \mod 2\pi^2.$

Proof. For the tetrahedron $T_{\frac{\pi}{2}}$ in the proof of Lemma 2.3, we have $a_j = i, z_0 = \frac{1+i}{2}$ and $V(i, i, i, i, i, i, i) = \frac{\pi^2}{8} = \operatorname{Vol}(T_{\frac{\pi}{2}})$ since $T_{\frac{\pi}{2}}$ is one-sixteenth of S^3 and the volume of S^3 with radius 1 is $2\pi^2$. Because V is analytic on some neighborhood N of $T_{\frac{\pi}{2}}$ in D_s , two functions V and Vol are identical on N by Proposition 2.4 and Schäfli differential equality. Moreover, Vol is analytic on D_s and so it is given by an adequate analytic continuation of V. We already showed in previous lemmas that all the terms in V except $\pi i \log z_0$ are analytic on D_s , and the analytic continuation of $\pi i \log z_0$ is $\pi i \log z_0 + 2k\pi^2$ for some integer k. Hence we get the proposition. \Box

Proof of Theorem 1.1. We prove Theorem 1.1 by investigating the real part of V. For $\theta \in [0, 2\pi] \subset \mathbb{R}$, the real part of $\text{Li}_2(e^{i\theta})$ is given by $\text{Re}(\text{Li}_2(e^{i\theta})) = \text{Re}(\text{Li}_2(e^{-i\theta})) = \theta^2/4 - \pi \theta/2 + \pi^2/6$. Substituting this in each dilog function of $\text{Re}(\Delta(a_1, a_2, \dots, a_6)), \text{ we get } \text{Re}(\Delta(a_1, a_2, \dots, a_6)) = -2\pi^2/3 + \sum_{j=1}^6 \pi \theta_j/2$ by using (2.4). We also know that $\text{Im} \log z_0 = -\arg(-q_2)$ since the numerator of z_0 in (1.2) is a negative real number. Hence we get Theorem 1.1 from Proposition 2.5. \Box

Remark 2.6. The function V is non-continuous at the points where the values of q_2 are positive real numbers.

3294

2.4. Proof of the formula in terms of edge lengths. We use the notation in Subsection 2.2.

Proof of Theorem 1.2. Let $\theta_1, \theta_2, \dots, \theta_6$ be the dihedral angles at the edges e_1, e_2 , \cdots , e_6 of T and let T^* be the dual tetrahedron of T given by [3, p. 294]. Then the dihedral angles of T^* are $\pi - l_4$, $\pi - l_5$, $\pi - l_6$, $\pi - l_1$, $\pi - l_2$, $\pi - l_3$ and the edge lengths of T^* are $\pi - \theta_4$, $\pi - \theta_5$, $\pi - \theta_6$, $\pi - \theta_1$, $\pi - \theta_2$, $\pi - \theta_3$. The relation of volumes of T and T^* is given by [3, p. 294] as follows:

$$\operatorname{Vol}(T) + \operatorname{Vol}(T^*) + \frac{1}{2} \sum_{j=1}^{6} l_j (\pi - \theta_j) = \pi^2$$

By Theorem 1.1, we have

$$\operatorname{Vol}(T^*) = -\operatorname{Re}(\widetilde{L}(b_1, b_2, \cdots, b_6, \widetilde{z}_0)) + \pi \left(\arg(-\widetilde{q}_2) + \frac{1}{2} \sum_{j=1}^6 (\pi - l_j) \right) - \frac{3}{2} \pi^2 \mod 2\pi^2.$$

Because $\frac{\partial}{\partial(\pi-l_j)}U(-b_4^{-1}, -b_5^{-1}, -b_6^{-1}, -b_1^{-1}, -b_2^{-1}, -b_3^{-1}, z)\Big|_{z=\tilde{z}_0} = (2\pi - (\pi - \theta_j))/2$ by (2.6) and $\frac{\partial}{\partial(\pi-l_j)} \operatorname{Re}\left(\Delta(-b_4^{-1}, -b_5^{-1}, -b_6^{-1}, -b_1^{-1}, -b_2^{-1}, -b_3^{-1})\right) = \pi/2$ by Lemma 2.1, we know that $\frac{\partial \operatorname{Re}\tilde{L}}{\partial l_j}\Big|_{z=\tilde{z}_0} = -\theta_j/2$. Hence

$$\operatorname{Vol}(T) = \operatorname{Re}\left(\widetilde{L}(b_1, b_2, \cdots, b_6, \widetilde{z}_0)\right) - \pi \operatorname{arg}(-\widetilde{q}_2) \\ - \sum_{j=1}^6 l_j \left. \frac{\partial \operatorname{Re}\left(\widetilde{L}(b_1, b_2, \cdots, b_6, z)\right)}{\partial l_j} \right|_{z=\widetilde{z}_0} - \frac{1}{2} \pi^2 \mod 2\pi^2,$$

and we get Theorem 1.2.

and we get Theorem 1.2.

References

- [1] Y. Cho and H. Kim, On the volume formula for hyperbolic tetrahedra, Discrete Comput. Geom. 22 (1999), no. 3, 347-366. MR1706606 (2000k:52008)
- [2] A. Kolpakov, A. Mednykh and M. Pashkevich, Volume formula for a \mathbb{Z}_2 -symmetric spherical tetrahedron through its edge lengths, to appear in Ark. Mat.
- [3] J. Milnor, The Schläfli differential equality, John Milnor Collected Papers, Volume 1, Geometry, Publish or Perish Inc., Houston, Texas, 1994, pp. 281–295. MR1277810 (95c:01043)
- [4] J. Murakami and A. Ushijima, A volume formula for hyperbolic tetrahedra in terms of edge lengths. J. Geom. 83 (2005), 153-163. MR2193233 (2006k:52012)
- [5]J. Murakami and M. Yano, On the volume of a hyperbolic and spherical tetrahedron. Comm. Anal. Geom. 13 (2005), 379-400. MR2154824 (2006d:52007)
- [6] A. Ushijima, A volume formula for generalized hyperbolic tetrahedra. In Non-Euclidean geometries, Math. Appl. (N.Y.) 581, Springer, New York, 2006, pp. 249–265. MR2191251 (2007h:52008)

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ENGINEERING, WASEDA UNIVER-SITY, 3-4-1 Ohkubo, Shinjuku-ku, Tokyo 169-8555, Japan

E-mail address: murakami@waseda.jp