

SIGNATURES, HEEGAARD FLOER CORRECTION TERMS AND QUASI-ALTERNATING LINKS

PAOLO LISCA AND BRENDAN OWENS

(Communicated by Daniel Ruberman)

ABSTRACT. Turaev showed that there is a well-defined map assigning to an oriented link L in the three-sphere a Spin structure \mathbf{t}_0 on $\Sigma(L)$, the two-fold cover of S^3 branched along L . We prove, generalizing results of Manolescu–Owens and Donald–Owens, that for an oriented quasi-alternating link L the signature of L equals minus four times the Heegaard Floer correction term of $(\Sigma(L), \mathbf{t}_0)$.

1. INTRODUCTION

Vladimir Turaev [21, §2.2] proved that there is a surjective map which associates to a link $L \subset S^3$ decorated with an orientation o a Spin structure $\mathbf{t}_{(L,o)}$ on $\Sigma(L)$, the double cover of S^3 branched along L . Moreover, he showed that the only other orientation on L which maps to $\mathbf{t}_{(L,o)}$ is $-o$, the overall reversed orientation. In other words, Turaev described a bijection between the set of quasi-orientations on L (i.e. orientations up to overall reversal) and the set $\text{Spin}(\Sigma(L))$ of Spin structures on $\Sigma(L)$. Each element $\mathbf{t} \in \text{Spin}(\Sigma(L))$ can be viewed as a Spin^c structure on $\Sigma(L)$, so if $\Sigma(L)$ is a rational homology sphere, then it makes sense to consider the rational number $d(\Sigma(L), \mathbf{t})$, where d is the correction term invariant defined by Ozsváth and Szabó [13]. Under the assumption that L is nonsplit alternating it was proved — in [10] when L is a knot and in [3] for any number of components of L — that

$$(*) \quad \sigma(L, o) = -4d(\Sigma(L), \mathbf{t}_{(L,o)}) \quad \text{for every orientation } o \text{ on } L,$$

where $\sigma(L, o)$ is the link signature. For an alternating link associated to a plumbing graph with no bad vertices, this follows from a combination of earlier results of Saveliev [19] and Stipsicz [20], each of whom showed that one of the quantities in $(*)$ is equal to the Neumann-Siebenmann $\bar{\mu}$ -invariant of the plumbing tree. The main purpose of this paper is to prove property $(*)$ for the family of *quasi-alternating links* introduced in [14]:

Definition 1. The *quasi-alternating* links are the links in S^3 with nonzero determinant defined recursively as follows:

- (1) the unknot is quasi-alternating;

Received by the editors March 7, 2013 and, in revised form, May 20, 2013.

2010 *Mathematics Subject Classification.* Primary 57M25, 57M27; Secondary 57Q60.

The present work is part of the first author's activities within CAST, a Research Network Program of the European Science Foundation, and the PRIN–MIUR research project 2010–2011 “Varietà reali e complesse: geometria, topologia e analisi armonica”.

The second author was supported in part by EPSRC grant EP/I033754/1.

- (2) if L_0, L_1 are quasi-alternating, $L \subset S^3$ is a link such that $\det L = \det L_0 + \det L_1$ and L, L_0, L_1 differ only inside a 3-ball as illustrated in Figure 1, then L is quasi-alternating.




FIGURE 1. L and its resolutions L_0 and L_1 .

Quasi-alternating links have recently been the object of considerable attention [1, 2, 4–6, 11, 16, 17, 22, 23]. Alternating links are quasi-alternating [14, Lemma 3.2], but (as shown in e.g. [1]) there exist infinitely many quasi-alternating, nonalternating links. Our main result is the following:

Theorem 1. *Let (L, o) be an oriented link. If L is quasi-alternating, then*

$$(1) \quad \sigma(L, o) = -4d(\Sigma(L), \mathbf{t}_{(L,o)}).$$

The contents of the paper are as follows. In Section 2 we first recall some basic facts on Spin structures and the existence of two natural 4-dimensional cobordisms, one from $\Sigma(L_1)$ to $\Sigma(L)$, the other from $\Sigma(L)$ to $\Sigma(L_0)$. Then, in Proposition 1 we show that for an orientation o on L for which the crossing in Figure 1 is positive, the Spin structure $\mathbf{t}_{(L,o)}$ extends to the first cobordism but not to the second one. In Section 3 we use this information together with the Heegaard Floer surgery exact triangle to prove Proposition 2, which relates the value of the correction term $d(\Sigma(L), \mathbf{t}_{(L,o)})$ with the value of an analogous correction term for $\Sigma(L_1)$. In Section 4 we restate and prove our main result, Theorem 1. The proof consists of an inductive argument based on Proposition 2 and the known relationship between the signatures of L and L_1 . The use of Proposition 2 is made possible by the fact that up to mirroring L one may always assume the crossing of Figure 1 to be positive. We close Section 4 with Corollary 3, which uses results of Rustamov and Mullins to relate Turaev’s torsion function for the two-fold branched cover of a quasi-alternating link L with the Jones polynomial of L .

2. TRIADS AND SPIN STRUCTURES

A Spin structure on an n -manifold M^n is a double cover of the oriented frame bundle of M with the added condition that if $n > 1$, then it restricts to the nontrivial double cover on fibres. A Spin structure on a manifold restricts to give a Spin structure on a codimension-one submanifold, or on a framed submanifold of codimension higher than one. As mentioned in the introduction, an orientation o on a link L in S^3 induces a Spin structure $\mathbf{t}_{(L,o)}$ on the double-branched cover $\Sigma(L)$, as in [21]. Recall also that there are two Spin structures on $S^1 = \partial D^2$: the nontrivial or *bounding* Spin structure, which is the restriction of the unique Spin structure on D^2 , and the trivial or *Lie* Spin structure, which does not extend over the disk. The restriction map from Spin structures on a solid torus to Spin structures on its boundary is injective; thus if two Spin structures on a closed 3-manifold agree

outside a solid torus, then they are the same. For more details on Spin structures see for example [7].

If Y is a 3-manifold with a Spin structure \mathfrak{t} and K is a knot in Y with framing λ , we may attach a 2-handle to K giving a surgery cobordism W from Y to $Y_\lambda(K)$. There is a unique Spin structure on $D^2 \times D^2$, which restricts to the bounding Spin structure on each framed circle $\partial D^2 \times \{\text{point}\}$ in $\partial D^2 \times D^2$. Thus the Spin structure on Y extends over W if and only if its restriction to K , viewed as a framed submanifold via the framing λ , is the bounding Spin structure. Note that this is equivalent, symmetrically, to the restriction of \mathfrak{t} to the submanifold λ framed by K being the bounding Spin structure. Moreover, the extension over W is unique if it exists.

Let L, L_0, L_1 be three links in S^3 differing only in a 3-ball B as in Figure 1. The double cover of B branched along the pair of arcs $B \cap L$ is a solid torus \tilde{B} with core C . The boundary of a properly embedded disk in B which separates the two branching arcs lifts to a disjoint pair of meridians of \tilde{B} . The preimage in $\Sigma(L)$ of the curve λ_0 shown in Figure 2 is a pair of parallel framings for C ; denote one of these by $\tilde{\lambda}_0$. Similarly, let $\tilde{\lambda}_1$ denote one of the components of the preimage in $\Sigma(L)$ of λ_1 . Since λ_0 is homotopic in $B - L$ to the boundary of a disk separating the two components of $L_0 \cap B$, we see that $\Sigma(L_0)$ is obtained from $\Sigma(L)$ by $\tilde{\lambda}_0$ -framed surgery on C . Similarly, λ_1 is homotopic in $B - L$ to the boundary of a disk separating the two components of $L_1 \cap B$, and $\Sigma(L_1)$ is obtained from $\Sigma(L)$ by $\tilde{\lambda}_1$ -framed surgery on C .

The two framings $\tilde{\lambda}_0$ and $\tilde{\lambda}_1$ differ by a meridian of C . In the terminology from [14], the manifolds $\Sigma(L), \Sigma(L_0)$ and $\Sigma(L_1)$ form a *triad* and there are surgery cobordisms

$$(2) \quad V : \Sigma(L_1) \rightarrow \Sigma(L) \quad \text{and} \quad W : \Sigma(L) \rightarrow \Sigma(L_0).$$

The surgery cobordism W is built by attaching a 2-handle to $\Sigma(L)$ along the knot C with framing $\tilde{\lambda}_0$. The cobordism V is built by attaching a 2-handle to $\Sigma(L_1)$. Dualising this handle structure, V is obtained by attaching a 2-handle to $\Sigma(L)$ along the knot C with framing $\tilde{\lambda}_1$ (and reversing orientation).




FIGURE 2. The loops λ_0 and λ_1 .

Proposition 1. *For any orientation o on L such that the crossing shown in Figure 1 is positive, the Spin structure $\mathfrak{t}_{(L,o)}$ extends to a unique Spin structure \mathfrak{s}_o on the cobordism V and does not admit an extension over W . The restriction of \mathfrak{s}_o to $\Sigma(L_1)$ is the Spin structure $\mathfrak{t}_{(L_1,o_1)}$, where o_1 is the orientation on L_1 induced by o .*

Proof. Let $\pi : \Sigma(L) \rightarrow S^3$ be the branched covering map. The Spin structure $\mathbf{t}_{(L,o)}$ is the lift $\tilde{\mathbf{s}}$ of the Spin structure restricted from S^3 to $S^3 - L$, twisted by $h \in H^1(\Sigma(L) - \pi^{-1}(L); \mathbb{Z}/2\mathbb{Z})$, where the value of h on a curve γ is the parity of half the sum of the linking numbers of $\pi \circ \gamma$ about the components of L (following Turaev [21, §2.2]). Suppose that the crossing in Figure 1 is positive as, for example, illustrated in Figure 3, so that the orientation o induces an orientation o_1 on L_1 .




FIGURE 3. The oriented link (L, o) together with the oriented resolution (L_1, o_1) and the unoriented resolution L_0 .

Then, we can compute from Figure 2 that $h(\tilde{\lambda}_1) = 0$ and $h(\tilde{\lambda}_0) = 1$. The Spin structure on S^3 restricts to the bounding structure on each of $\tilde{\lambda}_0$ and $\tilde{\lambda}_1$ using the 0–framing. The map π restricts to a diffeomorphism on neighbourhoods of $\tilde{\lambda}_0$ and $\tilde{\lambda}_1$. Therefore, the restriction of $\tilde{\mathbf{s}}$ to each of $\tilde{\lambda}_0$ and $\tilde{\lambda}_1$ using the pullback of the 0–framing is also the bounding structure. Also note that the preimage under π of a disk bounded by λ_i is an annulus with core C , so the framing of $\tilde{\lambda}_i$ given by C is the same as the pullback of the 0–framing.

The spin structure $\mathbf{t}_{(L,o)}$ is equal to $\tilde{\mathbf{s}}$ twisted by h . Since $\tilde{\mathbf{s}}$ restricts to the bounding spin structure on $\tilde{\lambda}_1$, and $h(\tilde{\lambda}_1) = 0$, we see that $\mathbf{t}_{(L,o)}$ restricts to the bounding Spin structure on $\tilde{\lambda}_1$ using the framing given by C . On the other hand since $h(\tilde{\lambda}_0) = 1$, $\mathbf{t}_{(L,o)}$ restricts to the Lie Spin structure on $\tilde{\lambda}_0$, again using the framing given by C . It follows that $\mathbf{t}_{(L,o)}$ admits a unique extension \mathbf{s}_o over the 2–handle giving the cobordism V , and does not extend over the cobordism W .

The restriction of \mathbf{s}_o to $\Sigma(L_1)$ coincides with $\mathbf{t}_{(L_1,o_1)}$ outside of the solid torus \tilde{B} , and therefore also on the closed manifold $\Sigma(L_1)$. □

3. RELATIONS BETWEEN CORRECTION TERMS

By [14, Proposition 2.1] we have the following exact triangle:

$$\begin{array}{ccc}
 \widehat{HF}(\Sigma(L_1)) & \xrightarrow{F_V} & \widehat{HF}(\Sigma(L)) \\
 & \searrow & \swarrow F_W \\
 & \widehat{HF}(\Sigma(L_0)) &
 \end{array}$$

where the maps F_V and F_W are induced by the surgery cobordisms of (2). (All the Heegaard Floer groups are taken with $\mathbb{Z}/2\mathbb{Z}$ coefficients.)

By [14, Proposition 3.3] (and notation as in that paper), if $L \subset S^3$ is a quasi–alternating link and L_0 and L_1 are resolutions of L as in Definition 1, then $\Sigma(L)$, $\Sigma(L_0)$ and $\Sigma(L_1)$ are L –spaces. Moreover, by assumption we have

$$(3) \quad |H^2(\Sigma(L); \mathbb{Z})| = |H^2(\Sigma(L_0); \mathbb{Z})| + |H^2(\Sigma(L_1); \mathbb{Z})|.$$

Since for every L -space Y we have $|H^2(Y; \mathbb{Z})| = \dim \widehat{HF}(Y)$, the Heegaard Floer surgery exact triangle reduces to a short exact sequence:

$$(4) \quad 0 \rightarrow \widehat{HF}(\Sigma(L_1)) \xrightarrow{F_V} \widehat{HF}(\Sigma(L)) \xrightarrow{F_W} \widehat{HF}(\Sigma(L_0)) \rightarrow 0.$$

The type of argument employed in the proof of the following proposition goes back to [9] and was also used in [20].

Proposition 2. *Let L be a quasi-alternating link and let L_0, L_1 be resolutions of L as in Definition 1. Let o be an orientation on L for which the crossing of Figure 1 is positive, and let o_1 be the induced orientation on L_1 . Then, the following holds:*

$$-4d(\Sigma(L), \mathbf{t}_{(L,o)}) = -4d(\Sigma(L_1), \mathbf{t}_{(L_1,o_1)}) - 1.$$

Proof. Since $\Sigma(L), \Sigma(L_1)$ and $\Sigma(L_0)$ are L -spaces, we may think of the Spin^c structures on these spaces as generators of their \widehat{HF} -groups, and we shall abuse our notation accordingly. Let $V : \Sigma(L_1) \rightarrow \Sigma(L)$ be the surgery cobordism of (2), and let \mathbf{s}_o be the unique Spin structure on V which extends $\mathbf{t}_{(L,o)}$ as in Proposition 1. Recall that, by definition, the map F_U associated to a cobordism $U : Y_1 \rightarrow Y_2$ is given by

$$F_U = \sum_{\mathbf{s} \in \text{Spin}^c(U)} F_{U,\mathbf{s}},$$

where $F_{U,\mathbf{s}} : \widehat{HF}(Y_1, \mathbf{t}_1) \rightarrow \widehat{HF}(Y_2, \mathbf{t}_2)$ and $\mathbf{t}_i = \mathbf{s}|_{Y_i}$ for $i = 1, 2$. We claim that

$$(5) \quad F_{V,\mathbf{s}_o}(\mathbf{t}_{(L_1,o_1)}) = \mathbf{t}_{(L,o)}.$$

The Heegaard Floer \widehat{HF} -groups admit a natural involution, usually denoted by \mathcal{J} . The maps induced by cobordisms are equivariant with respect to the $\mathbb{Z}/2\mathbb{Z}$ -actions associated to conjugation on Spin^c structures and the \mathcal{J} -map on the Heegaard Floer groups, in the sense that, if $\bar{x} := \mathcal{J}(x)$ for an element x , we have

$$(6) \quad F_{W,\bar{\mathbf{s}}}(\bar{x}) = \overline{F_{W,\mathbf{s}}(x)}$$

for each $\mathbf{s} \in \text{Spin}^c(W)$. Since by Proposition 1 there are no Spin structures on the surgery cobordism $W : \Sigma(L) \rightarrow \Sigma(L_0)$ of (2) which restrict to $\mathbf{t}_{(L,o)}$, the element $F_W(\mathbf{t}_{(L,o)}) \in \widehat{HF}(\Sigma(L_0))$ has no Spin component. In fact, since $\mathbf{t}_{(L,o)}$ is fixed under conjugation and we are working over $\mathbb{Z}/2\mathbb{Z}$, (6) implies that the contribution of each non- Spin $\mathbf{s} \in \text{Spin}^c(W)$ to a Spin component of $F_W(\mathbf{t}_{(L,o)})$ is cancelled by the contribution of $\bar{\mathbf{s}}$ to the same component. Therefore we may write

$$F_W(\mathbf{t}_{(L,o)}) = x + \bar{x}$$

for some $x \in \widehat{HF}(\Sigma(L_0))$. By the surjectivity of F_W there is some $y \in \widehat{HF}(\Sigma(L))$ with $F_W(y) = x$, therefore $F_W(\mathbf{t}_{(L,o)} + y + \bar{y}) = 0$, and by the exactness of (4) we have $\mathbf{t}_{(L,o)} + y + \bar{y} = F_V(z)$ for some $z \in \widehat{HF}(\Sigma(L_0))$. Since $F_V(\bar{z}) = \overline{F_V(z)} = F_V(z)$, the injectivity of F_V implies $z = \bar{z}$. Moreover, z must have some nonzero Spin component, otherwise we could write $z = u + \bar{u}$ and

$$F_V(u + \bar{u}) = \overline{F_V(u)} + \overline{F_V(\bar{u})} = \overline{F_V(u)} + F_V(u)$$

could not have the Spin component $\mathbf{t}_{(L,o)}$. This shows that there is a Spin structure $\mathbf{t} \in \widehat{HF}(\Sigma(L_1))$ such that $F_V(\mathbf{t}) = \mathbf{t}_{(L,o)}$. But, as we argued before for $F_W(\mathbf{t}_{(L,o)})$, in order for $F_V(\mathbf{t})$ to have a Spin component it must be the case that there is some Spin structure \mathbf{s} on V such that $F_{V,\mathbf{s}}(\mathbf{t}) = \mathbf{t}_{(L,o)}$. Applying Proposition 1 we conclude $\mathbf{s} = \mathbf{s}_o$ and therefore $\mathbf{t} = \mathbf{t}_{(L_1,o_1)}$. This establishes claim (5).

Using equation (3) and the fact that $\det(L_1) > 0$ it is easy to check that V is negative definite. The statement follows immediately from equation (5) and the degree–shift formula in Heegaard Floer theory [15, Theorem 7.1] using the fact that $c_1(\mathfrak{s}_o) = 0$, $\sigma(V) = -1$ and $\chi(V) = 1$. \square

4. THE MAIN RESULT AND A COROLLARY

Theorem 1. *Let (L, o) be an oriented link. If L is quasi–alternating, then*

$$(1) \quad \sigma(L, o) = -4d(\Sigma(L), \mathfrak{t}_{(L,o)}).$$

Proof. The statement trivially holds for the unknot, because the unknot has zero signature and the two–fold cover of S^3 branched along the unknot is S^3 , whose only correction term vanishes. If L is not the unknot and L is quasi–alternating, there are quasi–alternating links L_0 and L_1 such that $\det(L) = \det(L_0) + \det(L_1)$ and L , L_0 and L_1 are related as in Figure 1. To prove the theorem it suffices to show that if the statement holds for L_0 and L_1 , then it holds for L as well.

Denote by L^m the mirror image of L , and by o^m the orientation on L^m naturally induced by an orientation o on L . The orientation–reversing diffeomorphism from S^3 to itself taking L to L^m lifts to one from $\Sigma(L)$ to $\Sigma(L^m)$ sending $\mathfrak{t}_{(L,o)}$ to $\mathfrak{t}_{(L^m,o^m)}$. Thus by [8, Theorem 8.10] and [13, Proposition 4.2] we have

$$\sigma(L^m, o^m) = -\sigma(L, o)$$

and

$$4d(\Sigma(L^m), \mathfrak{t}_{(L^m,o^m)}) = 4d(-\Sigma(L), \mathfrak{t}_{(L,o)}) = -4d(\Sigma(L), \mathfrak{t}_{(L,o)}),$$

therefore equation (1) holds for (L, o) if and only if it holds for (L^m, o^m) . Hence, without loss of generality we may now fix an orientation o on L so that the crossing appearing in Figure 1 is positive.

Denote by o_1 the orientation on L_1 naturally induced by o . By [11, Lemma 2.1]

$$(7) \quad \sigma(L, o) = \sigma(L_1, o_1) - 1.$$

Since we are assuming that the statement holds for L_1 , we have

$$(8) \quad \sigma(L_1, o_1) = -4d(\Sigma(L_1), \mathfrak{t}_{(L_1,o_1)}).$$

Equations (7) and (8) together with Proposition 2 immediately imply equation (1). \square

Corollary 3. *Let (L, o) be an oriented, quasi–alternating link. Then,*

$$\tau(\Sigma(L), \mathfrak{t}_{(L,o)}) = -\frac{1}{12} \frac{V'_{(L,o)}(-1)}{V_{(L,o)}(-1)},$$

where τ is Turaev’s torsion function and $V_{(L,o)}(t)$ is the Jones polynomial of (L, o) .

Proof. By [18, Theorem 3.4] we have

$$(9) \quad d(\Sigma(L), \mathfrak{t}_{(L,o)}) = 2\chi(HF_{\text{red}}^+(\Sigma(L))) + 2\tau(\Sigma(L), \mathfrak{t}_{(L,o)}) - \lambda(\Sigma(L)),$$

where λ denotes the Casson–Walker invariant, normalized so that it takes value -2 on the Poincaré sphere oriented as the boundary of the negative E_8 plumbing. Moreover, since L is quasi–alternating $\Sigma(L)$ is an L –space; therefore the first

summand on the right-hand side of (9) vanishes. By [12, Theorem 5.1], when $\det(L) > 0$ we have

$$(10) \quad \lambda(\Sigma(L)) = -\frac{1}{6} \frac{V'_{(L,o)}(-1)}{V_{(L,o)}(-1)} + \frac{1}{4} \sigma(L, o).$$

Therefore, when (L, o) is an oriented quasi-alternating link, Theorem 1 together with equations (9) and (10) yield the statement. \square

ACKNOWLEDGEMENTS

The authors would like to thank the anonymous referee for suggestions which helped to improve the exposition.

REFERENCES

- [1] Abhijit Champanerkar and Ilya Kofman, *Twisting quasi-alternating links*, Proc. Amer. Math. Soc. **137** (2009), no. 7, 2451–2458, DOI 10.1090/S0002-9939-09-09876-1. MR2495282 (2010b:57005)
- [2] A. Champanerkar and P. Ordning, *A note on quasi-alternating Montesinos links*, arXiv preprint 1205.5261.
- [3] Andrew Donald and Brendan Owens, *Concordance groups of links*, Algebr. Geom. Topol. **12** (2012), no. 4, 2069–2093, DOI 10.2140/agt.2012.12.2069. MR3020201
- [4] Joshua Greene, *Homologically thin, non-quasi-alternating links*, Math. Res. Lett. **17** (2010), no. 1, 39–49, DOI 10.4310/MRL.2010.v17.n1.a4. MR2592726 (2011e:57007)
- [5] Joshua Evan Greene and Liam Watson, *Turaev torsion, definite 4-manifolds, and quasi-alternating knots*, Bull. Lond. Math. Soc. **45** (2013), no. 5, 962–972. MR3104988
- [6] S. Jablan and R. Sazdanović, *Quasi-alternating links and odd homology: computations and conjectures*, arXiv preprint 0901.0075.
- [7] Robion C. Kirby, *The topology of 4-manifolds*, Lecture Notes in Mathematics, vol. 1374, Springer-Verlag, Berlin, 1989. MR1001966 (90j:57012)
- [8] W. B. Raymond Lickorish, *An introduction to knot theory*, Graduate Texts in Mathematics, vol. 175, Springer-Verlag, New York, 1997. MR1472978 (98f:57015)
- [9] Paolo Lisca and András I. Stipsicz, *Ozsváth-Szabó invariants and tight contact three-manifolds. II*, J. Differential Geom. **75** (2007), no. 1, 109–141. MR2282726 (2008g:57029)
- [10] C. Manolescu and B. Owens, *A concordance invariant from the Floer homology of double branched covers*, International Mathematics Research Notices **2007** (2007), no. 20, Art. ID rnm077.
- [11] Ciprian Manolescu and Peter Ozsváth, *On the Khovanov and knot Floer homologies of quasi-alternating links*, Proceedings of Gökova Geometry-Topology Conference 2007, Gökova Geometry/Topology Conference (GGT), Gökova, 2008, pp. 60–81. MR2509750 (2010k:57029)
- [12] David Mullins, *The generalized Casson invariant for 2-fold branched covers of S^3 and the Jones polynomial*, Topology **32** (1993), no. 2, 419–438, DOI 10.1016/0040-9383(93)90029-U. MR1217078 (94h:57013)
- [13] Peter Ozsváth and Zoltán Szabó, *Absolutely graded Floer homologies and intersection forms for four-manifolds with boundary*, Adv. Math. **173** (2003), no. 2, 179–261, DOI 10.1016/S0001-8708(02)00030-0. MR1957829 (2003m:57066)
- [14] Peter Ozsváth and Zoltán Szabó, *On the Heegaard Floer homology of branched double-covers*, Advances in Mathematics **194** (2005), no. 1, 1–33.
- [15] Peter Ozsváth and Zoltán Szabó, *Holomorphic triangles and invariants for smooth four-manifolds*, Adv. Math. **202** (2006), no. 2, 326–400, DOI 10.1016/j.aim.2005.03.014. MR2222356 (2007i:57029)
- [16] K. Qazaqzeh, N. Chbili, and B. Qublan, *Characterization of quasi-alternating Montesinos links*, arXiv preprint 1205.4650.
- [17] K. Qazaqzeh, B. Qublan, and A. Jaradat, *A new property of quasi-alternating links*, arXiv preprint 1205.4291.
- [18] R. Rustamov, *Surgery formula for the renormalized Euler characteristic of Heegaard Floer homology*, arXiv preprint math/0409294.

- [19] Nikolai Saveliev, *A surgery formula for the $\bar{\mu}$ -invariant*, *Topology Appl.* **106** (2000), no. 1, 91–102, DOI 10.1016/S0166-8641(99)00075-9. MR1769335 (2001g:57025)
- [20] András I. Stipsicz, *On the $\bar{\mu}$ -invariant of rational surface singularities*, *Proc. Amer. Math. Soc.* **136** (2008), no. 11, 3815–3823, DOI 10.1090/S0002-9939-08-09439-2. MR2425720 (2009e:57024)
- [21] V. G. Turaev, *Classification of oriented Montesinos links via spin structures*, *Topology and geometry—Rohlin Seminar, Lecture Notes in Math.*, vol. 1346, Springer, Berlin, 1988, pp. 271–289, DOI 10.1007/BFb0082779. MR970080 (90d:57013)
- [22] Liam Watson, *A surgical perspective on quasi-alternating links*, *Low-dimensional and symplectic topology, Proc. Sympos. Pure Math.*, vol. 82, Amer. Math. Soc., Providence, RI, 2011, pp. 39–51. MR2768652
- [23] Tamara Widmer, *Quasi-alternating Montesinos links*, *J. Knot Theory Ramifications* **18** (2009), no. 10, 1459–1469, DOI 10.1142/S0218216509007518. MR2583805 (2011d:57025)

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PISA, LARGO BRUNO PONTECORVO 5, 56127 PISA, ITALY

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GLASGOW, UNIVERSITY GARDENS, GLASGOW G12 8QW, UNITED KINGDOM

E-mail address: `b.owens@maths.gla.ac.uk`