

ON MEASURE EXPANSIVE DIFFEOMORPHISMS

M. J. PACIFICO AND J. L. VIEITEZ

(Communicated by Nimish Shah)

ABSTRACT. Let $f : M \rightarrow M$ be a diffeomorphism defined on a compact boundaryless d -dimensional manifold M , $d \geq 2$. In this note we show that diffeomorphisms in a residual subset far from homoclinic tangencies are measure expansive. We also show that surface diffeomorphisms presenting homoclinic tangencies can be C^1 -approximated by non-measure expansive diffeomorphisms.

1. INTRODUCTION

The notion of expansiveness was introduced by Utz in the middle of the twentieth century; see [Ut]. Roughly speaking, a system is expansive if two orbits cannot remain close to each other under the action of the system. This notion is very important in the context of the theory of dynamical systems. For instance, it is responsible for many chaotic properties for homeomorphisms defined on compact spaces; see for instance [Hi, Hi2, Le, Ft, Vi] for more on this. There is extensive literature concerning expansive systems, and a classical result establishes that every hyperbolic f -invariant subset $\Lambda \subset M$ is expansive.

In light of the rich consequences of expansiveness in the dynamics of a system, it is natural to consider another notion of expansiveness. In [Mo] the author introduced the notion of *measure expansiveness*, generalizing the usual concept of expansiveness; see Definition 2.7.

In this paper we prove that there is a residual subset \mathcal{G} of $\text{Diff}^1(M) \setminus \overline{\{\mathcal{HT}\}}$ such that if $f \in \mathcal{G}$, then f is measure-expansive. Here \mathcal{HT} is the subset of $\text{Diff}^1(M)$ presenting a homoclinic tangency (see Definition 2.8).

Moreover we also show that surface diffeomorphisms presenting homoclinic tangencies associated to hyperbolic periodic points can be C^1 -approximated by non-measure-expansive diffeomorphisms.

2. PRELIMINARY RESULTS AND STATEMENT OF THE MAIN RESULT

Let us start with the different definitions of expansiveness we shall deal with. To this end we define for $x \in X$, where (X, d) is a compact metric space, the set

$$(1) \quad \Gamma_\epsilon(x, f) \equiv \{y \in X / d(f^n(x), f^n(y)) \leq \epsilon, n \in \mathbb{Z}\}.$$

Received by the editors February 12, 2013 and, in revised form, March 2, 2013 and June 6, 2013.

2010 *Mathematics Subject Classification*. Primary 37C29, 37D30.

The first author was partially supported by CNPq Brazil, Pronex on Dynamical Systems, FAPERJ, Balzan Research Project of J. Palis.

The second author was partially supported by Grupo de Investigación “Sistemas Dinámicos” CSIC (Universidad de la República), SNI-ANII, PEDECIBA, Uruguay.

We simply write $\Gamma_\epsilon(x)$ instead of $\Gamma_\epsilon(x, f)$ when it is understood to which f we are referring.

2.1. Expansiveness and robust expansiveness.

Definition 2.1. Let $f : X \rightarrow X$ be a homeomorphism defined on a compact metric space (X, d) . We say that f is an expansive homeomorphism if there is $\alpha > 0$ such that $\Gamma_\alpha(x) = \{x\}$ for all $x \in X$. Equivalently, given $x, y \in X$, $x \neq y$, there is $n \in \mathbb{Z}$ such that $\text{dist}(f^n(x), f^n(y)) > \alpha$.

For f a diffeomorphism one is interested in the relation between a given property in the underlying dynamics and its influence on the dynamics on the infinitesimal level, i.e., in the dynamics of the tangent map $Df : TM \rightarrow TM$. Usually one cannot expect that a sole notion on the underlying dynamics can guarantee any interesting feature on the infinitesimal level. Hence we ask for a robust property valid in a whole neighborhood of $f \in \text{Diff}^r(M)$, $r \geq 1$.

Definition 2.2. A compact f -invariant subset Λ is C^r -robustly expansive, $r \geq 1$, if and only if there exists a C^r -neighborhood $\mathcal{U}(f)$ of f such that for all $g \in \mathcal{U}(f)$ there exists a continuation of Λ_g , such that $g|_{\Lambda_g}$ is expansive.

We proved in [PPV, PPSV, SV] that when $\Lambda = H(p, f)$ is a robustly C^1 -expansive homoclinic class associated to a hyperbolic periodic point p , then $H(p, f)$ is hyperbolic (see Subsection 2.3).

2.2. Entropy expansiveness and robust entropy expansiveness. Another notion of expansiveness introduced by Bowen [Bo] is that of an entropy expansive homeomorphism $f : M \rightarrow M$, or h -expansive homomorphism for short.

Let K be a compact invariant subset of M and $\text{dist} : M \times M \rightarrow \mathbb{R}^+$ a distance in M compatible with its Riemannian structure. For $E, F \subset K$, $n \in \mathbb{N}$ and $\delta > 0$ we say that E (n, δ)-spans F with respect to f if for each $y \in F$ there is $x \in E$ such that $\text{dist}(f^j(x), f^j(y)) \leq \delta$ for all $j = 0, \dots, n-1$. Let $r_n(\delta, F)$ denote the minimum cardinality of a set that (n, δ)-spans F . Since K is compact $r_n(\delta, F) < \infty$. We define

$$h(f, F, \delta) \equiv \limsup_{n \rightarrow \infty} \frac{1}{n} \log(r_n(\delta, F))$$

and the topological entropy of f restricted to F as

$$h(f, F) \equiv \lim_{\delta \rightarrow 0} h(f, F, \delta).$$

The last limit exists since $h(f, F, \delta)$ increases as δ decreases to zero.

Definition 2.3. We say that f/K is *entropy-expansive*, or *h-expansive* for short, if and only if there exists $\epsilon > 0$ such that

$$h_f^*(\epsilon) \equiv \sup_{x \in K} h(f, \Gamma_\epsilon(x)) = 0.$$

As for the case of expansiveness we may define a notion of robust h -expansiveness.

Definition 2.4. If $f : M \rightarrow M$ is a C^r -diffeomorphism, $r \geq 1$, and $K \subset M$ is compact invariant, we say that f/K is robustly C^1 -entropy expansive if there is a C^1 -neighborhood \mathcal{U} of f and an open set $U \supset K$ such that if $g \in \mathcal{U}$, then there is $K_g \subset U$ such that g/K_g is entropy expansive. We say that K_g is a continuation of K (not necessarily unique).

We proved in [PaVi, PaVi2, DFPV] that if K is a homoclinic class $H(p, f)$ associated to a hyperbolic periodic point p , then it is robustly h -expansive if and only if it admits a finest dominated splitting

$$T_{H(p)}M = E \oplus F_1 \oplus \cdots \oplus F_k \oplus G$$

with F_j one-dimensional sub-bundles, E uniformly contracting and G uniformly expanding.

Another class of robust entropy expansive diffeomorphisms is that of Morse-Smale diffeomorphisms. Indeed, all of them have topological entropy zero in a robust way.

2.3. Domination, partial hyperbolicity, hyperbolicity. Recall the notion of a dominated splitting for a compact f -invariant subset $\Lambda \subset M$ of a diffeomorphism $f : M \rightarrow M$. It can be seen as a weak form of hyperbolicity.

Definition 2.5. We say that a compact f -invariant set $\Lambda \subset M$ admits a dominated splitting if the tangent bundle $T_\Lambda M$ has a continuous Df -invariant splitting $E \oplus F$ and there exist $C > 0$, $0 < \lambda < 1$, such that

$$\|Df^n|E(x)\| \cdot \|Df^{-n}|F(f^n(x))\| \leq C\lambda^n \quad \forall x \in \Lambda, n \geq 0.$$

When the dominated splitting can be written as a sum

$$(2) \quad T_\Lambda M = E_1 \oplus \cdots \oplus E_j \oplus E_{j+1} \oplus \cdots \oplus E_k,$$

we say that this sum is dominated if for all j the sum

$$(E_1 \oplus \cdots \oplus E_j) \oplus (E_{j+1} \oplus \cdots \oplus E_k)$$

is dominated.

If we cannot decompose in a non-trivial way any sub-bundle E_j appearing in equation (2), we say that it is the *finest* dominated splitting.

Next we define partial hyperbolicity and hyperbolicity.

Definition 2.6. We say that a compact f -invariant set $\Lambda \subset M$ is partially hyperbolic if the tangent bundle $T_\Lambda M$ has a dominated splitting $E^s \oplus F \oplus E^u$ and there exist $C > 0$, $0 < \lambda < 1$, such that for all vectors $v \in E^s$ we have $\|Df^n(v)\| \leq C\lambda^n \|v\|$ for all $n \geq 0$ and for all vectors $v \in E^u$ we have $\|Df^{-n}(v)\| \leq C\lambda^n \|v\|$ for all $n \geq 0$. Vectors in F are less expanded than vectors in E^u and less contracted than vectors in E^s (this follows from domination).

Remark 2.1. In the case that the central sub-bundle F is trivial, we say that Λ is *hyperbolic*.

2.4. Measure expansiveness. Next we introduce the notion of measure expansiveness given in [Mo].

Definition 2.7. Let $f : X \rightarrow X$ be a homeomorphism defined on a compact metric space (X, d) and μ a non-atomic probability measure defined on X (not necessarily f -invariant). We say that f is a μ -expansive homeomorphism if there is $\alpha > 0$ such that $\mu(\Gamma_\alpha(x)) = 0$ for all $x \in X$. Here $\Gamma_\alpha(x)$ is the set defined in equation (1).

We will show that C^1 generic diffeomorphisms far away from homoclinic tangencies are measure expansive. To that end we recall the definition of homoclinic tangencies.

Definition 2.8. A diffeomorphism $f : M \rightarrow M$ exhibits a homoclinic tangency if there is a hyperbolic periodic orbit \mathcal{O} whose invariant manifolds $W^s(\mathcal{O})$ and $W^u(\mathcal{O})$ have a non-transverse intersection.

We set \mathcal{HT} for the subset of $\text{Diff}^1 M$ constituted of diffeomorphisms presenting a homoclinic tangency. Given a subset A of $\text{Diff}^1 M$ we use the notation \overline{A} for the closure of A in $\text{Diff}^1 M$.

The main results in this paper are the following theorems:

Theorem A. *Let $f : M \rightarrow M$ be a C^1 -diffeomorphism defined on a compact manifold M . There is a \mathcal{G} residual subset of $\text{Diff}^1 M \setminus \overline{\mathcal{HT}}$ such that for any Borel probability measure μ (invariant by f or not) absolutely continuous with respect to Lebesgue, we have that there is $\delta > 0$ such that $\mu(\Gamma_\delta(x)) = 0$ for all $x \in M$. In particular, f is μ -expansive.*

Theorem B. *Let $f : M \rightarrow M$ be a C^1 -diffeomorphism defined on a compact surface M having a homoclinic tangency associated to a hyperbolic periodic orbit \mathcal{O} . Then there is an arbitrarily small C^1 -perturbation of f giving a diffeomorphism $F : M \rightarrow M$ which is not measure-expansive.*

3. PROOF OF THEOREM A

We start by stating some results proved elsewhere that will be used in the proof. Let $X = \text{Diff}^1(M) \setminus \overline{\mathcal{HT}}$. The following result is Theorem 1.1 of [CSY].

Theorem 3.1. *The diffeomorphism f in a dense \mathcal{G}_δ subset $\mathcal{G} \subset \text{Diff}^1(M) \setminus \overline{\mathcal{HT}}$ has the following properties.*

- (1) *Any aperiodic class \mathcal{C} is partially hyperbolic with a one-dimensional central bundle. Moreover, the Lyapunov exponent along E^c of any invariant measure supported on \mathcal{C} is zero.*
- (2) *Any homoclinic class $H(p)$ has a partially hyperbolic structure*

$$T_{\mathcal{C}}M = E^s \oplus E_1^c \oplus \dots \oplus E_k^c \oplus E^u.$$

Moreover, the minimal stable dimension of the periodic orbits of $H(p)$ is $\dim(E^s)$ or $\dim(E^s) + 1$. Similarly, the maximal stable dimension of the periodic orbits of $H(p)$ is $\dim(E^s) + k$ or $\dim(E^s) + k - 1$. For every i , $1 \leq i \leq k$, there exist periodic points in $H(p)$ whose Lyapunov exponent along E_i^c is arbitrarily close to 0. In particular, if $f \in \mathcal{G}$, then f is partially hyperbolic.

For $x \in \Lambda$ and $i \in \{1, \dots, k\}$ let us denote

$$(3) \quad E^{cs,i}(x) := E^s(x) \oplus E_1^c(x) \oplus \dots \oplus E_i^c(x); \quad E^{cu,i}(x) := E_i^c(x) \oplus \dots \oplus E_k^c(x) \oplus E^u(x).$$

We also let $E^{cs,0} = E^s$ and $E^{cu,k+1} = E^u$ and write $s = \dim(E^s)$ and $u = \dim(E^u)$.

Let us recall the properties of fake central manifolds \widehat{W}^{cs} due to Burns and Wilkinson [BW]; see also [DFPV].

Proposition 3.2. *Let $f : M \rightarrow M$ be a C^1 -diffeomorphism and Λ a compact f -invariant set with a partially hyperbolic splitting,*

$$T_\Lambda M = E^s \oplus E_1^c \oplus \dots \oplus E_k^c \oplus E^u.$$

Let $E^{cs,i}$ and $E^{cu,i}$ be as in equation (3) and consider their extensions $\tilde{E}^{cs,i}$ and $\tilde{E}^{cu,i}$ to a small neighborhood of Λ .

Then for any $\epsilon > 0$ there exist constants $R > r > r_1 > 0$ such that, for every $p \in \Lambda$, the neighborhood $B(p, r)$ is foliated by foliations $\widehat{W}^u(p)$, $\widehat{W}^s(p)$, $\widehat{W}^{cs,i}(p)$, and $\widehat{W}^{cu,i}(p)$, $i \in \{1, \dots, k\}$, such that for each $\beta \in \{u, s, (cs, i), (cu, i)\}$ the following properties hold.

- (i) Almost tangency of the invariant distributions. For each $q \in B(p, r)$, the leaf $\widehat{W}_p^\beta(q)$ is C^1 , and the tangent space $T_q \widehat{W}_p^\beta(q)$ lies in a cone of radius ϵ about $\tilde{E}^\beta(q)$.
- (ii) Coherence. \widehat{W}_p^s subfoliates $\widehat{W}_p^{cs,i}$ and \widehat{W}_p^u subfoliates $\widehat{W}_p^{cu,i}$ for each $i \in \{1, \dots, k\}$.
- (iii) Local invariance. For each $q \in B(p, r_1)$ we have

$$f(\widehat{W}_p^\beta(q, r_1)) \subset \widehat{W}_{f(p)}^\beta(f(q)) \text{ and } f^{-1}(\widehat{W}_p^\beta(q, r_1)) \subset \widehat{W}_{f^{-1}(p)}^\beta(f^{-1}(q));$$
 here $\widehat{W}_p^\beta(q, r_1)$ is the connected component of $\widehat{W}_p^\beta(q) \cap B(q, r_1)$ containing q .
- (iv) Uniqueness. $\widehat{W}_p^s(p) = W^s(p, r)$ and $\widehat{W}_p^u(p) = W^u(p, r)$.

Proof. See [BW, Section 3]. □

Given $j \in \{1, \dots, k\}$, using Proposition 3.2, we consider a small r and the submanifold

$$(4) \quad \widetilde{W}^{cs,j}(x) = \bigcup_{z \in \gamma_j(x)} \widehat{W}_x^{cs,j-1}(z, r).$$

This submanifold has dimension $s + j$ and is transverse to $\widehat{W}_x^{cu,j+1}(z)$ for all z close to x . Note that $\widetilde{W}^{cs,1}(x)$ is foliated by stable manifolds (recall that $\widehat{W}_x^{cs,0}(z) \subset W^s(z)$).

The next two lemmas follow straightforwardly from the fact that the angles between unitary vectors in the cone fields $\mathcal{C}(E^{cs,j})$ and $\mathcal{C}(E^{cu,j+1})$ are uniformly bounded away from zero; see [DFPV].

Lemma 3.3. *There is $\kappa > 0$ such that for every $j \in \{1, \dots, k\}$ and every $\delta > 0$ small enough the following property holds.*

For every $x \in \Lambda$, every $y \in B_\delta(x)$, and every local submanifold $N(x)$ of dimension $s + j$ tangent to the cone field $\mathcal{C}(E^{cs,j})$ containing x and $M(y)$ of dimension $(k - j) + u$ tangent to the cone field $\mathcal{C}(E^{cu,j+1})$ containing y , one has that $N(x) \cap M(y)$ is contained in $B_{\kappa\delta}(x)$.

Lemma 3.4. *There is $\kappa > 0$ such that for every $j \in \{1, \dots, k\}$ and every $\delta > 0$ small enough the following property holds.*

Take any $x \in \Lambda$ and the local manifold $\widetilde{W}^{cs,j}(x)$ in (4). For every $y \in B_\delta(x) \cap \widetilde{W}^{cs,j}(x)$ one has that $\gamma_j(x) \cap \widehat{W}_x^{cs,j-1}(y)$ is contained in $B_{\kappa\delta}(x)$. Here $\gamma_j(x)$ is a central curve with end points x and y .

As a consequence of Theorem 3.1 and Lemmas 3.3 and 3.4 we have

Theorem 3.5. *Let μ be a Borel probability measure defined on M and let $f \in \mathcal{G}$ where \mathcal{G} is as in Theorem 3.1. Then there is $\delta > 0$ such that $\mu(\Gamma_\delta(x)) = 0$ for all $x \in M$.*

Proof. Let μ be a Borel probability measure of M and choose $x \in \Omega(f)$. Then there is either an aperiodic class or a homoclinic class H in $\omega(x)$. In any case H is partially hyperbolic, since we are assuming that $f \in \mathcal{G}$.

Let $T_H M = E^s \oplus E_1^c \oplus \dots \oplus E_k^c \oplus E^u$, with E^s uniformly contracting and E^u uniformly expanding. Assume $\theta > 1$ is the minimum rate of expansion of E^u for $z \in \Lambda$. Let $c > 0$ be such that $(1 - c)\theta > 1$ and find $\delta > 0$ less than or equal to that of Lemmas 3.3 and 3.4 and also less than $r/2$ where $r > 0$ is given by Proposition 3.2, such that if

$$\text{dist}(x, y) \leq (\kappa + 1)\delta, \quad \text{then} \quad 1 - c \leq \frac{\|Df|_{E^u(y)}\|}{\|Df|_{E^u(x)}\|} \leq 1 + c.$$

For this choice of δ it holds that $\mu(\Gamma_\delta(x)) = 0$, for, if $y \in \Gamma_\delta(x)$, then letting y_u be the projection of y into $\widehat{W}^u(x)$ along $\widehat{W}^{cs,k}(y)$ (if it were the case that $\text{dist}(x, y_u) > 0$), then setting $\theta' = (1 - c)\theta$, we get for $n \geq 1$,

$$\text{dist}(f^n(x), f^n(y_u)) \geq (\theta')^n \text{dist}(x, y_u).$$

Since $\theta' > 1$ eventually $\text{dist}(f^n(x), f^n(y_u)) > \kappa \delta$, and hence by Lemmas 3.3 and 3.4 we obtain $\text{dist}(f^n(x), f^n(y)) > \delta$, contradicting the fact that $y \in \Gamma_\delta(x)$.

Thus $\Gamma_\delta(x)$ is contained in $\widehat{W}^{cs,k}(x)$. By backward iteration we also get that $\Gamma_\delta(x) \subset \widehat{W}^{cu,0}(x)$.

This implies that $\mu(\Gamma_\delta(x)) = 0$, finishing the proof for $x \in \Omega(x)$.

Assume now that x is any point in M . Then by forward iteration we find $N > 0$ such that $f^n(x) \in B(\omega(x), r/4)$, where $r > 0$ is as in Proposition 3.2. Since $\omega(x) \subset \Omega(f)$ we have that there is either an aperiodic class or a homoclinic class H which is partially hyperbolic such that $\omega(x) \subset H$, and therefore $f^n(x) \in B(H, r/4)$ for $n \geq N$. The result follows as in the case $x \in \Omega(f)$. Similarly, by backward iteration we find $N' > 0$ such that $f^{-n}(x) \in B(\alpha(x), r/4)$ for $n \geq N'$.

We may conclude, using similar estimations as in the case $x \in \Omega(f)$, that $\mu(\Gamma_\delta(x)) = 0$. □

Theorem A is an immediate consequence of Theorem 3.5 .

Remark 3.6. For the validity of Theorem 3.5 it is enough to have that E^s is uniformly contracting for the α -limit of x or that E^u is uniformly expanding for the ω -limit of x .

4. SURFACE DIFFEOMORPHISMS IN \mathcal{HT}

In the remainder of the paper M is a compact boundaryless surface.

Let $f : M \rightarrow M$ be a diffeomorphism and assume that f exhibits a homoclinic tangency associated to a hyperbolic periodic point p of f .

4.1. Horseshoes with positive Lebesgue measure. It is proved in [Bo2] that there is a C^1 horseshoe with positive Lebesgue measure. In [RY] is constructed such a horseshoe fattening up an invariant horseshoe Λ to have positive Lebesgue measure as Bowen did. They obtain this fattened horseshoe by modifying a diffeomorphism f defined in a square $B = [0, 1] \times [0, 1]$ so that $f|B$ gives a linear evenly spaced full shift on two symbols; see [RY, §1]. The perturbed diffeomorphism is C^1 close to the original one; see [RY, §3 and §4]. After that they embed Λ in a C^1 -diffeomorphism F defined on a surface [RY, §5]. Although this construction is

made to embed the horseshoe on a C^1 -Anosov diffeomorphism, the same can be done for any diffeomorphism.

Remark 4.1. Perhaps it is worthwhile to note that it is crucial that we are working in the C^1 -topology. Bowen [Bo1] proved that C^2 -diffeomorphisms have no horseshoes with positive volume.

4.2. Proof of Theorem B. We now make use of the construction in [RY] to prove that arbitrarily near a diffeomorphism exhibiting a homoclinic tangency there is one which is not measure-expansive.

We start by establishing some auxiliary lemmas proved elsewhere.

Lemma 4.2. *Given a C^1 -diffeomorphism $f : M \rightarrow M$ with a homoclinic tangency associated to a hyperbolic periodic point p , there is a C^1 near diffeomorphism f_1 presenting a flat homoclinic tangency; i.e., there is a small arc J contained in $W^s(p, f_1) \cap W^u(p, f_1)$.*

Proof. See [PaVi2, Proposition 2.6]. □

Lemma 4.3. *Given a C^1 -diffeomorphism $f_1 : M \rightarrow M$ with a flat homoclinic tangency associated to a hyperbolic periodic point p , there is a C^1 near diffeomorphism f_2 presenting a sequence of horseshoes $\widehat{\Lambda}_n$ such that for all $k \in \mathbb{Z}$, $\text{diam}(f^k(\widehat{\Lambda}_n)) < r_n$ with $r_n \rightarrow 0$ when $n \rightarrow \infty$.*

Proof. The proof is essentially the same as that of [PaVi2, Subsection 2.2]. □

Proposition 4.4. *Let $f_2 : M \rightarrow M$ as in the thesis of Lemma 4.3. There is a C^1 -diffeomorphism $F : M \rightarrow M$ arbitrarily near f_2 presenting a sequence of horseshoes Λ_n such that the Lebesgue measure $\mu(\Lambda_n) > 0$ and $\text{diam}(\Lambda_n) < 2r_n$, where r_n is as in Lemma 4.3.*

Proof. We profit from the construction made in [RY]. In fact we do not need to take care for the perturbed diffeomorphism to be Anosov, as is the case in [RY]. Hence, in our case, to fit the construction in the global picture of the perturbations is easier than in [RY, §5]. Since the support of the perturbation needed to fatten the horseshoe $\widehat{\Lambda}_n$ is contained in a box $B_n \supset \widehat{\Lambda}_n$ such that $\lim_{n \rightarrow \infty} \text{diam}(B_n) = 0$ (see [RY, §3]), it can be taken disjointly from the support of the previous perturbations needed to fatten $\widehat{\Lambda}_j$ for $j = 1, \dots, n-1$ (see [RY, §2 and §4]). From this it follows that F is C^1 -close to f_2 and has the desired sequence of horseshoes Λ_n with positive Lebesgue measure.

Moreover, the construction of Λ_n gives that the diameter of Λ_n is about the same as that of $\widehat{\Lambda}_n$, so that we can assure that $\text{diam}(\Lambda_n) < 2r_n$ from $\text{diam}(\widehat{\Lambda}_n) < r_n$. □

As a consequence we have

Theorem 4.5. *Let M be a smooth compact surface. Given a C^1 -diffeomorphism $f : M \rightarrow M$ exhibiting a homoclinic tangency associated to a hyperbolic periodic point p , it is C^1 -approximated by a diffeomorphism $F : M \rightarrow M$ such that F is not measure expansive with respect to any absolutely continuous invariant measure with respect to Lebesgue.*

Proof. Let $F : M \rightarrow M$ be the C^1 -diffeomorphism constructed as in Proposition 4.4 above. Then for every horseshoe Λ_n associated to F there is a hyperbolic periodic point $p_n \in \Lambda_n$ such that $\mu(\Gamma_{2r_n}(p_n)) \geq \mu(\Lambda_n) > 0$, where $\mu \ll \text{Leb}$ and $f^*\mu = \mu$. Since $r_n \rightarrow 0$, when $n \rightarrow \infty$ the proof follows. \square

Theorem 4.5 gives the proof of Theorem B.

ACKNOWLEDGEMENTS

The second author thanks IMPA and Instituto de Matematica, Universidade Federal do Rio de Janeiro for their kind hospitality during the preparation of this paper.

REFERENCES

- [Bo] Rufus Bowen, *Entropy-expansive maps*, Trans. Amer. Math. Soc. **164** (1972), 323–331. MR0285689 (44 #2907)
- [Bo1] R. Bowen, *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*, Lecture Notes in Mathematics, 470, Springer, 1975. MR0442989
- [Bo2] Rufus Bowen, *A horseshoe with positive measure*, Invent. Math. **29** (1975), no. 3, 203–204. MR0380890 (52 #1787)
- [BW] Keith Burns and Amie Wilkinson, *On the ergodicity of partially hyperbolic systems*, Ann. of Math. (2) **171** (2010), no. 1, 451–489, DOI 10.4007/annals.2010.171.451. MR2630044 (2011g:37075)
- [CSY] S. Crovisier, M. Sambarino, D. Yang, *Partial Hyperbolicity and Homoclinic Tangencies*, arXiv 1103.0869v1, 2011.
- [DFPV] Lorenzo J. Díaz, Todd Fisher, Maria José Pacifico, and José L. Vieitez, *Entropy-expansiveness for partially hyperbolic diffeomorphisms*, Discrete Contin. Dyn. Syst. **32** (2012), no. 12, 4195–4207, DOI 10.3934/dcds.2012.32.4195. MR2966742
- [Ft] Albert Fathi, *Expansiveness, hyperbolicity and Hausdorff dimension*, Comm. Math. Phys. **126** (1989), no. 2, 249–262. MR1027497 (90m:58159)
- [Hi] Koichi Hiraide, *Expansive homeomorphisms with the pseudo-orbit tracing property of n -tori*, J. Math. Soc. Japan **41** (1989), no. 3, 357–389, DOI 10.2969/jmsj/04130357. MR999503 (90h:58062)
- [Hi2] Koichi Hiraide, *Expansive homeomorphisms of compact surfaces are pseudo-Anosov*, Osaka J. Math. **27** (1990), no. 1, 117–162. MR1049828 (91b:58184)
- [Le] Jorge Lewowicz, *Expansive homeomorphisms of surfaces*, Bol. Soc. Brasil. Mat. (N.S.) **20** (1989), no. 1, 113–133, DOI 10.1007/BF02585472. MR1129082 (92i:58139)
- [Mo] C. A. Morales, *Measure expansive systems*, preprint, IMPA, D083, 2011.
- [PaVi] María José Pacifico and José L. Vieitez, *Entropy-expansiveness and domination for surface diffeomorphisms*, Rev. Mat. Complut. **21** (2008), no. 2, 293–317. MR2441955 (2009m:37090)
- [PaVi2] M. J. Pacifico and J. L. Vieitez, *Robust entropy expansiveness implies generic domination*, Nonlinearity **23** (2010), no. 8, 1971–1990, DOI 10.1088/0951-7715/23/8/009. MR2669634 (2011k:37047)
- [PPSV] M. J. Pacifico, E. R. Pujals, M. Sambarino, and J. L. Vieitez, *Robustly expansive codimension-one homoclinic classes are hyperbolic*, Ergodic Theory Dynam. Systems **29** (2009), no. 1, 179–200, DOI 10.1017/S0143385708000175. MR2470632 (2009k:37070)
- [PPV] M. J. Pacifico, E. R. Pujals, and J. L. Vieitez, *Robustly expansive homoclinic classes*, Ergodic Theory Dynam. Systems **25** (2005), no. 1, 271–300, DOI 10.1017/S0143385704000203. MR2122923 (2006k:37041)
- [RY] Clark Robinson and Lai Sang Young, *Nonabsolutely continuous foliations for an Anosov diffeomorphism*, Invent. Math. **61** (1980), no. 2, 159–176, DOI 10.1007/BF01390119. MR590160 (81m:58061)
- [SV] Martín Sambarino and José L. Vieitez, *Robustly expansive homoclinic classes are generically hyperbolic*, Discrete Contin. Dyn. Syst. **24** (2009), no. 4, 1325–1333, DOI 10.3934/dcds.2009.24.1325. MR2505706 (2010f:37052)

- [Vi] José L. Vieitez, *Lyapunov functions and expansive diffeomorphisms on 3D-manifolds*, Ergodic Theory Dynam. Systems **22** (2002), no. 2, 601–632, DOI 10.1017/S0143385702000305. MR1898808 (2003b:37049)
- [Ut] W. R. Utz, *Unstable homeomorphisms*, Proc. Amer. Math. Soc. **1** (1950), 769–774. MR0038022 (12,344b)

INSTITUTO DE METEMATICA, UNIVERSIDADE FEDERAL DO RIO DE JANEIRO, C. P. 68.530, CEP 21.945-970, RIO DE JANEIRO, RJ, BRAZIL
E-mail address: `pacifico@im.ufrj.br`

INSTITUTO DE MATEMATICA, FACULTAD DE INGENIERIA, UNIVERSIDAD DE LA REPUBLICA, CC30, CP 11300, MONTEVIDEO, URUGUAY
E-mail address: `jvieitez@fing.edu.uy`