ON SYZYGIES OF LINEAR SECTIONS

EUISUNG PARK

(Communicated by Irena Peeva)

ABSTRACT. In this paper, we study how the minimal free resolution of a closed subscheme $X \subset \mathbb{P}^r$ relates to that of a linear section $X \cap \Lambda \subset \Lambda = \mathbb{P}^s$ ($0 < s < r$). Our main result implies that the shape of the final non-zero row of the Betti diagram of X is preserved under taking the zero-dimensional and one-dimensional linear sections.

1. INTRODUCTION

Throughout this paper, we will work with projective schemes defined over an algebraically closed field K . We denote by $\mathbb{P}^r$ the projective r -space over K .

For a closed subscheme $X \subset \mathbb{P}^r$, let $\mathcal{I}_X$ and I_X be respectively the ideal sheaf and the homogeneous ideal of X in the homogeneous coordinate ring R of $\mathbb{P}^r$. We will denote by $m(X)$ the least integer m such that I_X is generated by forms of degree $\leq m$. Following Eisenbud et al. [EGHP], we will say that X satisfies the *condition* $N_{d,p}$, for some $d \geq 2$ and $p \geq 1$, if in the minimal free resolution

$$\cdots \rightarrow F_i \rightarrow \cdots \rightarrow F_2 \rightarrow F_1 \rightarrow I_X \rightarrow 0$$

of I_X as a graded R -module, the free modules F_i are generated by homogeneous elements of degree at most $d+i-1$ for $1 \leq i \leq p$. For example, X satisfies condition $N_{d,1}$ if and only if $m(X) \leq d$. Also, X satisfies condition $N_{d,p}$ for some $p \geq r+1$ if and only if it is d -regular in the sense of Castelnuovo-Mumford.

The aim of this article is to study how the minimal free resolution of $X \subset \mathbb{P}^r$ relates to that of the scheme-theoretic intersection $X \cap \Lambda \subset \Lambda = \mathbb{P}^s$ where $\Lambda = \mathbb{P}^s \subset \mathbb{P}^r$ ($0 < s < r$), is a linear subspace.

Maybe the first result in this direction is the well-known and elementary fact that if $s = 1$ and Λ is not contained in X then it holds that

$$(1.1) \quad \text{reg}(X) \geq m(X) \geq \text{length}(X \cap \Lambda) (= \text{reg}(X \cap \Lambda)).$$

For example, if Λ is a tri-secant line to X then I_X cannot be generated by quadrics. More recently, (1.1) was widely extended by Theorem 1.1 in [EGHP] which states that if X satisfies condition $N_{d,p}$, $s \leq p$ and $\dim(X \cap \Lambda) \leq 1$, then $X \cap \Lambda$ is d -regular. In particular, if $d = 2$ and $X \cap \Lambda$ is finite, then $\text{length}(X \cap \Lambda)$ is at most $s+1$. One can easily see that (1.1) is $p = 1$ case of this result.

Received by the editors October 31, 2012 and, in revised form, January 3, 2013.

2010 *Mathematics Subject Classification.* Primary 14N05, 13D02.

Key words and phrases. Minimal free resolution, Linear section.

Along this line, our main result in this paper is:

Theorem 1.1. *Let $X \subset \mathbb{P}^r$ be a closed subscheme which is m -regular and satisfies condition $N_{m-1,p}$ for some $p \geq 1$. Then for any linear subspace $\Lambda = \mathbb{P}^s \subset \mathbb{P}^r$ ($0 < s < r$) with $\dim X \cap \Lambda \leq 1$, the intersection $X \cap \Lambda \subset \mathbb{P}^s$ is m -regular and satisfies condition $N_{m-1,p}$.*

Briefly speaking, our result means that the shape of the final non-zero row of the Betti diagram of X is preserved under taking zero-dimensional and one-dimensional linear sections.

To compare our result with Theorem 1.1 in [EGHP], let us consider a smooth linearly normal curve $\mathcal{C} \subset \mathbb{P}^{d-g}$ of genus $g \geq 2$ and degree $d \geq 2g + 2$. Then $\mathcal{C}$ is 3-regular and satisfies condition $N_{2,d-2g-1}$. Furthermore, the following two statements are equivalent (cf. Theorem 2 in [GL]):

- (i) $\mathcal{C}$ fails to satisfy condition $N_{2,d-2g}$.
- (ii) Either
 - (ii.a) $\mathcal{C}$ is a hyperelliptic curve or
 - (ii.b) it admits a $(d - 2g + 2)$ -secant $(d - 2g)$ -plane.

The implication (ii.b) $\implies$ (i) is reproved by Theorem 1.1 in [EGHP]. On the other hand, if $\mathcal{C}$ is a hyperelliptic curve then it happens that there exists no $(d - 2g + 2)$ -secant $(d - 2g)$ -plane to $\mathcal{C}$. So, one can naturally ask if there is a geometric obstruction of condition $N_{2,d-2g}$ for a hyperelliptic curve $\mathcal{C}$. Along this line, it turns out to be always true that if $\mathcal{C}$ is a hyperelliptic curve then there exists a proper linear subspace Λ of $\mathbb{P}^{d-g}$ such that the intersection $\mathcal{C} \cap \Lambda$ fails to satisfy condition $N_{2,d-2g}$ (see Section 3). Thus $\mathcal{C}$ fails to satisfy condition $N_{2,d-2g}$ by Theorem 1.1. In consequence, we have

Corollary 1.2. *Let $\mathcal{C} \subset \mathbb{P}^{d-g}$ be a smooth linearly normal curve of genus $g \geq 2$ and degree $d \geq 2g + 2$. Then $\mathcal{C}$ fails to satisfy condition $N_{2,d-2g}$ if and only if there exists a proper linear subspace Λ of $\mathbb{P}^{d-g}$ such that the intersection $\mathcal{C} \cap \Lambda$ fails to satisfy condition $N_{2,d-2g}$.*

2. PROOF OF THEOREM 1.1

This section is devoted to giving a proof of Theorem 1.1.

Definition and Remark 2.1. (1) For a nonzero finitely generated graded R -module $M = \bigoplus_{\ell \in \mathbb{Z}} M_\ell$, let

$$\cdots \rightarrow F_i \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow M \rightarrow 0$$

be the minimal free resolution where $F_i = \bigoplus_{j \geq 0} R(-i-j)^{\beta_{i,j}(M)}$. The integer $\beta_{i,j}(M)$ is called the (i, j) -th graded Betti number of M . Note that

$$\beta_{i,j}(M) := \dim_K \operatorname{Tor}_i^R(M, K)_{i+j}.$$

(2) Let $\mathcal{F}$ be a nonzero coherent sheaf on $\mathbb{P}^r$, and set $F := \bigoplus_{n \geq 0} H^0(\mathbb{P}^r, \mathcal{F}(n))$. For $j \geq 1$, there is an exact sequence

$$0 \rightarrow \operatorname{Tor}_i^R(F, K)_{i+j} \rightarrow H^1(\mathbb{P}^r, \bigwedge^{i+1} \mathcal{M} \otimes \mathcal{F}(j-1)) \rightarrow \bigwedge^{i+1} V \otimes H^1(\mathbb{P}^r, \mathcal{F}(j-1))$$

where $\mathcal{M} = \Omega_{\mathbb{P}^r}(1)$ and $V = H^0(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(1))$. In particular, we have

$$\beta_{i,j}(F) = h^1(\mathbb{P}^r, \bigwedge^{i+1} \mathcal{M} \otimes \mathcal{F}(j-1))$$

if $h^1(\mathbb{P}^r, \mathcal{F}(j-1)) = 0$. For details, see Theorem 5.8 in [E].

(3) Let $X \subset \mathbb{P}^r$ be a closed subscheme. We denote by $\beta_{i,j}(X)$ the (i, j) -th graded Betti number of the homogeneous coordinate ring of X . If X is m -regular, then by (2) we have

$$(2.1) \quad \beta_{i,j}(X) = h^1(\mathbb{P}^r, \bigwedge^i \mathcal{M} \otimes \mathcal{I}_X(j)) \quad \text{for all } i \geq 1 \text{ and } j \geq m-1.$$

In particular, $h^1(\mathbb{P}^r, \bigwedge^i \mathcal{M} \otimes \mathcal{I}_X(j)) = \beta_{i,j}(X) = 0$ for all $i \geq 0$ and $j \geq m$.

Proposition 2.2. *For a closed subscheme $X \subset \mathbb{P}^r$, suppose that X is m -regular and satisfies condition $N_{m-1,p}$ for some $p \geq 1$. Then*

(1) $H^s(\mathbb{P}^r, \bigwedge^i \mathcal{M} \otimes \mathcal{I}_X(j)) = 0$ if either

$$\begin{cases} 1 \leq s \leq i & \text{and } j \geq m-s+1; & \text{or} \\ 1 \leq s \leq i, j = m-s & \text{and } 1 \leq i-s+1 \leq p; & \text{or} \\ s \geq i+1 & \text{and } j \geq m-s. \end{cases}$$

(2) Let $\mathcal{I}_\Lambda$ be the ideal sheaf of Λ in $\mathbb{P}^r$. If $\dim(X \cap \Lambda) \leq 1$, then

$$H^2(\mathbb{P}^r, \bigwedge^i \mathcal{M} \otimes \mathcal{I}_\Lambda \otimes \mathcal{I}_X(j)) = 0$$

whenever $j \geq m$ or $j = m-1$ and $1 \leq i \leq p$.

Proof. (1) We use induction on i and s . For $i = 0$, the statements hold since X is m -regular. Also for $s = 1$, see (2.1). Now, suppose that $i \geq 1$ and $s \geq 2$. By the m -regularity of X and the induction hypothesis, we have

$$H^s(\mathbb{P}^r, \mathcal{I}_X(j)) = H^{s-1}(\mathbb{P}^r, \bigwedge^{i-1} \mathcal{M} \otimes \mathcal{I}_X(j+1)) = 0.$$

Then we get the desired vanishing of $H^s(\mathbb{P}^r, \bigwedge^i \mathcal{M} \otimes \mathcal{I}_X(j))$ by using the short exact sequence

$$0 \rightarrow \bigwedge^i \mathcal{M} \otimes \mathcal{I}_X(j) \rightarrow \bigwedge^i V \otimes \mathcal{I}_X(j) \rightarrow \bigwedge^{i-1} \mathcal{M} \otimes \mathcal{I}_X(j+1) \rightarrow 0.$$

For the above exact sequence, see (5.1) in [E].

(2) Let I_Λ be the homogeneous ideal of Λ in R and $U = H^0(\mathbb{P}^r, \mathcal{I}_\Lambda(1))$. Then the sheafification of the minimal free resolution of I_Λ is of the form

$$\cdots \rightarrow \bigwedge^2 U \otimes \mathcal{O}_{\mathbb{P}^r}(-2) \rightarrow U \otimes \mathcal{O}_{\mathbb{P}^r}(-1) \rightarrow \mathcal{I}_\Lambda \rightarrow 0.$$

Consider the complex obtained by tensoring $\mathcal{I}_X$ to this long exact sequence:

$$\cdots \rightarrow \bigwedge^2 U \otimes \mathcal{I}_X(-2) \rightarrow U \otimes \mathcal{I}_X(-1) \rightarrow \mathcal{I}_\Lambda \otimes \mathcal{I}_X \rightarrow 0.$$

Since this is exact away from $X \cap \Lambda$ and $\dim(X \cap \Lambda) \leq 1$, we get the desired vanishing if

$$H^{2+t}(\mathbb{P}^r, \bigwedge^i \mathcal{M} \otimes \mathcal{I}_X(j-t-1)) = 0$$

for all $t \geq 0$ when $j \geq m$ or $j = m - 1$ and $1 \leq i \leq p$ (cf. Proposition B.1.2 (ii) in [L]). Thus the proof is completed by (1). $\square$

Proof of Theorem 1.1. Let $\mathcal{I}_{X \cap \Lambda}$ be the ideal sheaf of $X \cap \Lambda$ in Λ and let $\mathcal{M}_\Lambda = \Omega_\Lambda(1)$. By Definition and Remark 2.1, $X \cap \Lambda$ is m -regular and satisfies condition $N_{m-1,p}$ if

$$(2.2) \quad H^1(\Lambda, \bigwedge^i \mathcal{M}_\Lambda \otimes \mathcal{I}_{X \cap \Lambda}(j)) = 0 \quad \text{for } j \geq m \text{ and for } j = m - 1 \text{ and } 1 \leq i \leq p.$$

Also, since $\mathcal{M} \otimes \mathcal{O}_\Lambda$ is isomorphic to the direct sum of $\mathcal{M}_\Lambda$ and copies of $\mathcal{O}_\Lambda$, we obtain (2.2) by showing the vanishing

$$(2.3) \quad H^1(\mathbb{P}^r, \bigwedge^i \mathcal{M} \otimes \mathcal{I}_{X \cap \Lambda}(j)) = 0 \quad \text{for } j \geq m \text{ and for } j = m - 1 \text{ and } 1 \leq i \leq p.$$

To this aim, we use the tensor product of $\mathcal{I}_X$ with the short exact sequence $0 \rightarrow \mathcal{I}_\Lambda \rightarrow \mathcal{O}_{\mathbb{P}^r} \rightarrow \mathcal{O}_\Lambda \rightarrow 0$. Let $\mathcal{K}$ and $\mathcal{K}'$ denote the kernels of the surjective homomorphisms $\mathcal{I}_X \otimes \mathcal{O}_\Lambda \rightarrow \mathcal{I}_{X \cap \Lambda}$ and $\mathcal{I}_\Lambda \otimes \mathcal{I}_X \rightarrow \mathcal{I}_\Lambda \mathcal{I}_X$, respectively. By chopping the sequence $\mathcal{I}_\Lambda \otimes \mathcal{I}_X \rightarrow \mathcal{I}_X \rightarrow \mathcal{I}_X \otimes \mathcal{O}_\Lambda$ into short exact sequences in the usual way, we get the following chain of exact sequences:

$$\begin{array}{ccccccc} 0 & & & & & & \\ & \downarrow & & & & & \\ & \mathcal{K}' & & & 0 & & \\ & \downarrow & & & \downarrow & & \\ & \mathcal{I}_\Lambda \otimes \mathcal{I}_X & & & \mathcal{K} & & \\ & \downarrow & & & \downarrow & & \\ 0 \rightarrow & \mathcal{I}_\Lambda \mathcal{I}_X & \rightarrow & \mathcal{I}_X & \rightarrow & \mathcal{I}_X \otimes \mathcal{O}_\Lambda & \rightarrow 0 \\ & \downarrow & & & \downarrow & & \\ & 0 & & & \mathcal{I}_{X \cap \Lambda} & & \\ & & & & \downarrow & & \\ & & & & 0. & & \end{array}$$

Since $\text{Supp } \mathcal{K}$ and $\text{Supp } \mathcal{K}'$ are both contained in $X \cap \Lambda$ and $\dim(X \cap \Lambda) \leq 1$, it holds that for all $i \geq 0$ and $j \in \mathbb{Z}$,

- (i) $H^2(\bigwedge^i \mathcal{M} \otimes \mathcal{I}_\Lambda \otimes \mathcal{I}_X(j)) \cong H^2(\bigwedge^i \mathcal{M} \otimes \mathcal{I}_\Lambda \mathcal{I}_X(j))$,
- (ii) $H^1(\bigwedge^i \mathcal{M} \otimes \mathcal{I}_X(j)) \rightarrow H^1(\bigwedge^i \mathcal{M} \otimes \mathcal{I}_X \otimes \mathcal{O}_\Lambda(j)) \rightarrow H^2(\bigwedge^i \mathcal{M} \otimes \mathcal{I}_\Lambda \mathcal{I}_X(j))$ is exact, and
- (iii) $H^1(\bigwedge^i \mathcal{M} \otimes \mathcal{I}_X \otimes \mathcal{O}_\Lambda(j)) \rightarrow H^1(\bigwedge^i \mathcal{M} \otimes \mathcal{I}_{X \cap \Lambda}(j))$ is surjective.

Also Proposition 2.2 shows that

$$H^1(\mathbb{P}^r, \bigwedge^i \mathcal{M} \otimes \mathcal{I}_X(j)) = H^2(\mathbb{P}^r, \bigwedge^i \mathcal{M} \otimes \mathcal{I}_\Lambda \otimes \mathcal{I}_X(j)) = 0$$

if $j \geq m$ or if $j = m - 1$ and $1 \leq i \leq p$. This fact and the above (i) $\sim$ (iii) imply (2.3). This completes the proof that $X \cap \Lambda$ is m -regular and satisfies condition $N_{m-1,p}$. $\square$

3. HIGH DEGREE HYPERELLIPTIC CURVES

Let $\mathcal{C} \subset \mathbb{P}^{d-g}$ be a smooth linearly normal curve of genus $g \geq 2$ and degree $d \geq 2g + 2$. Then it is well known that $\mathcal{C}$ is 3-regular and satisfies condition $N_{2,d-2g-1}$. Furthermore, Theorem 2 in [GL] says that it fails to satisfy condition $N_{2,d-2g}$ if and only if either

- (a) $\mathcal{C}$ admits a $(d - 2g + 2)$ -secant $(d - 2g)$ -plane or
- (b) $\mathcal{C}$ is a hyperelliptic curve.

Theorem 1.1 in [EGHP] reproves the direction that if (a) holds, then $\mathcal{C}$ fails to satisfy condition $N_{2,d-2g}$.

On the other hand, if $\mathcal{C}$ is a hyperelliptic curve, then it happens that there exists no $(d - 2g + 2)$ -secant $(d - 2g)$ -plane to $\mathcal{C}$. In this case we have the following common property.

Proposition 3.1. *Let $\mathcal{C} \subset \mathbb{P}^{d-g}$ be as above. If $\mathcal{C}$ is a hyperelliptic curve, then there exists a proper linear subspace Λ of $\mathbb{P}^r$ such that the intersection $\mathcal{C} \cap \Lambda$ fails to satisfy condition $N_{2,d-2g}$.*

Proof. Let $f : \mathcal{C} \rightarrow \mathbb{P}^1$ be the double covering and let $A = f^* \mathcal{O}_{\mathbb{P}^1}(1)$. We can write $\mathcal{L} := \mathcal{O}_{\mathcal{C}}(1)$ uniquely as $\mathcal{L} = A^u \otimes B$ where B is an effective line bundle of degree $v := d - 2u$ satisfying the condition $H^0(\mathcal{C}, B \otimes A^{-1}) = 0$. Then $v \leq g + 1$ and $\mathcal{C}$ is contained in the rational normal surface scroll $S(u + v - g - 1, u)$. Furthermore, letting C_0 and $\mathfrak{f}$ denote respectively the minimal section of $S := S(u + v - g - 1, u)$ and a fiber of the projection map $\pi : S \rightarrow \mathbb{P}^1$, our curve $\mathcal{C}$ is linearly equivalent to $2C_0 + (2g + 2 - v)\mathfrak{f}$ as a divisor of S . For details, we refer the reader to [P, Section 3].

Now, let Λ be the span of C_0 . Thus $\Lambda = \mathbb{P}^{u+v-g-1}$. Consider the intersection $\Gamma := \mathcal{C} \cap \Lambda$. Since $S \cap \Lambda$ is exactly C_0 , the length of Γ is equal to $\mathcal{C} \cdot C_0 = v$. Also since Γ is contained in the rational normal curve $C_0 \subset \mathbb{P}^{u+v-g-1}$ and since $|\Gamma| = v = 2(u + v - g - 1) + 1 - (d - 2g - 1)$, it holds by [NP, Proposition 2.3] that Γ fails to satisfy condition $N_{2,d-2g}$. $\square$

Remark 3.2. (1) The proof of Corollary 1.2 comes immediately by combining Proposition 3.1 and Theorem 1.1.

(2) In the proof of Proposition 3.1, note that A^{g-1} is the canonical sheaf of $\mathcal{C}$. Thus $\mathcal{C}$ admits a $(d - 2g + 2)$ -secant $(d - 2g)$ -plane if and only if $H^0(\mathcal{C}, \mathcal{L} \otimes A^{-g+1}) \neq 0$ if and only if $u \geq g - 1$. On the other hand, if $u < g - 1$ (and hence $d \geq 3g - 3$) then there exists no $(d - 2g + 2)$ -secant $(d - 2g)$ -plane to $\mathcal{C}$.

ACKNOWLEDGEMENT

This research was supported by the Korea Research Foundation Grant funded by the Korean Government (NRF-2013R1A1A2008445).

REFERENCES

- [E] David Eisenbud, *The geometry of syzygies. A second course in commutative algebra and algebraic geometry*, Graduate Texts in Mathematics, vol. 229, Springer-Verlag, New York, 2005. MR2103875 (2005h:13021)
- [EGHP] David Eisenbud, Mark Green, Klaus Hulek, and Sorin Popescu, *Restricting linear syzygies: algebra and geometry*, Compos. Math. **141** (2005), no. 6, 1460–1478, DOI 10.1112/S0010437X05001776. MR2188445 (2006m:14072)

- [GL] M. Green and R. Lazarsfeld, *Some results on the syzygies of finite sets and algebraic curves*, Compositio Math. **67** (1988), no. 3, 301–314. MR959214 (90d:14034)
- [L] Robert Lazarsfeld, *Positivity in algebraic geometry. I*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 48, Springer-Verlag, Berlin, 2004. Classical setting: line bundles and linear series. MR2095471 (2005k:14001a)
- [NP] Uwe Nagel and Yves Pitteloud, *On graded Betti numbers and geometrical properties of projective varieties*, Manuscripta Math. **84** (1994), no. 3-4, 291–314, DOI 10.1007/BF02567458. MR1291122 (95i:13017)
- [P] Euisung Park, *Higher syzygies of hyperelliptic curves*, J. Pure Appl. Algebra **214** (2010), no. 2, 101–111, DOI 10.1016/j.jpaa.2009.04.006. MR2559684 (2010i:14057)

DEPARTMENT OF MATHEMATICS, KOREA UNIVERSITY, SEOUL 136-701, REPUBLIC OF KOREA
E-mail address: euisungpark@korea.ac.kr