

ON THE LENGTH OF FINITE GROUPS AND OF FIXED POINTS

E. I. KHUKHRO AND P. SHUMYATSKY

(Communicated by Pham Huu Tiep)

ABSTRACT. The generalized Fitting height of a finite group G is the least number $h = h^*(G)$ such that $F_h^*(G) = G$, where the $F_i^*(G)$ is the generalized Fitting series: $F_1^*(G) = F^*(G)$ and $F_{i+1}^*(G)$ is the inverse image of $F^*(G/F_i^*(G))$. It is proved that if G admits a soluble group of automorphisms A of coprime order, then $h^*(G)$ is bounded in terms of $h^*(C_G(A))$, where $C_G(A)$ is the fixed-point subgroup, and the number of prime factors of $|A|$ counting multiplicities. The result follows from the special case when $A = \langle \varphi \rangle$ is of prime order, where it is proved that $F^*(C_G(\varphi)) \leq F_9^*(G)$.

The nonsoluble length $\lambda(G)$ of a finite group G is defined as the minimum number of nonsoluble factors in a normal series each of whose factors is either soluble or is a direct product of nonabelian simple groups. It is proved that if A is a group of automorphisms of G of coprime order, then $\lambda(G)$ is bounded in terms of $\lambda(C_G(A))$ and the number of prime factors of $|A|$ counting multiplicities.

1. INTRODUCTION

The structure of an arbitrary finite group G can be described in terms of length parameters related to the normal series with ‘nice’ sections. One way of constructing such a series is based on the generalized Fitting subgroup $F^*(G)$. Recall that $F^*(G)$ is the product of the Fitting subgroup $F(G)$ and all subnormal quasisimple subgroups; here a group is quasisimple if it is perfect and its quotient by the centre is a nonabelian simple group. Then the *generalized Fitting series* of G is defined starting from $F_1^*(G) = F^*(G)$, and then by induction, with $F_{i+1}^*(G)$ being the inverse image of $F^*(G/F_i^*(G))$. The least number h such that $F_h^*(G) = G$ is naturally defined as the *generalized Fitting height* $h^*(G)$ of G . Clearly, if G is soluble, then $h^*(G) = h(G)$ is the ordinary Fitting height of G . Bounding the generalized Fitting height of a finite group G greatly facilitates using the classification of finite simple groups (and is itself often obtained using the classification). One such example is the reduction of the Restricted Burnside Problem to soluble and nilpotent groups in the Hall–Higman paper [1], where the generalized Fitting height was in effect bounded for groups of a given exponent (using the classification as a conjecture at the time). A similar example is Wilson’s reduction of the problem of local finiteness of periodic profinite groups to pro- p groups in [11].

Received by the editors May 2, 2014 and, in revised form, May 28, 2014.

1991 *Mathematics Subject Classification.* Primary 20D45.

Key words and phrases. Finite groups, nonsoluble length, generalized Fitting height, coprime automorphism group.

This work was supported by CNPq-Brazil. The first author thanks CNPq-Brazil and the University of Brasilia for support and hospitality that he enjoyed during his visits to Brasilia.

Another useful, more rough, length parameter is the *nonsoluble length* $\lambda(G)$ of a finite group G , which is defined as the minimum number of nonsoluble factors in a normal series, each of whose factors either is soluble or is a direct product of nonabelian simple groups. More precisely, consider normal series

$$1 = G_0 \leq G_1 \leq \cdots \leq G_{2h+1} = G$$

in which for i even the factor G_{i+1}/G_i is soluble (possibly trivial), and for i odd the factor G_{i+1}/G_i is a (nonempty) direct product of nonabelian simple groups. Then the nonsoluble length $\lambda(G)$ is equal to the minimum possible value of h . (In particular, the group is soluble if and only if its nonsoluble length is 0.) Bounding the nonsoluble length was one of the steps in Wilson's paper [11]; more recently we used this parameter in the study of both finite and profinite groups in [4, 5].

In the present paper we consider the generalized Fitting height and nonsoluble length of a finite group G relative to the same parameters of the fixed-point subgroup $C_G(A)$ of a soluble group of automorphisms A of coprime order. The results are modelled on Thompson's paper [8], where in the case of soluble groups G , A of coprime orders, a bound for the Fitting height of G was obtained in terms of the Fitting height of the fixed-point subgroup $C_G(A)$ and the number of prime factors of $|A|$ counting multiplicities.

Theorem 1.1. *Suppose that a finite group G admits a soluble group of automorphisms A of coprime order. Then its generalized Fitting height $h^*(G)$ is bounded in terms of the generalized Fitting height $h^*(C_G(A))$ of the fixed-point subgroup $C_G(A)$ and the number of prime factors of $|A|$ counting multiplicities.*

In the proof we use the underlying result of Thompson [8] for soluble groups in the special case when $A = \langle \varphi \rangle$ is of prime order; then $F(C_G(\varphi)) \leq F_4(G)$. In fact, we also obtain an analogous result for the generalized Fitting height.

Theorem 1.2. *Suppose that a finite group G admits an automorphism φ of prime order coprime to $|G|$. Then $F^*(C_G(\varphi)) \leq F_9^*(G)$.*

The proof of Theorem 1.2 uses the following consequences of the classification: the validity of Schreier's conjecture on solubility of outer automorphism groups of non-abelian finite simple groups and the result of Wang and Chen [10] that in a finite nonabelian simple group the fixed-point subgroup of an automorphism of coprime order cannot be a nilpotent group. Theorem 1.1 follows from Theorem 1.2 by a straightforward induction, which furnishes the bound $h^*(G) \leq 9^{\alpha(|A|)} \cdot h^*(C_G(A)) + (9^{\alpha(|A|)} - 1)/8$, where $|A|$ is the product of $\alpha(|A|)$ primes. We have no reason to believe that this bound is anywhere near a sharp one. It is worth mentioning that in the soluble case the original bound of Thompson, $h(G) \leq 5^{\alpha(|A|)} \cdot h(C_G(A))$ (and $h(G) \leq 5^{\alpha(|A|)}$ if $h(C_G(A)) = 0$ when $C_G(A) = 1$), was later drastically improved by various authors, with the first linear bound obtained by Kurzweil [6], and the definitive sharp bound by Turull [9]. We hope that similar improvements can be made for the bound in Theorem 1.1.

We now state a similar result for the nonsoluble length.

Theorem 1.3. *Suppose that a finite group G admits a group of automorphisms A of coprime order. Then its nonsoluble length $\lambda(G)$ is bounded in terms of the nonsoluble length $\lambda(C_G(A))$ of the fixed-point subgroup $C_G(A)$ and the number of prime factors of $|A|$ counting multiplicities.*

The proof of Theorem 1.3 follows from the main case, where $A = \langle \varphi \rangle$ is of prime order, and $G = [G, \varphi]$, where we prove that $\lambda(G) \leq \lambda(C_G(\varphi)) + 1$. Straightforward induction then gives the bound $\lambda(G) \leq 2^{\alpha(|A|)}(\lambda(C_G(A)) + 1) - 1$ in Theorem 1.3. The proof uses the classification as much as the validity of Schreier's conjecture on solubility of outer automorphism groups of nonabelian finite simple groups.

The coprimeness conditions are unavoidable in all our theorems. For Theorems 1.1 and 1.2 this was shown even for soluble groups by examples in Thompson's paper [8]. For Theorem 1.3 one can take a repeated wreath product $G = S \wr (S \wr (\dots (S \wr S) \dots))$ of a simple group S containing an element a of prime order with soluble centralizer $C_S(a)$. Let $b \in S$ be a conjugate of a different from a . Then the element (b, a, a, \dots, a) in the base of the last wreath product has a soluble centralizer in G , while the nonsoluble length of G is unbounded.

Throughout the paper we use without special references the well-known property of coprime actions: if a group A acts by automorphisms on a finite group G of coprime order, $(|A|, |G|) = 1$, then $C_{G/N}(A) = C_G(A)N/N$ for any A -invariant normal subgroup N . We usually use the same letter to denote induced automorphisms of invariant sections.

2. GENERALIZED FITTING HEIGHT

In this section we obtain bounds for the generalized Fitting height in terms of the generalized Fitting height of the fixed-point subgroup of a group of automorphisms. Recall that the generalized Fitting subgroup $F^*(G)$ of a finite group G is the product of the Fitting subgroup $F(G)$ and the characteristic subgroup $E(G)$, which is a central product of all subnormal quasisimple subgroups of G , that is, $E(G) = \prod Q_i$ over all Q_i such that Q_i is subnormal in G , $Z(Q_i) \leq [Q_i, Q_i]$, and $S_i = Q_i/Z(Q_i)$ is a nonabelian simple group. Then $[F(G), E] = 1$ and $E(G)/Z(E(G)) \cong F^*(G)/F(G)$ is the direct product of the S_i . Acting by conjugation, the group G permutes the factors Q_i and $C_G(F^*(G)) \leq F(G)$. The following fact (see, for example, [4, Lemma 2.1]) is a well-known consequence of Schreier's conjecture on solubility of outer automorphism groups of nonabelian finite simple groups confirmed by the classification.

Lemma 2.1. *Let $L/S(G) = F^*(G/S(G))$ be the generalized Fitting subgroup of the quotient by the soluble radical $S(G)$ of a finite group G , and let K be the kernel of the permutational action of G on the set of subnormal simple factors of $L/S(G)$. Then K/L is soluble. \square*

The following elementary lemma may also be known.

Lemma 2.2. *Let N be a soluble normal subgroup of a finite group G . If $x \in F^*(G)$, then $x \in F(N\langle x \rangle)$.*

Proof. We claim that $[N\langle x \rangle, x, \dots, x] = 1$ if x is repeated sufficiently many times. Write $x = yz$, where $y \in F(G)$ and $z \in E(G)$. Since $[N\langle x \rangle, x] \leq N \cap F^*(G) \leq F(G)$ and $[F(G), z] = 1$, we have

$$\begin{aligned} [N, x, x, \dots, x] &\leq [F(G), yz, \dots, yz] \\ &= [F(G), y, \dots, y]. \end{aligned}$$

The commutator on the right becomes trivial if it is sufficiently long, because $F(G)$ is nilpotent. As a result, $x \in F(N\langle x \rangle)$ by Baer's theorem (see [2, Satz III.6.15]). \square

We shall use without special references the following well-known properties of the generalized Fitting subgroups relative to normal subgroups: if $N \trianglelefteq G$, then $F^*(N) = F^*(G) \cap N$ and $F^*(G)N/N \leq F^*(G/N)$. These and other properties follow, for example, from the fact that $F^*(G)$ is the set of all elements of G that induce inner automorphisms on every chief factor of G ; see, for example, [3, Ch. X, §13]. It is easy to see that similar properties hold for the higher terms of the generalized Fitting series: if $N \trianglelefteq G$, then $F_i^*(N) = F_i^*(G) \cap N$ and $F_i^*(G)N/N \leq F_i^*(G/N)$.

Proof of Theorem 1.2. We have a finite group G admitting an automorphism φ of prime order p coprime to $|G|$. We need to show that $F^*(C_G(\varphi)) \leq F_9^*(G)$. Recall that $S(G)$ denotes the soluble radical of G . Let L be the inverse image of $F^*(G/S(G))$ and let $L/S(G) = S_1 \times \cdots \times S_m$, where the S_i are nonabelian simple groups. Let K denote the kernel of the permutational action of G on $\{S_1, \dots, S_m\}$. The quotient K/L is soluble by Lemma 2.1.

Lemma 2.3. *We have the inclusion $F^*(C_G(\varphi)) \leq K$.*

Proof. We argue by contradiction. Let $x \in F^*(C_G(\varphi)) \setminus K$. Consider the quotient $\bar{G} = G/S(G)$. Since $C_{\bar{G}}(\varphi)$ is the image of $C_G(\varphi)$, then also $\bar{x} \in F^*(C_{\bar{G}}(\varphi)) \setminus \bar{K}$. Thus we can assume that $S(G) = 1$ and $L = S_1 \times \cdots \times S_m$. Then x must permute these factors nontrivially. Since $x \in C_G(\varphi)$, the element x also permutes the orbits of φ in the permutational action on $\{S_1, \dots, S_m\}$. If x really moves some nontrivial φ -orbit, say, $\{S_{i_1}, \dots, S_{i_p}\}$, then x moves the ‘diagonal’ nonabelian simple group $D = C_G(\varphi) \cap (S_{i_1} \times \cdots \times S_{i_p})$. Since $S_{i_1} \times \cdots \times S_{i_p}$ is a subnormal subgroup, D is subnormal in $C_G(\varphi)$ and therefore is normal in $F^*(C_G(\varphi))$, contrary to being moved by x .

Thus, x stabilizes every nontrivial φ -orbit $\{S_{i_1}, \dots, S_{i_p}\}$. Since the centralizer of φ as a cycle of length p in the symmetric group on p symbols is $\langle \varphi \rangle$ and p is coprime to $|G|$, it follows that x must leave each factor S_{i_1}, \dots, S_{i_p} fixed.

Therefore x must really move some of the one-element orbits of φ , say, $S_i^x = S_j \neq S_i$, where both S_i and S_j are φ -invariant. By the result of Wang and Chen [10] based on the classification, $C_{S_i}(\varphi)$ cannot be a nilpotent group. If $x \in F(C_G(\varphi))$, then we obtain a contradiction, since $[C_{S_i}(\varphi), x] \leq F(C_G(\varphi))$, and yet the projection of $[C_{S_i}(\varphi), x]$ onto S_i covers $C_{S_i}(\varphi)$, which is not nilpotent. It remains to consider the case where $x \in F^*(C_G(\varphi)) \setminus F(C_G(\varphi))$. Then at least one of the quasisimple components Q of $F^*(C_G(\varphi))$ is not contained in K and moves S_i . Let M be a minimal normal subgroup of $C_{S_i}(\varphi)$. Since $C_{S_i}(\varphi)$ is subnormal in $C_G(\varphi)$, it follows that $M \leq F^*(C_G(\varphi))$. If M is nilpotent, then $[M, Q] = 1$. If M is nonsoluble, then it is a product of quasisimple components of $F^*(C_G(\varphi))$ which are all different from $Q \not\leq L$, and then, too, $[M, Q] = 1$. In either case we obtained a contradiction with $[M, Q]$ having nontrivial projection onto S_i . □

We now complete the proof of Theorem 1.2. Recall that $L/S(G) = F^*(G/S(G))$ and K is the kernel of the permutational action of G on the set of subnormal simple factors of $L/S(G)$. By Lemma 2.3 we know that $F^*(C_G(\varphi))$ is contained in K . Therefore $F^*(C_G(\varphi)) \leq F^*(C_K(\varphi))$. Since $F_9^*(K) \leq F_9^*(G)$, it is sufficient to prove that $F^*(C_K(\varphi)) \leq F_9^*(K)$. Thus, we can assume from the outset that $G = K$.

Let $x \in F^*(C_G(\varphi))$ and let $X = \langle x^G \rangle$ be the normal closure of x . It is sufficient to prove that $X \leq F_9^*(G)$. Consider the soluble subgroup $H = S(X)\langle x \rangle$.

By Lemma 2.2 applied to $C_G(\varphi)$ with $N = C_{S(X)}(\varphi)$, we obtain that

$$x \in F(C_{S(X)}(\varphi)\langle x \rangle) = F(C_H(\varphi)).$$

Then $x \in F_4(H)$ by Thompson’s theorem [8].

Therefore, $[x, S(X)] \leq F_4(H) \cap S(X) \leq F_4(S(X)) \leq F_4(G)$. In the quotient $\bar{G} = G/F_4(G)$ this means that $\bar{x} \in C_{\bar{G}}(S(\bar{X}))$. Since $X = \langle x^G \rangle$, it follows that $S(\bar{X}) \leq Z(\bar{X})$. Then $F^*(\bar{X}) = S(\bar{X})(\bar{L} \cap \bar{X})$ because $L/S(G)$ is a direct product of simple groups and G/L is soluble. Therefore $\bar{X}/F^*(\bar{X})$ is a soluble group. Since $F^*(\bar{X}) \leq F^*(\bar{G})$, we obtain that the image \tilde{X} of X in $\tilde{G} = G/F_5^*(G)$ is a soluble group.

We now apply Thompson’s theorem [8] to \tilde{X} . Namely, since $\tilde{x} \in F^*(C_{\tilde{X}}(\varphi)) = F(C_{\tilde{X}}(\varphi))$, we have $\tilde{x} \in F_4(\tilde{X})$. Then $\tilde{X} = F_4(\tilde{X}) \leq F_4(\tilde{G})$. As a result, $X \leq F_9^*(G)$, as required. □

Proof of Theorem 1.1. We have a finite group G admitting a soluble group of automorphisms A of coprime order. We are going to prove that the generalized Fitting height $h^*(G)$ satisfies the inequality $h^*(G) \leq 9^\alpha \cdot h^*(C_G(A)) + (9^\alpha - 1)/8$, where $|A|$ is a product of $\alpha = \alpha(|A|)$ primes (counting multiplicities). We proceed by induction on $\alpha(|A|)$. Let $h = h^*(C_G(A))$. When $\alpha(|A|) = 1$, it follows from Theorem 1.2 that $C_G(A) \leq F_{9h}^*(G)$, and then $G/F_{9h}^*(G)$ is nilpotent by Thompson’s theorem [7]. Thus, $h^*(G) \leq 9h + 1$. For $\alpha(|A|) > 1$, let A_0 be a normal subgroup of prime index in A . Then $C_G(A_0)$ admits the group of automorphisms A/A_0 of prime order and $C_{C_G(A_0)}(A/A_0) = C_G(A)$. By the above, $h^*(C_G(A_0)) \leq 9h + 1$. By induction applied to A_0 we obtain

$$\begin{aligned} h^*(G) &\leq 9^{\alpha-1} \cdot (9h + 1) + (9^{\alpha-1} - 1)/8 \\ &= 9^\alpha \cdot h + (9^\alpha - 1)/8, \end{aligned}$$

as required. □

3. NONSOLUBLE LENGTH

Recall that the nonsoluble length $\lambda(G)$ of a finite group G is the minimum number of nonsoluble factors in a normal series each of whose factors is either soluble or is a direct product of nonabelian simple groups. Consider the ‘upper nonsoluble series’ of G , which by definition starts from the soluble radical $M_1 = S(G)$, then L_1 is the full inverse image of $F^*(G/M_1)$, and then by induction M_k is the full inverse image of the soluble radical of G/L_{k-1} and L_k the full inverse image of $F^*(G/M_k)$. It is easy to show that $\lambda(G) = m$, where m is the first positive integer such that $M_{m+1} = G$. In the normal series

$$(1) \quad 1 = L_0 \leq M_1 < L_1 \leq M_2 < \dots \leq M_{m+1} = G$$

each quotient $U_i = L_i/M_i$ is a (nontrivial) direct product of nonabelian simple groups, and each quotient M_i/L_{i-1} is soluble (possibly trivial). Since $L_i/M_i = F^*(G/M_i)$, the properties mentioned at the beginning of §2 apply. In particular, if we write one of those nonsoluble quotients as a direct product $U_k = S_1 \times \dots \times S_t$ of nonabelian simple groups S_i , then the set of these factors S_i is unique, characterized as the set of subnormal simple subgroups of G/M_k . The group G and its automorphisms permute these subnormal factors, and for brevity we simply speak of their orbits on U_k meaning orbits in this permutational action. The subgroup L_k/M_k contains its centralizer in G/M_k . The kernel of the permutational action

of G/L_k on $\{S_1, \dots, S_t\}$ is soluble by Lemma 2.1. Therefore the inverse image of this kernel is contained in M_{k+1} . We shall routinely use this fact without special references.

We shall need the following technical lemma.

Lemma 3.1. *Let N be a normal subgroup of a finite group G and K/N a simple subnormal subgroup of G/N . Let D be a subgroup of G such that $K = DN$. Suppose that $D \leq L_j(G)N$ and $D \not\leq M_j(G)N$. Then $[D, D^x]NM_j(G) = KM_j(G)$ for any $x \in L_j(G)N$.*

Proof. We can pass to the quotient $G/M_j(G)$ and without loss of generality assume that $j = 1$ and G has no nontrivial normal soluble subgroups. It follows that $K = SN$, where S is a subnormal simple subgroup of G (contained in $L_1(G)$). The group N permutes the subnormal simple factors of L_1 contained in K and normalizes $L_1 \cap N$; therefore N also normalizes S . Hence, $K = S \times N$. Now it is clear that the projection of D onto S is equal to S and hence so is that of D^x for any $x \in L_1(G)N$. Hence the projection of $[D, D^x]$ also equals S , as S is nonabelian simple. □

Proof of Theorem 1.3. We have a finite group G admitting a group of automorphisms A of coprime order. We wish to show that its nonsoluble length $\lambda(G)$ is bounded in terms of $\lambda(C_G(A))$ and the number of prime factors of $|A|$ counting multiplicities. We can of course assume that G is nonsoluble, whence A then has odd order and therefore is soluble by the Feit–Thompson theorem. The result will follow from the case of $|A|$ being a prime by straightforward induction on $|A|$. Thus the bulk of the proof is the following proposition.

Proposition 3.2. *Suppose that a finite group G admits an automorphism φ of prime order p coprime to $|G|$ such that $[G, \varphi] = G$. Then $\lambda(G) \leq \lambda(C(\varphi)) + 1$.*

Clearly, we can assume that G is nonsoluble; in particular, $|\varphi| \geq 3$.

We consider the ‘upper nonsoluble series’ for $C_G(\varphi)$ constructed in the same way as (1) was constructed for G , with its terms denoted by

$$(2) \quad 1 = \lambda_0 \leq \mu_1 < \lambda_1 \leq \mu_2 < \dots \leq \mu_{l+1} = C_G(\varphi),$$

so that the quotients λ_i/μ_i are (nontrivial) direct products of nonabelian simple groups, and the quotients μ_i/λ_{i-1} (possibly trivial) are soluble. Our task is to show that the nonsoluble length $\lambda(G)$ of G , which is equal to m in (1), is at most $\lambda(C_G(\varphi)) + 1$, that is, at most $l + 1$.

Lemma 3.3. *For every $k \leq m - 1$, the automorphism φ has nontrivial orbits (of length $p = |\varphi|$) on the quotient U_k .*

Proof. For $k \leq m - 1$, the group G/L_k is nonsoluble and therefore by Lemma 2.1 acts nontrivially by permutations on the set of subnormal simple factors of U_k . Since $G = [G, \varphi]$ by hypothesis, the automorphism φ also has nontrivial orbits (of course, of length $p = |\varphi|$) on the set of these factors. Indeed, if φ belonged to the kernel of the permutational action of $G\langle\varphi\rangle$, then $[G, \varphi]$ would also belong to this kernel, which is a normal subgroup. □

Consider one of the nonsoluble quotients $U_k = L_k/M_k$ such that the automorphism φ has a nontrivial orbit $\{S_1, \dots, S_p\}$ of length $p = |\varphi|$ on the set of subnormal factors of U_k . Let $S = S_1 \times \dots \times S_p$, and let $D = C_S(\varphi)$ be the diagonal subgroup of S , which is a nonabelian simple group (isomorphic to S_1).

Lemma 3.4. *There is an index i such that for any φ -invariant subgroup F with $FM_k/M_k \geq D$ there is a subgroup $\hat{D} \leq C_F(\varphi)$ such that*

- (1) $\hat{D}M_k/M_k = D,$
- (2) $\hat{D} \leq \lambda_i M_k,$
- (3) $\hat{D} \not\leq \mu_i M_k.$

Proof. Recall that since the action of φ on G is coprime, the fixed points of φ in any φ -invariant section are images of fixed points of φ in G . Let \tilde{D} be any subgroup of $C_F(\varphi)$ such that $\tilde{D}M_k/M_k = D$.

Consider the quotient $\tilde{G} = G/M_k$. The images $\bar{\lambda}_i$ and $\bar{\mu}_i$ of the terms of the series (2) form a normal series of $C_G(\varphi)M_k/M_k$ with $\bar{\lambda}_i/\bar{\mu}_i$ being direct products (possibly empty) of nonabelian simple groups, and $\bar{\mu}_i/\bar{\lambda}_{i-1}$ being soluble. Since $D = \tilde{D}M_k/M_k$ is a nonabelian simple subgroup of $C_G(\varphi)M_k/M_k$, there is an index i such that $D \leq \bar{\lambda}_i$ and $D \not\leq \bar{\mu}_i$. This means that $\tilde{D} \leq \lambda_i M_k$ and $\tilde{D} \not\leq \mu_i M_k$. \square

We fix the notation \hat{D} for a subgroup of $C_F(\varphi)$ constructed for a diagonal subgroup D of a nontrivial φ -orbit in $U_k = L_k/M_k$ in accordance with Lemma 3.4 (for some F); we choose \hat{D} to be minimal by inclusion subject to satisfying Lemma 3.4. We then say for brevity that \hat{D} corresponds to D (with or without specifying for which F); these subgroups will also be denoted by $D_k = D$ and $\hat{D}_k = \hat{D}$, where the index only indicates the level of the quotient $U_k = L_k/M_k$. The subgroups \hat{D}_k can be chosen in a special way described in the following lemma. Note that since $\hat{D}_k \leq C_G(\varphi)$, the subgroup \hat{D}_k permutes the orbits of φ in its permutational action on the set of simple subnormal factors of U_{k-1} .

Lemma 3.5. *Suppose that $k \geq 2$ and D_k is a diagonal subgroup in a nontrivial φ -orbit on U_k . Then there exists a subgroup \hat{D}_k corresponding to D_k that does not stabilize some nontrivial φ -orbits on U_{k-1} .*

Proof. Note that the automorphism φ does have nontrivial orbits on U_{k-1} by Lemma 3.3.

Recall that $D_k = C_S(\varphi)$ is the ‘diagonal’ in some $S = S_1 \times \dots \times S_p$, where $S_i^\varphi = S_{i+1 \pmod p}$. Choose a minimal φ -invariant subgroup \hat{S} such that $\hat{S}M_k/M_k = S$. (Subgroups with these properties obviously do exist: say, the inverse image of S .) Let $\hat{D}_k \leq C_{\hat{S}}(\varphi)$ be the subgroup corresponding to D_k for $F = \hat{S}$ in the sense of Lemma 3.4. We claim that \hat{D}_k is a required subgroup. We argue by contradiction: suppose that \hat{D}_k stabilizes all nontrivial φ -orbits on U_{k-1} . Let $\{R_1, \dots, R_p\}$ be a nontrivial φ -orbit on U_{k-1} . Since the centralizer of φ as a cycle of length p in the symmetric group on p symbols is $\langle \varphi \rangle$ and p is coprime to $|G|$, it follows that \hat{D}_k must leave each factor R_1, \dots, R_p fixed.

On the other hand, since the kernel of the permutational action on U_{k-1} is contained in M_k by Lemma 2.1, there is an element $d \in \hat{D}_k \setminus M_k$ that moves some of the subnormal simple factors of U_{k-1} . Let R_0 be such a factor that is fixed by φ but moved by d .

In the permutational action of $\hat{S}\langle \varphi \rangle$ on the set of subnormal factors of U_{k-1} , we now focus on the action of $\hat{S}\langle \varphi \rangle$ on its orbit containing R_0 . Since S is the only proper normal subgroup of $S\langle \varphi \rangle$ and $d \notin M_k$, the kernel of this transitive action is contained in $\hat{S} \cap M_k$; in particular, φ acts nontrivially. Let H be the stabilizer

of R_0 in $\hat{S}\langle\varphi\rangle$. We have $\varphi \in H$, but H does not contain \hat{S} , as it does not contain $d \in \hat{D}_k$. Since φ acts nontrivially, there are elements $x \in \hat{S}$ such that $\varphi \notin H^x$.

We claim that

$$(3) \quad S \leq HM_k/M_k.$$

Indeed, by our supposition, whenever $\varphi \notin H^y$ for $y \in \hat{S}$, we must have $\hat{D}_k \leq H^y$. In other words, $[\varphi, x] \notin H$ implies $\hat{D}_k^x \leq H$. First assume that there is no x with this property whose image modulo M_k is a nontrivial element of S_1 . Then $[S_1, \varphi] \leq \bar{H}$, where $\bar{H} = HM_k/M_k$. We obtain that modulo M_k , in the coordinates of $S_1 \times \dots \times S_p$, all elements $(a, a^{-1}, 1, \dots)$ are in \bar{H} . (Henceforth in this paragraph the dots denote 1s.) Multiplying $(a, a^{-1}, 1, \dots)$ by $(b, b^{-1}, 1, \dots)$ and then by $((ab)^{-1}, (ab), 1, \dots)$ for some noncommuting elements $a, b \in S_1$, we obtain an element $(1, g, 1, \dots) \in \bar{H}$ with $g \neq 1$. Then conjugating by $(1, a, a^{-1}, \dots) \in \bar{H}$ we obtain $(1, S_2, 1, \dots) \leq \bar{H}$. Here the element $(1, a, a^{-1}, \dots)$ belongs to \bar{H} because $(1, a, a^{-1}, \dots) = (a, a^{-1}, 1, \dots)^\varphi \in H^\varphi = H$, as $\varphi \in H$; also recall that $|\varphi| > 2$. From $(1, S_2, 1, \dots) \leq \bar{H}$ by the action of φ we obtain $S \leq \bar{H}$.

Thus we can assume that there is $x \in \hat{S}$ whose image modulo M_k is a nontrivial element of S_1 such that $[\varphi, x] \notin H$, and therefore, $\hat{D}_k^x \leq H$. This means that modulo M_k , in the coordinates of $S_1 \times \dots \times S_p$, all elements (b^x, b, b, \dots) lie in \bar{H} and there is b with $b^x \neq b$. Using the action of φ we obtain $(b, b^x, b, \dots) \in \bar{H}$, whence $(c, c^{-1}, 1, \dots) \in \bar{H}$ for $c = [b, x] \neq 1$. Conjugating by (b, b, b^x, \dots) we obtain $(c^b, (c^{-1})^b, 1, \dots) \in \bar{H}$ for any b . We can choose $b_1, b_2 \in S_1$ such that $[c^{b_1}, c^{b_2}] \neq 1$ and therefore also $[(c^{-1})^{b_1}, c^{b_2}] \neq 1$. We conjugate $(c^{b_1}, (c^{-1})^{b_1}, 1, \dots)$ by $(1, c^{b_2}, \dots)$ and obtain $(c^{b_1}, ((c^{-1})^{b_1})^{c^{b_2}}, 1, \dots) \in \bar{H}$. Multiplying by the inverse of $(c^{b_1}, (c^{-1})^{b_1}, 1, \dots)$ we obtain $(1, g, 1, 1, \dots) \in \bar{H}$ with $g \neq 1$. Conjugating by (b^x, b, b, \dots) for all b we again obtain $S_2 \leq \bar{H}$, and then $S \leq \bar{H}$, completing the proof of (3).

Now consider $H \cap \hat{S}$, which is a φ -invariant subgroup. By (3) above, $\hat{S} \leq HM_k$; we claim that also $\hat{S} \leq (H \cap \hat{S})M_k$. For any $s \in \hat{S}$ we have $s = ha$ for $h \in H$ and $a \in M_k$. But $H \leq \hat{S}\langle\varphi\rangle$, so $h = s_1\varphi^i$ for some i and $s_1 \in \hat{S}$, and then $s = s_1\varphi^i a$. Clearly, $\varphi^i = 1$, so $s = s_1 a$, where $s_1 = h \in H \cap \hat{S}$. Finally, $H \cap \hat{S}$ is a proper subgroup of \hat{S} , since $d \notin H$. This contradicts the choice of \hat{S} as a minimal φ -invariant subgroup such that $\hat{S}M_k/M_k = S$. □

Proof of Proposition 3.2. The idea is to construct a chain of subgroups $\hat{D}_{m-1}, \dots, \hat{D}_1$, working from the second highest quotient in (1) downwards, in such a way that each of them ‘marks’ a strictly higher quotient among the λ_i/μ_i in the series (2) of $C_G(\varphi)$ than the next one with a smaller index.

The proposition is obviously correct if $m \leq 1$. Therefore we assume that $m \geq 2$. By Lemma 3.3 the automorphism φ has a nontrivial orbit on the set of subnormal factors of L_{m-1}/M_{m-1} . If $m \geq 3$, let $\hat{D}_{m-1} \leq C_G(\varphi)$ be a subgroup of level $m - 1$ constructed by Lemma 3.5 in such a way that \hat{D}_{m-1} does not stabilize some nontrivial φ -orbit on L_{m-2}/M_{m-2} . Let D_{m-2} be the diagonal in one of these orbits which is not stabilized by \hat{D}_{m-1} . Then let $\hat{D}_{m-2} \leq C_G(\varphi)$ be a subgroup of level $m - 2$ corresponding to D_{m-2} by Lemma 3.5 that does not stabilize some nontrivial φ -orbit on L_{m-3}/M_{m-3} . Continuing in this manner, we obtain a sequence of subgroups $\hat{D}_{m-1}, \dots, \hat{D}_1$ such that for every k the subgroup \hat{D}_k is constructed by Lemma 3.5 from the diagonal D_k of a nontrivial φ -orbit on the

subnormal factors of L_k/M_k , which is not stabilized by \hat{D}_{k+1} . Therefore for every $k \geq 2$ there exists $x \in \hat{D}_k$ such that D_{k-1}^x is the ‘diagonal’ in a different orbit from the orbit where D_{k-1} is the ‘diagonal’. Then of course,

$$(4) \quad [\hat{D}_{k-1}, \hat{D}_{k-1}^x] \leq M_{k-1}.$$

We claim that \hat{D}_k is not contained in $\mu_k M_k$ for $k = 1, \dots, m - 1$. Clearly, $\hat{D}_1 \not\leq \mu_1 M_1$, since \hat{D}_1 is not soluble. By induction suppose that $\hat{D}_{k-1} \not\leq \mu_{k-1} M_{k-1}$. Let j be the index given by Lemma 3.4 such that $\hat{D}_{k-1} \leq \lambda_j M_{k-1}$ and $\hat{D}_{k-1} \not\leq \mu_j M_{k-1}$. Clearly, $k - 1 \leq j$. Let $N = C_G(\varphi) \cap M_{k-1}$; then also

$$(5) \quad \hat{D}_{k-1} \not\leq \mu_j N.$$

To perform the induction step we argue by contradiction and suppose that $\hat{D}_k \leq \mu_k M_k$. Since μ_k/λ_{k-1} is soluble and $\hat{D}_k M_k/M_k = D_k$ is nonabelian simple, we must have $\hat{D}_k \leq \lambda_{k-1} M_k$. Since $k - 1 \leq j$, it follows that $\hat{D}_k \leq \lambda_j M_k$ and therefore $\hat{D}_k \leq \lambda_j(M_k \cap C_G(\varphi))$.

The group $C_G(\varphi)$ acts on the set of simple factors of the image of λ_j/μ_j in $C_G(\varphi)/N$. By the above, the image D_{k-1} of \hat{D}_{k-1} is one of these factors. Clearly, λ_j is in the kernel of this action by Lemma 3.1. It is easy to see that the image of $L_{k-1} \cap C_G(\varphi)$ stabilizes D_{k-1} and therefore as a normal subgroup is in the kernel of the action on the orbit containing D_{k-1} . Since M_k/L_{k-1} is soluble, we obtain that the image of $\lambda_j(M_k \cap C_G(\varphi))$ in this action on the orbit containing D_{k-1} is a soluble group. But $\hat{D}_k \leq \lambda_j(M_k \cap C_G(\varphi))$, so the image of \hat{D}_k is soluble. By minimality this image is actually trivial. Then (4) implies $\hat{D}_{k-1} \leq \mu_j N$, in contradiction with (5).

Thus, indeed \hat{D}_k is not contained in $\mu_k M_k$ for $k = 1, \dots, m - 1$. In particular, \hat{D}_{m-1} is not contained in $\mu_{m-1} M_{m-1}$, and therefore $\lambda_{m-1} \neq \mu_{m-1}$, whence $\lambda(C_G(\varphi)) \geq m - 1$. □

We now complete the proof of Theorem 1.3. Actually we prove by induction on $\alpha = \alpha(|A|)$ that $\lambda(G) \leq 2^\alpha(\lambda(C_G(A)) + 1) - 1$. Let $\lambda = \lambda(C_G(A))$ for brevity. When $\alpha(|A|) = 1$, that is, $A = \langle \varphi \rangle$ is of prime order, then $\lambda([G, \varphi]) \leq \lambda + 1$ by Proposition 3.2. Then $\lambda(G) \leq 2\lambda + 1$, since $G = [G, \varphi]C_G(\varphi)$ and $[G, \varphi]$ is a normal subgroup. For $\alpha(|A|) > 1$, let A_0 be a normal subgroup of prime index in A . Then $C_G(A_0)$ admits the group of automorphisms A/A_0 of prime order and $C_{C_G(A_0)}(A/A_0) = C_G(A)$. By the above, $\lambda(C_G(A_0)) \leq 2\lambda + 1$. It remains to apply the induction hypothesis to A_0 :

$$\begin{aligned} \lambda(G) &\leq 2^{\alpha-1}(\lambda(C_G(A_0)) + 1) - 1 \leq 2^{\alpha-1}(2\lambda + 1 + 1) - 1 \\ &= 2^{\alpha-1} \cdot 2\lambda + 2^{\alpha-1} \cdot 2 - 1 = 2^\alpha(\lambda + 1) - 1. \end{aligned} \quad \square$$

ACKNOWLEDGEMENT

The authors thank the referee for a careful reading and helpful comments.

REFERENCES

[1] P. Hall and Graham Higman, *On the p-length of p-soluble groups and reduction theorems for Burnside’s problem*, Proc. London Math. Soc. (3) **6** (1956), 1–42. MR0072872 (17,344b)
 [2] B. Huppert, *Endliche Gruppen. I* (German), Die Grundlehren der Mathematischen Wissenschaften, Band 134, Springer-Verlag, Berlin-New York, 1967. MR0224703 (37 #302)

- [3] Bertram Huppert and Norman Blackburn, *Finite groups. III*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 243, Springer-Verlag, Berlin-New York, 1982. MR662826 (84i:20001b)
- [4] E. I. Khukhro and P. Shumyatsky, *Nonsoluble and non- p -soluble length of finite groups*, to appear in Israel J. Math.; [arXiv:1310.2434](https://arxiv.org/abs/1310.2434).
- [5] E. I. Khukhro and P. Shumyatsky, *Words and pronilpotent subgroups in profinite groups*, J. Aust. Math. Soc. **97** (2014), no. 3, 343–364, DOI 10.1017/S1446788714000317. MR3270773
- [6] Hans Kurzweil, *p -Automorphismen von auflösbaren p' -Gruppen* (German), Math. Z. **120** (1971), 326–354. MR0284503 (44 #1728)
- [7] John Thompson, *Finite groups with fixed-point-free automorphisms of prime order*, Proc. Nat. Acad. Sci. U.S.A. **45** (1959), 578–581. MR0104731 (21 #3484)
- [8] John G. Thompson, *Automorphisms of solvable groups*, J. Algebra **1** (1964), 259–267. MR0173710 (30 #3920)
- [9] Alexandre Turull, *Fitting height of groups and of fixed points*, J. Algebra **86** (1984), no. 2, 555–566, DOI 10.1016/0021-8693(84)90048-6. MR732266 (85i:20021)
- [10] Yan Ming Wang and Zhong Mu Chen, *Solubility of finite groups admitting a coprime order operator group* (English, with Italian summary), Boll. Un. Mat. Ital. A (7) **7** (1993), no. 3, 325–331. MR1249108 (95f:20038)
- [11] John S. Wilson, *On the structure of compact torsion groups*, Monatsh. Math. **96** (1983), no. 1, 57–66, DOI 10.1007/BF01298934. MR721596 (85a:22007)

SOBOLEV INSTITUTE OF MATHEMATICS, NOVOSIBIRSK, 630 090, RUSSIA – AND – UNIVERSITY OF LINCOLN, LINCOLN, UNITED KINGDOM.

E-mail address: khukhro@yahoo.co.uk

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRASILIA, DF 70910-900, BRAZIL

E-mail address: pavel@unb.br